

AE

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19988 A1

(51) International Patent Classification⁷: C12N 15/12, C07K 14/47, C12N 1/21, 5/10, C12Q 1/68, A61K 38/17

(74) Agents: CARROLL, Alice, O. et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02421 (US).

(21) International Application Number: PCT/US00/25135

(22) International Filing Date:
14 September 2000 (14.09.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/398,829 17 September 1999 (17.09.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02149 (US).

(72) Inventors: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). MCCOY, John, M.; 56 Howard Street, Reading, MA 01867 (US). LAVALLIE, Edward, R.; 113 Ann Lee Road, Harvard, MA 01451 (US). COLLINS-RACIE, Lisa, A.; 124 School Street, Acton, MA 01720 (US). EVANS, Cheryl; 18801 Bent Willow Circle, Germantown, MD 20874 (US). MERBERG, David; 2 Orchard Drive, Acton, MA 01720 (US). TREACY, Maurice; 12 Foxrock Court, Dublin 18 (IE). BOWMAN, Michael, R.; 63 Gloucester Road, Westwood, MA 02090 (US). SPAULDING, Vikki; 11 Meadowbank Road, Billerica, MA 01821 (US). AGOSTINO, Michael, J.; 26 Wolcott Avenue, Andover, MA 01810 (US).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- With (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/19988 A1

(54) Title: SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

(57) Abstract: Novel polynucleotides and the proteins encoded thereby are disclosed.

5 SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

BACKGROUND OF THE INVENTION

Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The
10 now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as
15 signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins and the polynucleotides
20 encoding them that the present invention is directed.

SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1
 from nucleotide 282 to nucleotide 565;
 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1
 from nucleotide 342 to nucleotide 565;
30 (d) a polynucleotide comprising the nucleotide sequence of the full-length
 protein coding sequence of clone AX65_22 deposited with the ATCC under accession
 number 98196;
 (e) a polynucleotide encoding the full-length protein encoded by the cDNA
 insert of clone AX65_22 deposited with the ATCC under accession number 98196;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AX65_22 deposited with the ATCC under accession number 98196;

5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AX65_22 deposited with the ATCC under accession number 98196;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;

10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:1.

20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 282 to nucleotide 565; the nucleotide sequence of SEQ ID NO:1 from nucleotide 342 to nucleotide 565; the nucleotide sequence of the full-length protein coding sequence of clone AX65_22 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone AX65_22 deposited with the ATCC under accession
25 number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AX65_22 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably
30 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:2.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
35 NO:1 and SEQ ID NO:3.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X

SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:1;

(ab) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(ac) the nucleotide sequence of the cDNA insert of clone AX65_22 deposited with the ATCC under accession number 98196;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:1;

(bb) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bc) the nucleotide sequence of the cDNA insert of clone AX65_22 deposited with the ATCC under accession number 98196;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:1 and SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:1 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:1 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:1. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 282 to nucleotide 565, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence
5 of SEQ ID NO:1 from nucleotide 282 to nucleotide 565, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 282 to nucleotide 565. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1 from nucleotide 342 to nucleotide 565, and extending contiguously from a nucleotide sequence corresponding to the 5'
10 end of said sequence of SEQ ID NO:1 from nucleotide 342 to nucleotide 565, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 342 to nucleotide 565.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:2;
- (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AX65_22 deposited with the ATCC under accession number 98196;

20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:2. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a protein comprising
25 a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:4;
- 30 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:4 from nucleotide 192 to nucleotide 2318;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:4 from nucleotide 653 to nucleotide 825;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD335_14 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD335_14 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:5;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:5 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:5;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:4.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:4 from nucleotide 192 to nucleotide 2318; the nucleotide sequence of SEQ ID NO:4 from nucleotide 653 to nucleotide 825; the nucleotide sequence of the full-length protein coding sequence of clone BD335_14 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BD335_14 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:5 from amino acid 148 to amino acid 240. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid

sequence of SEQ ID NO:5 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:5, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:5 having biological activity, the fragment comprising the amino acid sequence from amino
5 acid 349 to amino acid 358 of SEQ ID NO:5.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:4.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:4, but excluding the poly(A) tail at the 3'
15 end of SEQ ID NO:4; and
- (ab) the nucleotide sequence of the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
25 consisting of:
- (ba) SEQ ID NO:4, but excluding the poly(A) tail at the 3' end of SEQ ID NO:4; and
- (bb) the nucleotide sequence of the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196;
- 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
35 sequence corresponding to the cDNA sequence of SEQ ID NO:4, and extending contiguously from

a nucleotide sequence corresponding to the 5' end of SEQ ID NO:4 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:4, but excluding the poly(A) tail at the 3' end of SEQ ID NO:4. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:4 from nucleotide 192 to nucleotide 2318, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:4 from nucleotide 192 to nucleotide 2318, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:4 from nucleotide 192 to nucleotide 2318. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:4 from nucleotide 653 to nucleotide 825, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:4 from nucleotide 653 to nucleotide 825, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:4 from nucleotide 653 to nucleotide 825.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:5;
- (b) the amino acid sequence of SEQ ID NO:5 from amino acid 148 to amino acid 240;
- (c) a fragment of the amino acid sequence of SEQ ID NO:5, the fragment comprising eight contiguous amino acids of SEQ ID NO:5; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BD335_14 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:5 or the amino acid sequence of SEQ ID NO:5 from amino acid 148 to amino acid 240. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:5 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:5, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:5 having biological activity, the fragment comprising the amino acid sequence from amino acid 349 to amino acid 358 of SEQ ID NO:5.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 206 to nucleotide 391;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG241_1 deposited with the ATCC under accession number 98196;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG241_1 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG241_1 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG241_1 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:7.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 206 to nucleotide 391; the nucleotide sequence of the full-length protein coding sequence of clone BG241_1 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BG241_1 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG241_1 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide encoding a protein comprising a fragment of the amino acid

sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:8.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7, SEQ ID NO:6, and SEQ ID NO:9.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X
10 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:6;

(ab) SEQ ID NO:7;

(ac) SEQ ID NO:9, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:9; and

15 (ad) the nucleotide sequence of the cDNA insert of clone
BG241_1 deposited with the ATCC under accession number 98196;

(ii) hybridizing said probe(s) to human genomic DNA in conditions
at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

20 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in
6X SSC at 65 degrees C to a nucleotide sequence selected from the group
consisting of:

25 (ba) SEQ ID NO:6;

(bb) SEQ ID NO:7;

(bc) SEQ ID NO:9, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:9; and

(bd) the nucleotide sequence of the cDNA insert of clone
30 BG241_1 deposited with the ATCC under accession number 98196;

(ii) hybridizing said primer(s) to human genomic DNA in conditions
at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:6 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 206 to nucleotide 391, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 206 to nucleotide 391, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 206 to nucleotide 391.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BG241_1 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:10;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:10 from nucleotide 302 to nucleotide 1762;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:10 from nucleotide 389 to nucleotide 1762;

- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:10 from nucleotide 1723 to nucleotide 2050;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BL187_4 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BL187_4 deposited with the ATCC under accession number 98196;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:11;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:11 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:11;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:10.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:10 from nucleotide 302 to nucleotide 1762; the nucleotide sequence of SEQ ID NO:10 from nucleotide 389 to nucleotide 1762; the nucleotide sequence of SEQ ID NO:10 from nucleotide 1723 to nucleotide 2050; the nucleotide sequence of the full-length protein coding sequence of clone BL187_4 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BL187_4 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention

provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:11 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:11, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:11 having biological activity, the fragment comprising the amino acid sequence from amino acid 238 to amino acid 247 of SEQ ID NO:11.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:10.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:10, but excluding the poly(A) tail at the 3' end of SEQ ID NO:10; and

(ab) the nucleotide sequence of the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:10, but excluding the poly(A) tail at the 3' end of SEQ ID NO:10; and

(bb) the nucleotide sequence of the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:10, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:10 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:10, but excluding the poly(A) tail at the 3' end of SEQ ID NO:10. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:10 from nucleotide 302 to nucleotide 1762, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:10 from nucleotide 302 to nucleotide 1762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:10 from nucleotide 302 to nucleotide 1762. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:10 from nucleotide 389 to nucleotide 1762, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:10 from nucleotide 389 to nucleotide 1762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:10 from nucleotide 389 to nucleotide 1762. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:10 from nucleotide 1723 to nucleotide 2050, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:10 from nucleotide 1723 to nucleotide 2050, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:10 from nucleotide 1723 to nucleotide 2050.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:11;
- (b) a fragment of the amino acid sequence of SEQ ID NO:11, the fragment comprising eight contiguous amino acids of SEQ ID NO:11; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BL187_4 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:11. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:11 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:11, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:11 having biological activity, the fragment comprising the amino acid sequence from amino acid 238 to amino acid 247 of SEQ ID NO:11.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:12;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:12
5 from nucleotide 2 to nucleotide 2290;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:12
from nucleotide 134 to nucleotide 2290;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:12
from nucleotide 1 to nucleotide 309;
- 10 (e) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone BL249_18 deposited with the ATCC under accession
number 98196;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone BL249_18 deposited with the ATCC under accession number 98196;
- 15 (g) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone BL249_18 deposited with the ATCC under accession number
98196;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone BL249_18 deposited with the ATCC under accession number 98196;
- 20 (i) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:13;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:13 having biological activity, the fragment comprising eight
contiguous amino acids of SEQ ID NO:13;
- 25 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i)
or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the
length of SEQ ID NO:12.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:12
35 from nucleotide 2 to nucleotide 2290; the nucleotide sequence of SEQ ID NO:12 from nucleotide

134 to nucleotide 2290; the nucleotide sequence of SEQ ID NO:12 from nucleotide 1 to nucleotide 309; the nucleotide sequence of the full-length protein coding sequence of clone BL249_18 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BL249_18 deposited with the ATCC under
5 accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BL249_18 deposited with the ATCC under accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:13 from amino acid 3 to amino acid 102. In further preferred embodiments, the present
10 invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:13 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:13, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:13 having biological activity, the fragment comprising the amino acid sequence from amino
15 acid 376 to amino acid 385 of SEQ ID NO:13.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:12.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 25 (aa) SEQ ID NO:12, but excluding the poly(A) tail at the 3' end of SEQ ID NO:12; and
- (ab) the nucleotide sequence of the cDNA insert of clone BL249_18 deposited with the ATCC under accession number 98196;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
35 consisting of:

-16-

(ba) SEQ ID NO:12, but excluding the poly(A) tail at the 3' end of SEQ ID NO:12; and

(bb) the nucleotide sequence of the cDNA insert of clone BL249_18 deposited with the ATCC under accession number 98196;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
10 sequence corresponding to the cDNA sequence of SEQ ID NO:12, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:12 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:12, but excluding the poly(A) tail at the 3' end of SEQ ID NO:12. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:12 from nucleotide
15 2 to nucleotide 2290, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:12 from nucleotide 2 to nucleotide 2290, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:12 from nucleotide 2 to nucleotide 2290. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID
20 NO:12 from nucleotide 134 to nucleotide 2290, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:12 from nucleotide 134 to nucleotide 2290, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:12 from nucleotide 134 to nucleotide 2290. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
25 sequence of SEQ ID NO:12 from nucleotide 1 to nucleotide 309, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:12 from nucleotide 1 to nucleotide 309, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:12 from nucleotide 1 to nucleotide 309.

In other embodiments, the present invention provides a composition comprising a protein,
30 wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:13;

(b) the amino acid sequence of SEQ ID NO:13 from amino acid 3 to amino acid 102;

(c) a fragment of the amino acid sequence of SEQ ID NO:13, the fragment
35 comprising eight contiguous amino acids of SEQ ID NO:13; and

- (d) the amino acid sequence encoded by the cDNA insert of clone BL249_18 deposited with the ATCC under accession number 98196;
the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:13 or the amino acid sequence of SEQ ID NO:13 from amino acid 3 to amino acid 102. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:13 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:13, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:13 having biological activity, the fragment comprising the amino acid sequence from amino acid 376 to amino acid 385 of SEQ ID NO:13.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15;
(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 459 to nucleotide 539;
(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BO71_1 deposited with the ATCC under accession number 98196;
(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196;
(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BO71_1 deposited with the ATCC under accession number 98196;
(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196;
(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:15.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 459 to nucleotide 539; the nucleotide sequence of the full-length protein coding sequence of clone BO71_1 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BO71_1 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 8 to amino acid 17 of SEQ ID NO:16.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15, SEQ ID NO:14, and SEQ ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

25 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:14;

(ab) SEQ ID NO:15;

30 (ac) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and

(ad) the nucleotide sequence of the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

35 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:14;

(bb) SEQ ID NO:15;

(bc) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and

(bd) the nucleotide sequence of the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:14 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 459 to nucleotide 539, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 459 to nucleotide 539, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide 459 to nucleotide 539.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:16;

(b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and

(c) the amino acid sequence encoded by the cDNA insert of clone BO71_1 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 8 to amino acid 17 of SEQ ID NO:16.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:18;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:18 from nucleotide 1237 to nucleotide 1944;

15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:18 from nucleotide 737 to nucleotide 1072;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BO365_2 deposited with the ATCC under accession number 98196;

20 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BO365_2 deposited with the ATCC under accession number 98196;

25 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:19;

30 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:19;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

35 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:18.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:18 from nucleotide 1237 to nucleotide 1944; the nucleotide sequence of SEQ ID NO:18 from nucleotide 737 to nucleotide 1072; the nucleotide sequence of the full-length protein coding sequence of clone BO365_2 deposited with the ATCC under accession number 98196; or the
10 nucleotide sequence of a mature protein coding sequence of clone BO365_2 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the
15 amino acid sequence of SEQ ID NO:19 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:19, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment comprising the amino acid sequence from amino acid 113 to amino acid 122 of SEQ ID NO:19.

20 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:18.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- 25 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:18, but excluding the poly(A) tail at the 3' end of SEQ ID NO:18; and
- 30 (ab) the nucleotide sequence of the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- 35 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:18, but excluding the poly(A) tail at the 3' end of SEQ ID NO:18; and

(bb) the nucleotide sequence of the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:18, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:18 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:18, but excluding the poly(A) tail at the 3' end of SEQ ID NO:18. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:18 from nucleotide 1237 to nucleotide 1944, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:18 from nucleotide 1237 to nucleotide 1944, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:18 from nucleotide 1237 to nucleotide 1944. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:18 from nucleotide 737 to nucleotide 1072, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:18 from nucleotide 737 to nucleotide 1072, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:18 from nucleotide 737 to nucleotide 1072.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:19;

(b) a fragment of the amino acid sequence of SEQ ID NO:19, the fragment comprising eight contiguous amino acids of SEQ ID NO:19; and

(c) the amino acid sequence encoded by the cDNA insert of clone BO365_2 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:19. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment preferably comprising eight (more preferably
 5 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:19, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment comprising the amino acid sequence from amino acid 113 to amino acid 122 of SEQ ID NO:19.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:20;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:20 from nucleotide 68 to nucleotide 328;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BV51_1 deposited with the ATCC under accession
 15 number 98196;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BV51_1 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV51_1 deposited with the ATCC under accession number
 20 98196;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BV51_1 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:21;
- 25 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:21 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:21;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 30 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:20.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:20
5 from nucleotide 68 to nucleotide 328; the nucleotide sequence of the full-length protein coding sequence of clone BV51_1 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BV51_1 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV51_1
10 deposited with the ATCC under accession number 98196. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:21 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:21, or a polynucleotide encoding a protein comprising a fragment of the amino acid
15 sequence of SEQ ID NO:21 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:21.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:20 and SEQ ID NO:22.

Further embodiments of the invention provide isolated polynucleotides produced
20 according to a process selected from the group consisting of:

(a) a process comprising the steps of:
(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (aa) SEQ ID NO:20;
(ab) SEQ ID NO:22, but excluding the poly(A) tail at the 3' end of SEQ ID NO:22; and
(ac) the nucleotide sequence of the cDNA insert of clone BV51_1 deposited with the ATCC under accession number 98196;
30 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:20;

5 (bb) SEQ ID NO:22, but excluding the poly(A) tail at the 3' end of SEQ ID NO:22; and

(bc) the nucleotide sequence of the cDNA insert of clone BV51_1 deposited with the ATCC under accession number 98196;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:20 and SEQ ID NO:22, and
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:20 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:22, but excluding the poly(A) tail at the 3' end of SEQ ID NO:22. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:20, and extending contiguously from a nucleotide sequence corresponding to the 5' end of
20 SEQ ID NO:20 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:20. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:20 from nucleotide 68 to nucleotide 328, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:20 from nucleotide 68 to nucleotide 328, to a nucleotide
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:20 from nucleotide 68 to nucleotide 328.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:21;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:21, the fragment comprising eight contiguous amino acids of SEQ ID NO:21; and

(c) the amino acid sequence encoded by the cDNA insert of clone BV51_1 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein
35 comprises the amino acid sequence of SEQ ID NO:21. In further preferred embodiments, the

present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:21 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:21, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:21 having biological activity, the fragment
5 comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:21.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:24;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:24
10 from nucleotide 57 to nucleotide 396;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BV140_3 deposited with the ATCC under accession number 98196;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA
15 insert of clone BV140_3 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV140_3 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert
20 of clone BV140_3 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:25;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:25 having biological activity, the fragment comprising eight
25 contiguous amino acids of SEQ ID NO:25;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:24.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:24 from nucleotide 57 to nucleotide 396; the nucleotide sequence of the full-length protein coding sequence of clone BV140_3 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BV140_3 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV140_3 deposited with the ATCC under accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:25 from amino acid 29 to amino acid 57. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:25 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:25, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:25 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:25.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:24, SEQ ID NO:23, and SEQ ID NO:26.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:23;
 - (ab) SEQ ID NO:24;
 - (ac) SEQ ID NO:26, but excluding the poly(A) tail at the 3' end of SEQ ID NO:26; and
 - (ad) the nucleotide sequence of the cDNA insert of clone BV140_3 deposited with the ATCC under accession number 98196;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:23;
- 5 (bb) SEQ ID NO:24;
- (bc) SEQ ID NO:26, but excluding the poly(A) tail at the 3' end of SEQ ID NO:26; and
- (bd) the nucleotide sequence of the cDNA insert of clone BV140_3 deposited with the ATCC under accession number 98196;
- 10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
15 sequence corresponding to the cDNA sequences of SEQ ID NO:23, SEQ ID NO:24, and SEQ ID NO:26, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:23 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:26, but excluding the poly(A) tail at the 3' end of SEQ ID NO:26. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the
20 cDNA sequence of SEQ ID NO:24, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:24 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:24. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:24 from nucleotide 57 to nucleotide 396, and extending contiguously from a nucleotide sequence
25 corresponding to the 5' end of said sequence of SEQ ID NO:24 from nucleotide 57 to nucleotide 396, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:24 from nucleotide 57 to nucleotide 396.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:25;
- (b) the amino acid sequence of SEQ ID NO:25 from amino acid 29 to amino acid 57;
- (c) a fragment of the amino acid sequence of SEQ ID NO:25, the fragment comprising eight contiguous amino acids of SEQ ID NO:25; and

(d) the amino acid sequence encoded by the cDNA insert of clone BV140_3 deposited with the ATCC under accession number 98196;
the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:25 or the amino acid sequence of SEQ ID NO:25 from amino acid 29 to amino acid 57. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:25 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:25, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:25 having biological activity, the fragment comprising the amino acid sequence from amino acid 51 to amino acid 60 of SEQ ID NO:25.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 328;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 1 to nucleotide 197;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BV141_2 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BV141_2 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV141_2 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BV141_2 deposited with the ATCC under accession number 98196;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

5 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:27.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 328; the nucleotide sequence of SEQ ID NO:27 from
10 nucleotide 1 to nucleotide 197; the nucleotide sequence of the full-length protein coding sequence of clone BV141_2 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone BV141_2 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV141_2 deposited with the
15 ATCC under accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28 from amino acid 1 to amino acid 37. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight
20 (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:28.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
25 NO:27.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X
30 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

(ab) the nucleotide sequence of the cDNA insert of clone
35 BV141_2 deposited with the ATCC under accession number 98196;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

5 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

(bb) the nucleotide sequence of the cDNA insert of clone BV141_2 deposited with the ATCC under accession number 98196;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:27 to a nucleotide sequence
20 corresponding to the 3' end of SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 328, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 328, to a
25 nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 101 to nucleotide 328. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 1 to nucleotide 197, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 1 to
30 nucleotide 197, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 1 to nucleotide 197.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:28;

(b) the amino acid sequence of SEQ ID NO:28 from amino acid 1 to amino acid 37;

(c) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

5 (d) the amino acid sequence encoded by the cDNA insert of clone BV141_2 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28 or the amino acid sequence of SEQ ID NO:28 from amino acid 1 to amino acid 37. In further preferred embodiments, the present
10 invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 33 to amino acid 42 of SEQ ID NO:28.

15 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 28 to nucleotide 351;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 328 to nucleotide 351;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CC194_4 deposited with the ATCC under accession number 98196;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CC194_4 deposited with the ATCC under accession number 98196;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CC194_4 deposited with the ATCC under accession number 98196;

30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CC194_4 deposited with the ATCC under accession number 98196;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:30;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:29.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:29 from nucleotide 28 to nucleotide 351; the nucleotide sequence of SEQ ID NO:29 from nucleotide 15 328 to nucleotide 351; the nucleotide sequence of the full-length protein coding sequence of clone CC194_4 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone CC194_4 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CC194_4 deposited with the ATCC under 20 accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:30 from amino acid 56 to amino acid 108. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more 25 preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 49 to amino acid 58 of SEQ ID NO:30.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID 30 NO:29 and SEQ ID NO:31.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:29;
 - 5 (ab) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
 - (ac) the nucleotide sequence of the cDNA insert of clone CC194_4 deposited with the ATCC under accession number 98196;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions
- 10 at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
 - 15 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:29;
 - (bb) SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31; and
 - 20 (bc) the nucleotide sequence of the cDNA insert of clone CC194_4 deposited with the ATCC under accession number 98196;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - 25 (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:29 and SEQ ID NO:31, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:31, but excluding the poly(A)
- 30 tail at the 3' end of SEQ ID NO:31. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:29. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide
- 35 sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 28 to

nucleotide 351, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 28 to nucleotide 351, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 28 to nucleotide 351. Also preferably the polynucleotide isolated according to the above process
5 comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 328 to nucleotide 351, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 328 to nucleotide 351, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 328 to nucleotide 351.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
- (b) the amino acid sequence of SEQ ID NO:30 from amino acid 56 to amino acid 108;
- 15 (c) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CC194_4 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein
20 comprises the amino acid sequence of SEQ ID NO:30 or the amino acid sequence of SEQ ID NO:30 from amino acid 56 to amino acid 108. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a protein comprising a
25 fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 49 to amino acid 58 of SEQ ID NO:30.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:32;
- 30 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:32 from nucleotide 338 to nucleotide 1198;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:32 from nucleotide 467 to nucleotide 1058;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone DA136_11 deposited with the ATCC under accession number 98196;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone DA136_11 deposited with the ATCC under accession number 98196;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:33;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:33 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:33;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:32.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:32 from nucleotide 338 to nucleotide 1198; the nucleotide sequence of SEQ ID NO:32 from nucleotide 467 to nucleotide 1058; the nucleotide sequence of the full-length protein coding sequence of clone DA136_11 deposited with the ATCC under accession number 98196; or the nucleotide sequence of a mature protein coding sequence of clone DA136_11 deposited with the ATCC under accession number 98196. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:33 from amino acid 124 to amino acid 182. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a

fragment of the amino acid sequence of SEQ ID NO:33 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:33, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:33 having biological activity, the fragment comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:33.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:32.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 15 (aa) SEQ ID NO:32, but excluding the poly(A) tail at the 3' end of SEQ ID NO:32; and
 - (ab) the nucleotide sequence of the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 25 (ba) SEQ ID NO:32, but excluding the poly(A) tail at the 3' end of SEQ ID NO:32; and
 - (bb) the nucleotide sequence of the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196;
 - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:32, and extending contiguously

from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:32 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:32, but excluding the poly(A) tail at the 3' end of SEQ ID NO:32. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:32 from nucleotide
5 338 to nucleotide 1198, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:32 from nucleotide 338 to nucleotide 1198, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:32 from nucleotide 338 to nucleotide 1198. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID
10 NO:32 from nucleotide 467 to nucleotide 1058, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:32 from nucleotide 467 to nucleotide 1058, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:32 from nucleotide 467 to nucleotide 1058.

In other embodiments, the present invention provides a composition comprising a protein,
15 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:33;
- (b) the amino acid sequence of SEQ ID NO:33 from amino acid 124 to amino acid 182;
- (c) a fragment of the amino acid sequence of SEQ ID NO:33, the fragment
20 comprising eight contiguous amino acids of SEQ ID NO:33; and
- (d) the amino acid sequence encoded by the cDNA insert of clone DA136_11 deposited with the ATCC under accession number 98196;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:33 or the amino acid sequence of SEQ ID
25 NO:33 from amino acid 124 to amino acid 182. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:33 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:33, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:33 having biological activity, the fragment
30 comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:33.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:34;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:34 from nucleotide 437 to nucleotide 1159;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:34 from nucleotide 515 to nucleotide 1159;
- 5 (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:34 from nucleotide 539 to nucleotide 1099;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AR415_4 deposited with the ATCC under accession number 98232;
- 10 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AR415_4 deposited with the ATCC under accession number 98232;
- 15 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:35;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:35 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:35;
- 20 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:34.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:34 from nucleotide 437 to nucleotide 1159; the nucleotide sequence of SEQ ID NO:34 from nucleotide 515 to nucleotide 1159; the nucleotide sequence of SEQ ID NO:34 from nucleotide 539 to nucleotide 1099; the nucleotide sequence of the full-length protein coding sequence of clone AR415_4 deposited with the ATCC under accession number 98232; or the nucleotide

35

sequence of a mature protein coding sequence of clone AR415_4 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:35 from amino acid 51 to amino acid 221. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:35 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:35, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:35 having biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:35.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:34.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:34, but excluding the poly(A) tail at the 3' end of SEQ ID NO:34; and

(ab) the nucleotide sequence of the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:34, but excluding the poly(A) tail at the 3' end of SEQ ID NO:34; and

(bb) the nucleotide sequence of the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

5 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:34, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:34 to a nucleotide sequence
10 corresponding to the 3' end of SEQ ID NO:34, but excluding the poly(A) tail at the 3' end of SEQ ID NO:34. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:34 from nucleotide 437 to nucleotide 1159, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:34 from nucleotide 437 to nucleotide 1159, to a
15 nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:34 from nucleotide 437 to nucleotide 1159. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:34 from nucleotide 515 to nucleotide 1159, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:34 from nucleotide 515 to
20 nucleotide 1159, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:34 from nucleotide 515 to nucleotide 1159. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:34 from nucleotide 539 to nucleotide 1099, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:34 from
25 nucleotide 539 to nucleotide 1099, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:34 from nucleotide 539 to nucleotide 1099.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:35;
- (b) the amino acid sequence of SEQ ID NO:35 from amino acid 51 to amino acid 221;
- (c) a fragment of the amino acid sequence of SEQ ID NO:35, the fragment comprising eight contiguous amino acids of SEQ ID NO:35; and
- 35 (d) the amino acid sequence encoded by the cDNA insert of clone AR415_4 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:35 or the amino acid sequence of SEQ ID NO:35 from amino acid 51 to amino acid 221. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:35
5 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:35, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:35 having biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:35.

In one embodiment, the present invention provides a composition comprising an isolated
10 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:36;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:36 from nucleotide 59 to nucleotide 376;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:36
15 from nucleotide 179 to nucleotide 376;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AS63_29 deposited with the ATCC under accession number 98232;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
20 insert of clone AS63_29 deposited with the ATCC under accession number 98232;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AS63_29 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
25 of clone AS63_29 deposited with the ATCC under accession number 98232;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:37;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment comprising eight
30 contiguous amino acids of SEQ ID NO:37;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:36.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:36 from nucleotide 59 to nucleotide 376; the nucleotide sequence of SEQ ID NO:36 from nucleotide 179 to nucleotide 376; the nucleotide sequence of the full-length protein coding sequence of clone AS63_29 deposited with the ATCC under accession number 98232; or the nucleotide sequence
10 of a mature protein coding sequence of clone AS63_29 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AS63_29 deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:37 from
15 amino acid 1 to amino acid 91. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:37, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having
20 biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:37.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:36 and SEQ ID NO:38.

Further embodiments of the invention provide isolated polynucleotides produced
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:36;
(ab) SEQ ID NO:38, but excluding the poly(A) tail at the 3' end of SEQ ID NO:38; and
(ac) the nucleotide sequence of the cDNA insert of clone AS63_29 deposited with the ATCC under accession number 98232;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

5 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:36;

10 (bb) SEQ ID NO:38, but excluding the poly(A) tail at the 3' end of SEQ ID NO:38; and

(bc) the nucleotide sequence of the cDNA insert of clone AS63_29 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:36 and SEQ ID NO:38, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:36 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:38, but excluding the poly(A) tail at the 3' end of SEQ ID NO:38. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:36, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:36 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:36. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:36 from nucleotide 59 to nucleotide 376, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:36 from nucleotide 59 to nucleotide 376, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:36 from nucleotide 59 to nucleotide 376. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:36 from nucleotide 179 to nucleotide 376, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:36 from nucleotide 179 to nucleotide

376, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:36 from nucleotide 179 to nucleotide 376.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 5 (a) the amino acid sequence of SEQ ID NO:37;
- (b) the amino acid sequence of SEQ ID NO:37 from amino acid 1 to amino acid 91;
- (c) a fragment of the amino acid sequence of SEQ ID NO:37, the fragment comprising eight contiguous amino acids of SEQ ID NO:37; and
- 10 (d) the amino acid sequence encoded by the cDNA insert of clone AS63_29 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:37 or the amino acid sequence of SEQ ID NO:37 from amino acid 1 to amino acid 91. In further preferred embodiments, the present
15 invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:37, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:37.

20 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 198 to nucleotide 2039;
- 25 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 490 to nucleotide 809;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AY304_14 deposited with the ATCC under accession number 98561;
- 30 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AY304_14 deposited with the ATCC under accession number 98561;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AY304_14 deposited with the ATCC under accession number 98561;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AY304_14 deposited with the ATCC under accession number 98561;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
15 the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:39.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 198 to nucleotide 2039; the nucleotide sequence of SEQ ID NO:39 from nucleotide 490 to nucleotide 809; the nucleotide sequence of the full-length protein coding
20 sequence of clone AY304_14 deposited with the ATCC under accession number 98561; or the nucleotide sequence of a mature protein coding sequence of clone AY304_14 deposited with the ATCC under accession number 98561. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AY304_14 deposited with the ATCC under accession number 98561. In yet other preferred embodiments,
25 the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40 from amino acid 106 to amino acid 204. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino
30 acids of SEQ ID NO:40, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 302 to amino acid 311 of SEQ ID NO:40.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:39.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X

5 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and

10 (ab) the nucleotide sequence of the cDNA insert of clone AY304_14 deposited with the ATCC under accession number 98561;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

15 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and

(bb) the nucleotide sequence of the cDNA insert of clone AY304_14 deposited with the ATCC under accession number 98561;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:39 to a nucleotide sequence
30 corresponding to the 3' end of SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 198 to nucleotide 2039, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 198 to nucleotide 2039, to a
35 nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from

nucleotide 198 to nucleotide 2039. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 490 to nucleotide 809, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 490 to nucleotide 809, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 490 to nucleotide 809.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
- 10 (b) the amino acid sequence of SEQ ID NO:40 from amino acid 106 to amino acid 204;
- (c) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AY304_14
- 15 deposited with the ATCC under accession number 98561;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40 or the amino acid sequence of SEQ ID NO:40 from amino acid 106 to amino acid 204. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40

20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 302 to amino acid 311 of SEQ ID NO:40.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 2027;
- 30 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 2027;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- 15 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:41.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 2027; the nucleotide sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 2027; the nucleotide sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431; the nucleotide sequence of the full-length protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232; or the nucleotide sequence
- 30 of a mature protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42 from
- 35 amino acid 1 to amino acid 110. In further preferred embodiments, the present invention provides

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having
5 biological activity, the fragment comprising the amino acid sequence from amino acid 316 to amino acid 325 of SEQ ID NO:42.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:41.

Further embodiments of the invention provide isolated polynucleotides produced
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and

(ab) the nucleotide sequence of the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41; and

30 (bb) the nucleotide sequence of the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:41 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 207, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 207, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 207. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 207, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 207, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 207. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
 - (b) the amino acid sequence of SEQ ID NO:42 from amino acid 1 to amino acid 110;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:42 or the amino acid sequence of SEQ ID NO:42 from amino acid 1 to amino acid 110. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a protein comprising a

fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 316 to amino acid 325 of SEQ ID NO:42.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:44;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:44 from nucleotide 566 to nucleotide 631;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BO432_4 deposited with the ATCC under accession
10 number 98232;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BO432_4 deposited with the ATCC under accession number
15 98232;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:45;
- 20 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:45 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:45;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 25 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:44.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:44 from nucleotide 566 to nucleotide 631; the nucleotide sequence of the full-length protein coding sequence of clone BO432_4 deposited with the ATCC under accession number 98232; or the
35 nucleotide sequence of a mature protein coding sequence of clone BO432_4 deposited with the

ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:45 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:45, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:45 having biological activity, the fragment comprising the amino acid sequence from amino acid 6 to amino acid 15 of SEQ ID NO:45.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:44, SEQ ID NO:43, and SEQ ID NO:46.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:43;
 - (ab) SEQ ID NO:44;
 - (ac) SEQ ID NO:46, but excluding the poly(A) tail at the 3' end of SEQ ID NO:46; and
 - (ad) the nucleotide sequence of the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:43;
 - (bb) SEQ ID NO:44;
 - (bc) SEQ ID NO:46, but excluding the poly(A) tail at the 3' end of SEQ ID NO:46; and

- (bd) the nucleotide sequence of the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:46, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:43 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:46, but excluding the poly(A) tail at the 3' end of SEQ ID NO:46. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:44, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:44 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:44. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:44 from nucleotide 566 to nucleotide 631, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:44 from nucleotide 566 to nucleotide 631, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:44 from nucleotide 566 to nucleotide 631.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:45;
- (b) a fragment of the amino acid sequence of SEQ ID NO:45, the fragment comprising eight contiguous amino acids of SEQ ID NO:45; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone BO432_4 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:45. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:45 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:45, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:45 having biological activity, the fragment comprising the amino acid sequence from amino acid 6 to amino acid 15 of SEQ ID NO:45.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47
5 from nucleotide 45 to nucleotide 428;
 - (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BO538_2 deposited with the ATCC under accession number 98232;
 - (d) a polynucleotide encoding the full-length protein encoded by the cDNA
10 insert of clone BO538_2 deposited with the ATCC under accession number 98232;
 - (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BO538_2 deposited with the ATCC under accession number 98232;
 - (f) a polynucleotide encoding a mature protein encoded by the cDNA insert
15 of clone BO538_2 deposited with the ATCC under accession number 98232;
 - (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
 - (h) a polynucleotide encoding a protein comprising a fragment of the amino
20 acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
 - (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
 - (j) a polynucleotide which encodes a species homologue of the protein of (g)
or (h) above ;
 - (k) a polynucleotide that hybridizes under stringent conditions to any one of
25 the polynucleotides specified in (a)-(h); and
 - (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:47.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 45 to nucleotide 428; the nucleotide sequence of the full-length protein coding sequence of clone BO538_2 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone BO538_2 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide
35 encodes the full-length or a mature protein encoded by the cDNA insert of clone BO538_2

deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48 from amino acid 52 to amino acid 128. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a
5 fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 59 to amino acid 68 of SEQ ID NO:48.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:47 and SEQ ID NO:49.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:47;
 - (ab) SEQ ID NO:49, but excluding the poly(A) tail at the 3'
20 end of SEQ ID NO:49; and
 - (ac) the nucleotide sequence of the cDNA insert of clone BO538_2 deposited with the ATCC under accession number 98232;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
30 consisting of:
 - (ba) SEQ ID NO:47;
 - (bb) SEQ ID NO:49, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:49; and
 - (bc) the nucleotide sequence of the cDNA insert of clone
35 BO538_2 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:47 and SEQ ID NO:49, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49. Also preferably the polynucleotide isolated according to the
10 above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:47. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 45 to
15 nucleotide 428, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 45 to nucleotide 428, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 45 to nucleotide 428.

In other embodiments, the present invention provides a composition comprising a protein,
20 wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:48;

(b) the amino acid sequence of SEQ ID NO:48 from amino acid 52 to amino acid 128;

(c) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment
25 comprising eight contiguous amino acids of SEQ ID NO:48; and

(d) the amino acid sequence encoded by the cDNA insert of clone BO538_2 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48 or the amino acid sequence of SEQ ID
30 NO:48 from amino acid 52 to amino acid 128. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment
35 comprising the amino acid sequence from amino acid 59 to amino acid 68 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:50;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:50
5 from nucleotide 144 to nucleotide 566;
 - (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BR595_4 deposited with the ATCC under accession number 98232;
 - (d) a polynucleotide encoding the full-length protein encoded by the cDNA
10 insert of clone BR595_4 deposited with the ATCC under accession number 98232;
 - (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BR595_4 deposited with the ATCC under accession number 98232;
 - (f) a polynucleotide encoding a mature protein encoded by the cDNA insert
15 of clone BR595_4 deposited with the ATCC under accession number 98232;
 - (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:51;
 - (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment comprising eight
20 contiguous amino acids of SEQ ID NO:51;
 - (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
 - (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
 - (k) a polynucleotide that hybridizes under stringent conditions to any one of
25 the polynucleotides specified in (a)-(h); and
 - (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:50.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:50 from nucleotide 144 to nucleotide 566; the nucleotide sequence of the full-length protein coding sequence of clone BR595_4 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone BR595_4 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide
35 encodes the full-length or a mature protein encoded by the cDNA insert of clone BR595_4

deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:51 from amino acid 39 to amino acid 141. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a
5 fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:51, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:51.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:50 and SEQ ID NO:52.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:50;
 - (ab) SEQ ID NO:52, but excluding the poly(A) tail at the 3'
20 end of SEQ ID NO:52; and
 - (ac) the nucleotide sequence of the cDNA insert of clone BR595_4 deposited with the ATCC under accession number 98232;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in
30 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:50;
 - (bb) SEQ ID NO:52, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:52; and
 - (bc) the nucleotide sequence of the cDNA insert of clone
35 BR595_4 deposited with the ATCC under accession number 98232;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:50 and SEQ ID NO:52, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:50 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:52, but excluding the poly(A) tail at the 3' end of SEQ ID NO:52. Also preferably the polynucleotide isolated according to the
- 10 above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:50, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:50 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:50. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:50 from nucleotide 144 to
- 15 nucleotide 566, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:50 from nucleotide 144 to nucleotide 566, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:50 from nucleotide 144 to nucleotide 566.

- In other embodiments, the present invention provides a composition comprising a protein,
- 20 wherein said protein comprises an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:51;
 - (b) the amino acid sequence of SEQ ID NO:51 from amino acid 39 to amino acid 141;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:51, the fragment
 - 25 comprising eight contiguous amino acids of SEQ ID NO:51; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BR595_4 deposited with the ATCC under accession number 98232;

- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:51 or the amino acid sequence of SEQ ID
- 30 NO:51 from amino acid 39 to amino acid 141. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:51, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment
- 35 comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:51.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53
5 from nucleotide 232 to nucleotide 1041;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53
from nucleotide 460 to nucleotide 1041;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53
from nucleotide 590 to nucleotide 1163;
- 10 (e) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone CI490_2 deposited with the ATCC under accession
number 98232;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone CI490_2 deposited with the ATCC under accession number 98232;
- 15 (g) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone CI490_2 deposited with the ATCC under accession number
98232;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone CI490_2 deposited with the ATCC under accession number 98232;
- 20 (i) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:54;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising eight
contiguous amino acids of SEQ ID NO:54;
- 25 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i)
or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j); and
- 30 (n) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the
length of SEQ ID NO:53.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:53
35 from nucleotide 232 to nucleotide 1041; the nucleotide sequence of SEQ ID NO:53 from

nucleotide 460 to nucleotide 1041; the nucleotide sequence of SEQ ID NO:53 from nucleotide 590 to nucleotide 1163; the nucleotide sequence of the full-length protein coding sequence of clone CI490_2 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone CI490_2 deposited with the ATCC under
5 accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CI490_2 deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:54 from amino acid 133 to amino acid 270. In further preferred embodiments, the present
10 invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino
15 acid 130 to amino acid 139 of SEQ ID NO:54.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:53.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:53, but excluding the poly(A) tail at the 3'
 - 25 end of SEQ ID NO:53; and
 - (ab) the nucleotide sequence of the cDNA insert of clone CI490_2 deposited with the ATCC under accession number 98232;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
35 consisting of:

(ba) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and

(bb) the nucleotide sequence of the cDNA insert of clone CI490_2 deposited with the ATCC under accession number 98232;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:53 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide
15 232 to nucleotide 1041, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 232 to nucleotide 1041, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 232 to nucleotide 1041. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID
20 NO:53 from nucleotide 460 to nucleotide 1041, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 460 to nucleotide 1041, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 460 to nucleotide 1041. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
25 sequence of SEQ ID NO:53 from nucleotide 590 to nucleotide 1163, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 590 to nucleotide 1163, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 590 to nucleotide 1163.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) the amino acid sequence of SEQ ID NO:54 from amino acid 133 to amino acid 270;

35 (c) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(d) the amino acid sequence encoded by the cDNA insert of clone CI490_2 deposited with the ATCC under accession number 98232;
the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:54 or the amino acid sequence of SEQ ID NO:54 from amino acid 133 to amino acid 270. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 130 to amino acid 139 of SEQ ID NO:54.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 268 to nucleotide 624;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 624;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CI522_1 deposited with the ATCC under accession number 98232;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CI522_1 deposited with the ATCC under accession number 98232;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CI522_1 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CI522_1 deposited with the ATCC under accession number 98232;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

5 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:55.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 268 to nucleotide 624; the nucleotide sequence of SEQ ID NO:55 from nucleotide
10 325 to nucleotide 624; the nucleotide sequence of the full-length protein coding sequence of clone CI522_1 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone CI522_1 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CI522_1 deposited with the ATCC under
15 accession number 98232. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a polynucleotide
20 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:56.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:55 and SEQ ID NO:57.

Further embodiments of the invention provide isolated polynucleotides produced
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:55;

(ab) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(ac) the nucleotide sequence of the cDNA insert of clone CI522_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

5 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:55;

10 (bb) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and

(bc) the nucleotide sequence of the cDNA insert of clone CI522_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:55 and SEQ ID NO:57, and
20 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55, and extending contiguously from a nucleotide sequence corresponding to the 5' end of
25 SEQ ID NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:55. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 268 to nucleotide 624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 268 to nucleotide 624, to a nucleotide
30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 268 to nucleotide 624. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 624, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide

624, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide 325 to nucleotide 624.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 5 (a) the amino acid sequence of SEQ ID NO:56;
- (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
- (c) the amino acid sequence encoded by the cDNA insert of clone CI522_1 deposited with the ATCC under accession number 98232;
- 10 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a protein comprising
- 15 a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:56.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:58;
- 20 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:58 from nucleotide 288 to nucleotide 710;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:58 from nucleotide 868 to nucleotide 1887;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CN238_1 deposited with the ATCC under accession
- 25 number 98232;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CN238_1 deposited with the ATCC under accession number 98232;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein
- 30 coding sequence of clone CN238_1 deposited with the ATCC under accession number 98232;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CN238_1 deposited with the ATCC under accession number 98232;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence
- 35 of SEQ ID NO:59;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:59;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:58.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:58 from nucleotide 288 to nucleotide 710; the nucleotide sequence of SEQ ID NO:58 from
15 nucleotide 868 to nucleotide 1887; the nucleotide sequence of the full-length protein coding sequence of clone CN238_1 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone CN238_1 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CN238_1
20 deposited with the ATCC under accession number 98232. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:59 from amino acid 1 to amino acid 109. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment
25 preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:59, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:59.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
30 NO:58.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:58, but excluding the poly(A) tail at the 3' end of SEQ ID NO:58; and

(ab) the nucleotide sequence of the cDNA insert of clone CN238_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:58, but excluding the poly(A) tail at the 3' end of SEQ ID NO:58; and

(bb) the nucleotide sequence of the cDNA insert of clone CN238_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:58, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:58 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:58, but excluding the poly(A) tail at the 3' end of SEQ ID NO:58. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:58 from nucleotide 288 to nucleotide 710, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:58 from nucleotide 288 to nucleotide 710, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:58 from nucleotide 288 to nucleotide 710. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:58 from nucleotide 868 to nucleotide 1887, and extending contiguously from a nucleotide

sequence corresponding to the 5' end of said sequence of SEQ ID NO:58 from nucleotide 868 to nucleotide 1887, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:58 from nucleotide 868 to nucleotide 1887.

In other embodiments, the present invention provides a composition comprising a protein,
5 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:59;
- (b) the amino acid sequence of SEQ ID NO:59 from amino acid 1 to amino acid 109;
- (c) a fragment of the amino acid sequence of SEQ ID NO:59, the fragment
10 comprising eight contiguous amino acids of SEQ ID NO:59; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CN238_1 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:59 or the amino acid sequence of SEQ ID
15 NO:59 from amino acid 1 to amino acid 109. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:59, or a protein comprising a
20 fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:59.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:60;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:60
25 from nucleotide 87 to nucleotide 1871;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:60 from nucleotide 628 to nucleotide 1882;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CO390_1 deposited with the ATCC under accession
30 number 98232;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CO390_1 deposited with the ATCC under accession number
35 98232;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:61;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:61;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:60.
- 15

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:60 from nucleotide 87 to nucleotide 1871; the nucleotide sequence of SEQ ID NO:60 from nucleotide 628 to nucleotide 1882; the nucleotide sequence of the full-length protein coding sequence of clone CO390_1 deposited with the ATCC under accession number 98232; or the nucleotide sequence of a mature protein coding sequence of clone CO390_1 deposited with the ATCC under accession number 98232. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232. In yet other preferred embodiments,

25 the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:61 from amino acid 182 to amino acid 248. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:61, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment comprising the amino acid sequence from amino acid 292 to amino acid 301 of SEQ ID NO:61.

30

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:60.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X

5 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:60, but excluding the poly(A) tail at the 3' end of SEQ ID NO:60; and

10 (ab) the nucleotide sequence of the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

15 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:60, but excluding the poly(A) tail at the 3' end of SEQ ID NO:60; and

(bb) the nucleotide sequence of the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:60, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:60 to a nucleotide sequence
30 corresponding to the 3' end of SEQ ID NO:60, but excluding the poly(A) tail at the 3' end of SEQ ID NO:60. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:60 from nucleotide 87 to nucleotide 1871, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:60 from nucleotide 87 to nucleotide 1871, to a
35 nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:60 from

nucleotide 87 to nucleotide 1871. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:60 from nucleotide 628 to nucleotide 1882, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:60 from nucleotide 628 to nucleotide 1882, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:60 from nucleotide 628 to nucleotide 1882.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:61;
- (b) the amino acid sequence of SEQ ID NO:61 from amino acid 182 to amino acid 248;
- (c) a fragment of the amino acid sequence of SEQ ID NO:61, the fragment comprising eight contiguous amino acids of SEQ ID NO:61; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CO390_1 deposited with the ATCC under accession number 98232;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:61 or the amino acid sequence of SEQ ID NO:61 from amino acid 182 to amino acid 248. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:61, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment comprising the amino acid sequence from amino acid 292 to amino acid 301 of SEQ ID NO:61.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:62;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:62 from nucleotide 68 to nucleotide 430;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:62 from nucleotide 128 to nucleotide 430;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AJ20_2 deposited with the ATCC under accession number 98261;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AJ20_2 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:63;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:63 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:63;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:62.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:62 from nucleotide 68 to nucleotide 430; the nucleotide sequence of SEQ ID NO:62 from nucleotide 128 to nucleotide 430; the nucleotide sequence of the full-length protein coding sequence of clone AJ20_2 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone AJ20_2 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:63 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:63, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:63 having biological activity, the fragment comprising the amino acid sequence from amino acid 55 to amino acid 64 of SEQ ID NO:63.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:62 and SEQ ID NO:64.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:62;
- 10 (ab) SEQ ID NO:64, but excluding the poly(A) tail at the 3' end of SEQ ID NO:64; and
- (ac) the nucleotide sequence of the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions
- 15 at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
- 20 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:62;
- (bb) SEQ ID NO:64, but excluding the poly(A) tail at the 3' end of SEQ ID NO:64; and
- 25 (bc) the nucleotide sequence of the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:62 and SEQ ID NO:64, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:62 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:64, but excluding the poly(A)

35 tail at the 3' end of SEQ ID NO:64. Also preferably the polynucleotide isolated according to the

above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:62, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:62 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:62. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:62 from nucleotide 68 to nucleotide 430, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:62 from nucleotide 68 to nucleotide 430, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:62 from nucleotide 68 to nucleotide 430. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:62 from nucleotide 128 to nucleotide 430, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:62 from nucleotide 128 to nucleotide 430, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:62 from nucleotide 128 to nucleotide 430.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:63;
- (b) a fragment of the amino acid sequence of SEQ ID NO:63, the fragment comprising eight contiguous amino acids of SEQ ID NO:63; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AJ20_2 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:63. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:63 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:63, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:63 having biological activity, the fragment comprising the amino acid sequence from amino acid 55 to amino acid 64 of SEQ ID NO:63.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:66;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:66 from nucleotide 289 to nucleotide 780;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AR440_1 deposited with the ATCC under accession number 98261;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AR440_1 deposited with the ATCC under accession number 98261;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AR440_1 deposited with the ATCC under accession number 98261;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AR440_1 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:67;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:67 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:67;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:66.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:66 from nucleotide 289 to nucleotide 780; the nucleotide sequence of the full-length protein coding sequence of clone AR440_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone AR440_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AR440_1 deposited with the ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:67 from amino acid 1 to amino acid 160. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:67 having biological activity, the fragment

preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:67, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:67 having biological activity, the fragment comprising the amino acid sequence from amino acid 77 to amino acid 86 of SEQ ID NO:67.

- 5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:66 and SEQ ID NO:65.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:65;
- (ab) SEQ ID NO:66, but excluding the poly(A) tail at the 3'
- 15 end of SEQ ID NO:66; and
- (ac) the nucleotide sequence of the cDNA insert of clone AR440_1 deposited with the ATCC under accession number 98261;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 25 consisting of:
- (ba) SEQ ID NO:65;
- (bb) SEQ ID NO:66, but excluding the poly(A) tail at the 3' end of SEQ ID NO:66; and
- (bc) the nucleotide sequence of the cDNA insert of clone
- 30 AR440_1 deposited with the ATCC under accession number 98261;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:65 and SEQ ID NO:66, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:65 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:66, but excluding the poly(A) tail at the 3' end of SEQ ID NO:66. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:66, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:66 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:66, but excluding the poly(A) tail at the 3' end of SEQ ID NO:66. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:66 from nucleotide 289 to nucleotide 780, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:66 from nucleotide 289 to nucleotide 780, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:66 from nucleotide 289 to nucleotide 780.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:67;
- (b) the amino acid sequence of SEQ ID NO:67 from amino acid 1 to amino acid 160;
- (c) a fragment of the amino acid sequence of SEQ ID NO:67, the fragment comprising eight contiguous amino acids of SEQ ID NO:67; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AR440_1 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:67 or the amino acid sequence of SEQ ID NO:67 from amino acid 1 to amino acid 160. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:67 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:67, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:67 having biological activity, the fragment comprising the amino acid sequence from amino acid 77 to amino acid 86 of SEQ ID NO:67.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:68;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:68 from nucleotide 76 to nucleotide 1050;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:68 from nucleotide 331 to nucleotide 567;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AS164_1 deposited with the ATCC under accession number 98261;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AS164_1 deposited with the ATCC under accession number 98261;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AS164_1 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AS164_1 deposited with the ATCC under accession number 98261;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:69;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:69;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:68.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:68 from nucleotide 76 to nucleotide 1050; the nucleotide sequence of SEQ ID NO:68 from nucleotide 331 to nucleotide 567; the nucleotide sequence of the full-length protein coding sequence of clone AS164_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone AS164_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AS164_1

30

35

deposited with the ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:69 from amino acid 87 to amino acid 164. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a
5 fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:69, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment comprising the amino acid sequence from amino acid 157 to amino acid 166 of SEQ ID NO:69.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:68.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:68, but excluding the poly(A) tail at the 3' end of SEQ ID NO:68; and
 - 20 (ab) the nucleotide sequence of the cDNA insert of clone AS164_1 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
 - 25 and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 30 (ba) SEQ ID NO:68, but excluding the poly(A) tail at the 3' end of SEQ ID NO:68; and
 - (bb) the nucleotide sequence of the cDNA insert of clone AS164_1 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions
 - 35 at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:68, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:68 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:68, but excluding the poly(A) tail at the 3' end of SEQ ID NO:68. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:68 from nucleotide 76 to nucleotide 1050, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:68 from nucleotide 76 to nucleotide 1050, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:68 from nucleotide 76 to nucleotide 1050. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:68 from nucleotide 331 to nucleotide 567, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:68 from nucleotide 331 to nucleotide 567, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:68 from nucleotide 331 to nucleotide 567.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:69;
- (b) the amino acid sequence of SEQ ID NO:69 from amino acid 87 to amino acid 164;
- (c) a fragment of the amino acid sequence of SEQ ID NO:69, the fragment comprising eight contiguous amino acids of SEQ ID NO:69; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AS164_1 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:69 or the amino acid sequence of SEQ ID NO:69 from amino acid 87 to amino acid 164. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:69, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment comprising the amino acid sequence from amino acid 157 to amino acid 166 of SEQ ID NO:69.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70
5 from nucleotide 242 to nucleotide 1060;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70
from nucleotide 596 to nucleotide 1060;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70
from nucleotide 10 to nucleotide 373;
- 10 (e) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone AX8_1 deposited with the ATCC under accession
number 98261;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone AX8_1 deposited with the ATCC under accession number 98261;
- 15 (g) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone AX8_1 deposited with the ATCC under accession number
98261;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone AX8_1 deposited with the ATCC under accession number 98261;
- 20 (i) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:71;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:71 having biological activity, the fragment comprising eight
contiguous amino acids of SEQ ID NO:71;
- 25 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i)
or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the
length of SEQ ID NO:70.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:70
35 from nucleotide 242 to nucleotide 1060; the nucleotide sequence of SEQ ID NO:70 from

nucleotide 596 to nucleotide 1060; the nucleotide sequence of SEQ ID NO:70 from nucleotide 10 to nucleotide 373; the nucleotide sequence of the full-length protein coding sequence of clone AX8_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone AX8_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AX8_1 deposited with the ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:71 from amino acid 1 to amino acid 44. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:71, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment comprising the amino acid sequence from amino acid 131 to amino acid 140 of SEQ ID NO:71.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:70.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:70, but excluding the poly(A) tail at the 3' end of SEQ ID NO:70; and
 - (ab) the nucleotide sequence of the cDNA insert of clone AX8_1 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:70, but excluding the poly(A) tail at the 3' end of SEQ ID NO:70; and

(bb) the nucleotide sequence of the cDNA insert of clone AX8_1 deposited with the ATCC under accession number 98261;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
10 sequence corresponding to the cDNA sequence of SEQ ID NO:70, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:70 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:70, but excluding the poly(A) tail at the 3' end of SEQ ID NO:70. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:70 from nucleotide
15 242 to nucleotide 1060, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:70 from nucleotide 242 to nucleotide 1060, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:70 from nucleotide 242 to nucleotide 1060. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID
20 NO:70 from nucleotide 596 to nucleotide 1060, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:70 from nucleotide 596 to nucleotide 1060, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:70 from nucleotide 596 to nucleotide 1060. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
25 sequence of SEQ ID NO:70 from nucleotide 10 to nucleotide 373, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:70 from nucleotide 10 to nucleotide 373, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:70 from nucleotide 10 to nucleotide 373.

In other embodiments, the present invention provides a composition comprising a protein,
30 wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:71;

(b) the amino acid sequence of SEQ ID NO:71 from amino acid 1 to amino acid 44;

(c) a fragment of the amino acid sequence of SEQ ID NO:71, the fragment
35 comprising eight contiguous amino acids of SEQ ID NO:71; and

(d) the amino acid sequence encoded by the cDNA insert of clone AX8_1 deposited with the ATCC under accession number 98261; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:71 or the amino acid sequence of SEQ ID NO:71 from amino acid 1 to amino acid 44. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:71, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment comprising the amino acid sequence from amino acid 131 to amino acid 140 of SEQ ID NO:71.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72 from nucleotide 773 to nucleotide 928;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72 from nucleotide 815 to nucleotide 928;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD176_3 deposited with the ATCC under accession number 98261;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD176_3 deposited with the ATCC under accession number 98261;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD176_3 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD176_3 deposited with the ATCC under accession number 98261;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:73;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:73;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

5 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:72.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:72 from nucleotide 773 to nucleotide 928; the nucleotide sequence of SEQ ID NO:72 from nucleotide
10 815 to nucleotide 928; the nucleotide sequence of the full-length protein coding sequence of clone BD176_3 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone BD176_3 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD176_3 deposited with the ATCC under
15 accession number 98261. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:73, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having
20 biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:73.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:72 and SEQ ID NO:74.

Further embodiments of the invention provide isolated polynucleotides produced
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:72;

(ab) SEQ ID NO:74, but excluding the poly(A) tail at the 3' end of SEQ ID NO:74; and

(ac) the nucleotide sequence of the cDNA insert of clone BD176_3 deposited with the ATCC under accession number 98261;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

5 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:72;

10 (bb) SEQ ID NO:74, but excluding the poly(A) tail at the 3' end of SEQ ID NO:74; and

(bc) the nucleotide sequence of the cDNA insert of clone BD176_3 deposited with the ATCC under accession number 98261;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:72 and SEQ ID NO:74, and
20 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:72 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:74, but excluding the poly(A) tail at the 3' end of SEQ ID NO:74. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:72, and extending contiguously from a nucleotide sequence corresponding to the 5' end of
25 SEQ ID NO:72 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:72. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:72 from nucleotide 773 to nucleotide 928, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:72 from nucleotide 773 to nucleotide 928, to a nucleotide
30 sequence corresponding to the 3' end of said sequence of SEQ ID NO:72 from nucleotide 773 to nucleotide 928. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:72 from nucleotide 815 to nucleotide 928, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:72 from nucleotide 815 to nucleotide

928, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:72 from nucleotide 815 to nucleotide 928.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 5 (a) the amino acid sequence of SEQ ID NO:73;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:73, the fragment comprising eight contiguous amino acids of SEQ ID NO:73; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone BD176_3 deposited with the ATCC under accession number 98261;
- 10 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:73. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:73, or a protein comprising
- 15 a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:73.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- 20 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 174 to nucleotide 440;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 1 to nucleotide 313;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD339_1 deposited with the ATCC under accession number 98261;
- 25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD339_1 deposited with the ATCC under accession number 98261;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD339_1 deposited with the ATCC under accession number 98261;
- 30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD339_1 deposited with the ATCC under accession number 98261;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- 35

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:75.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 174 to nucleotide 440; the nucleotide sequence of SEQ ID NO:75 from
15 nucleotide 1 to nucleotide 313; the nucleotide sequence of the full-length protein coding sequence of clone BD339_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone BD339_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD339_1 deposited with the
20 ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76 from amino acid 1 to amino acid 46. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight
25 (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:76.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
30 NO:75.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

(ab) the nucleotide sequence of the cDNA insert of clone BD339_1 deposited with the ATCC under accession number 98261;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

(bb) the nucleotide sequence of the cDNA insert of clone BD339_1 deposited with the ATCC under accession number 98261;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:75 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 174 to nucleotide 440, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 174 to nucleotide 440, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 174 to nucleotide 440. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 1 to nucleotide 313, and extending contiguously from a nucleotide

sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 1 to nucleotide 313, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 1 to nucleotide 313.

In other embodiments, the present invention provides a composition comprising a protein,
5 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
- (b) the amino acid sequence of SEQ ID NO:76 from amino acid 1 to amino acid 46;
- (c) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment
10 comprising eight contiguous amino acids of SEQ ID NO:76; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BD339_1 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:76 or the amino acid sequence of SEQ ID
15 NO:76 from amino acid 1 to amino acid 46. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment
20 comprising the amino acid sequence from amino acid 39 to amino acid 48 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77
25 from nucleotide 509 to nucleotide 619;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 1 to nucleotide 580;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD427_1 deposited with the ATCC under accession
30 number 98261;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD427_1 deposited with the ATCC under accession number 98261;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD427_1 deposited with the ATCC under accession number
35 98261;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD427_1 deposited with the ATCC under accession number 98261;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- 5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:77.
- 15
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 509 to nucleotide 619; the nucleotide sequence of SEQ ID NO:77 from nucleotide 1 to nucleotide 580; the nucleotide sequence of the full-length protein coding sequence of clone BD427_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone BD427_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD427_1 deposited with the ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 24. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 13 to amino acid 22 of SEQ ID NO:78.
- 20
- 25
- 30

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:77.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and

(ab) the nucleotide sequence of the cDNA insert of clone BD427_1 deposited with the ATCC under accession number 98261;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and

(bb) the nucleotide sequence of the cDNA insert of clone BD427_1 deposited with the ATCC under accession number 98261;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:77 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 509 to nucleotide 619, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 509 to nucleotide 619, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from

nucleotide 509 to nucleotide 619. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 1 to nucleotide 580, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 1 to nucleotide 580, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 1 to nucleotide 580.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
- 10 (b) the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 24;
- (c) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BD427_1
- 15 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:78 or the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 24. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, 20 most preferably thirty) contiguous amino acids of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 13 to amino acid 22 of SEQ ID NO:78.

In one embodiment, the present invention provides a composition comprising an isolated 25 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 300 to nucleotide 360;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BL229_22 deposited with the ATCC under accession number 98261;
- 30 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261;

- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BL229_22 deposited with the ATCC under accession number 98261;
- 5 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- 10 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 15 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:79.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:79 from nucleotide 300 to nucleotide 360; the nucleotide sequence of the full-length protein coding sequence of clone BL229_22 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone BL229_22 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide
- 25 encodes the full-length or a mature protein encoded by the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ
- 30 ID NO:80, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 5 to amino acid 14 of SEQ ID NO:80.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:79 and SEQ ID NO:81.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X

5 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:79;

(ab) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and

10 (ac) the nucleotide sequence of the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

20 (ba) SEQ ID NO:79;

(bb) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and

(bc) the nucleotide sequence of the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261;

25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:79 and SEQ ID NO:81, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID

35 NO:79, and extending contiguously from a nucleotide sequence corresponding to the 5' end of

SEQ ID NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:79. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 300 to nucleotide 360, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 300 to nucleotide 360, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 300 to nucleotide 360.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:80;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone BL229_22 deposited with the ATCC under accession number 98261;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:80. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a protein comprising
- 20 a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 5 to amino acid 14 of SEQ ID NO:80.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:82;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:82 from nucleotide 604 to nucleotide 771;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:82 from nucleotide 1 to nucleotide 684;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
- 30 protein coding sequence of clone BV123_16 deposited with the ATCC under accession number 98261;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV123_16 deposited with the ATCC under accession number 98261;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:83;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:83 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:83;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:82.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:82 from nucleotide 604 to nucleotide 771; the nucleotide sequence of SEQ ID NO:82 from nucleotide 1 to nucleotide 684; the nucleotide sequence of the full-length protein coding sequence of clone BV123_16 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature protein coding sequence of clone BV123_16 deposited with the ATCC under
- 25 accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:83 from amino acid 1 to amino acid 27. In further preferred embodiments, the present
- 30 invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:83 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:83, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:83 having biological activity, the fragment comprising the amino acid sequence from amino
- 35 acid 23 to amino acid 32 of SEQ ID NO:83.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:82.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 10 (aa) SEQ ID NO:82, but excluding the poly(A) tail at the 3' end of SEQ ID NO:82; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 15 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 20 (ba) SEQ ID NO:82, but excluding the poly(A) tail at the 3' end of SEQ ID NO:82; and
 - (bb) the nucleotide sequence of the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261;
 - 25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:82, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:82 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:82, but excluding the poly(A) tail at the 3' end of SEQ ID NO:82. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:82 from nucleotide 30 604 to nucleotide 771, and extending contiguously from a nucleotide sequence corresponding to

the 5' end of said sequence of SEQ ID NO:82 from nucleotide 604 to nucleotide 771, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:82 from nucleotide 604 to nucleotide 771. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:82 from nucleotide 1 to nucleotide 684, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:82 from nucleotide 1 to nucleotide 684, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:82 from nucleotide 1 to nucleotide 684.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:83;
- (b) the amino acid sequence of SEQ ID NO:83 from amino acid 1 to amino acid 27;
- (c) a fragment of the amino acid sequence of SEQ ID NO:83, the fragment comprising eight contiguous amino acids of SEQ ID NO:83; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BV123_16 deposited with the ATCC under accession number 98261;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:83 or the amino acid sequence of SEQ ID NO:83 from amino acid 1 to amino acid 27. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:83 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:83, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:83 having biological activity, the fragment comprising the amino acid sequence from amino acid 23 to amino acid 32 of SEQ ID NO:83.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:84;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:84 from nucleotide 43 to nucleotide 297;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:84 from nucleotide 94 to nucleotide 297;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:84 from nucleotide 1 to nucleotide 379;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CH377_1 deposited with the ATCC under accession number 98261;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CH377_1 deposited with the ATCC under accession number 98261;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:85;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:85 having biological activity, the fragment comprising eight
15 contiguous amino acids of SEQ ID NO:85;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:84.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:84 from nucleotide 43 to nucleotide 297; the nucleotide sequence of SEQ ID NO:84 from nucleotide 94 to nucleotide 297; the nucleotide sequence of SEQ ID NO:84 from nucleotide 1 to nucleotide 379; the nucleotide sequence of the full-length protein coding sequence of clone CH377_1 deposited with the ATCC under accession number 98261; or the nucleotide sequence of a mature
30 protein coding sequence of clone CH377_1 deposited with the ATCC under accession number 98261. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID
35 NO:85 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:85, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:85 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:85.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:84.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:84, but excluding the poly(A) tail at the 3' end of SEQ ID NO:84; and
 - 15 (ab) the nucleotide sequence of the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
 - 20 and
 - (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 25 (ba) SEQ ID NO:84, but excluding the poly(A) tail at the 3' end of SEQ ID NO:84; and
 - (bb) the nucleotide sequence of the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions
 - 30 at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:84, and extending contiguously
35 from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:84 to a nucleotide sequence

corresponding to the 3' end of SEQ ID NO:84, but excluding the poly(A) tail at the 3' end of SEQ ID NO:84. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:84 from nucleotide 43 to nucleotide 297, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:84 from nucleotide 43 to nucleotide 297, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:84 from nucleotide 43 to nucleotide 297. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:84 from nucleotide 94 to nucleotide 297, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:84 from nucleotide 94 to nucleotide 297, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:84 from nucleotide 94 to nucleotide 297. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:84 from nucleotide 1 to nucleotide 379, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:84 from nucleotide 1 to nucleotide 379, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:84 from nucleotide 1 to nucleotide 379.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:85;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:85, the fragment comprising eight contiguous amino acids of SEQ ID NO:85; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone CH377_1 deposited with the ATCC under accession number 98261;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:85. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:85 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:85, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:85 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:85.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 390 to nucleotide 563;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD441_1 deposited with the ATCC under accession number 98264;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD441_1 deposited with the ATCC under accession number 98264;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:87.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:87 from nucleotide 390 to nucleotide 563; the nucleotide sequence of the full-length protein coding sequence of clone BD441_1 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone BD441_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:88.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:87, SEQ ID NO:86, and SEQ ID NO:89.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:86;
 - (ab) SEQ ID NO:87;
 - 15 (ac) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and
 - (ad) the nucleotide sequence of the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions
 - 20 at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in
 - 25 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:86;
 - (bb) SEQ ID NO:87;
 - (bc) SEQ ID NO:89, but excluding the poly(A) tail at the 3'
 - 30 end of SEQ ID NO:89; and
 - (bd) the nucleotide sequence of the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions
 - at least as stringent as 4X SSC at 50 degrees C;
 - 35 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:86, SEQ ID NO:87, and SEQ ID NO:89, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:86 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:87 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:87. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 390 to nucleotide 563, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from nucleotide 390 to nucleotide 563, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 390 to nucleotide 563.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
- (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BD441_1 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:88. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:88.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:90;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:90 from nucleotide 583 to nucleotide 756;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD441_2 deposited with the ATCC under accession number 98264;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD441_2 deposited with the ATCC under accession number 98264;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD441_2 deposited with the ATCC under accession number 98264;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD441_2 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:91;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:91 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:91;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:90.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:90 from nucleotide 583 to nucleotide 756; the nucleotide sequence of the full-length protein coding sequence of clone BD441_2 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone BD441_2 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD441_2 deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:91 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:91, or a polynucleotide encoding a protein comprising a fragment of the amino acid

sequence of SEQ ID NO:91 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:91.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:90 and SEQ ID NO:92.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X
10 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:90;

(ab) SEQ ID NO:92, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:92; and

(ac) the nucleotide sequence of the cDNA insert of clone
15 BD441_2 deposited with the ATCC under accession number 98264;

(ii) hybridizing said probe(s) to human genomic DNA in conditions
at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in
6X SSC at 65 degrees C to a nucleotide sequence selected from the group
consisting of:

(ba) SEQ ID NO:90;

25 (bb) SEQ ID NO:92, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:92; and

(bc) the nucleotide sequence of the cDNA insert of clone
BD441_2 deposited with the ATCC under accession number 98264;

(ii) hybridizing said primer(s) to human genomic DNA in conditions
30 at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
sequence corresponding to the cDNA sequences of SEQ ID NO:90 and SEQ ID NO:92, and
35 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:90

to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:92, but excluding the poly(A) tail at the 3' end of SEQ ID NO:92. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:90, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:90 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:90. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:90 from nucleotide 583 to nucleotide 756, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:90 from nucleotide 583 to nucleotide 756, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:90 from nucleotide 583 to nucleotide 756.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:91;
- (b) a fragment of the amino acid sequence of SEQ ID NO:91, the fragment comprising eight contiguous amino acids of SEQ ID NO:91; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BD441_2 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:91. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:91 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:91, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:91 having biological activity, the fragment comprising the amino acid sequence from amino acid 24 to amino acid 33 of SEQ ID NO:91.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 426 to nucleotide 581;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 495 to nucleotide 581;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 354 to nucleotide 503;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG102_3 deposited with the ATCC under accession number 98264;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG102_3 deposited with the ATCC under accession number 98264;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- 15 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:93.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:93 from nucleotide 426 to nucleotide 581; the nucleotide sequence of SEQ ID NO:93 from nucleotide 495 to nucleotide 581; the nucleotide sequence of SEQ ID NO:93 from nucleotide 354 to nucleotide 503; the nucleotide sequence of the full-length protein coding sequence of clone BG102_3 deposited with the ATCC under accession number 98264; or the nucleotide sequence
- 30 of a mature protein coding sequence of clone BG102_3 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:94 from
- 35 amino acid 1 to amino acid 26. In further preferred embodiments, the present invention provides

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having
5 biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:94.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:93.

Further embodiments of the invention provide isolated polynucleotides produced
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

(ab) the nucleotide sequence of the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

30 (bb) the nucleotide sequence of the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:93 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 426 to nucleotide 581, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 426 to nucleotide 581, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 426 to nucleotide 581. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 495 to nucleotide 581, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 495 to nucleotide 581, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 495 to nucleotide 581. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 354 to nucleotide 503, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 354 to nucleotide 503, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 354 to nucleotide 503.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
 - (b) the amino acid sequence of SEQ ID NO:94 from amino acid 1 to amino acid 26;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BG102_3 deposited with the ATCC under accession number 98264;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:94 or the amino acid sequence of SEQ ID NO:94 from amino acid 1 to amino acid 26. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a protein comprising a

fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:94.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95
from nucleotide 112 to nucleotide 978;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95
from nucleotide 436 to nucleotide 1048;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone BK158_1 deposited with the ATCC under accession
number 98264;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone BK158_1 deposited with the ATCC under accession number 98264;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone BK158_1 deposited with the ATCC under accession number
98264;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone BK158_1 deposited with the ATCC under accession number 98264;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:96;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising eight
contiguous amino acids of SEQ ID NO:96;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g)
above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h)
or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
length of SEQ ID NO:95.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:95
35 from nucleotide 112 to nucleotide 978; the nucleotide sequence of SEQ ID NO:95 from

nucleotide 436 to nucleotide 1048; the nucleotide sequence of the full-length protein coding sequence of clone BK158_1 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone BK158_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide
5 encodes the full-length or a mature protein encoded by the cDNA insert of clone BK158_1 deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ
10 ID NO:96, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 139 to amino acid 148 of SEQ ID NO:96.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:95.

15 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X
20 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

- (ab) the nucleotide sequence of the cDNA insert of clone
BK158_1 deposited with the ATCC under accession number 98264;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in
30 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

- (bb) the nucleotide sequence of the cDNA insert of clone BK158_1 deposited with the ATCC under accession number 98264;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:95 to a nucleotide sequence

10 corresponding to the 3' end of SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 112 to nucleotide 978, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 112 to nucleotide 978, to a

15 nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 112 to nucleotide 978. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 436 to nucleotide 1048, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 436 to

20 nucleotide 1048, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 436 to nucleotide 1048.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:96;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BK158_1 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein

30 comprises the amino acid sequence of SEQ ID NO:96. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:96, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment

35 comprising the amino acid sequence from amino acid 139 to amino acid 148 of SEQ ID NO:96.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97
5 from nucleotide 16 to nucleotide 492;
 - (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BP163_1 deposited with the ATCC under accession number 98264;
 - (d) a polynucleotide encoding the full-length protein encoded by the cDNA
10 insert of clone BP163_1 deposited with the ATCC under accession number 98264;
 - (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BP163_1 deposited with the ATCC under accession number 98264;
 - (f) a polynucleotide encoding a mature protein encoded by the cDNA insert
15 of clone BP163_1 deposited with the ATCC under accession number 98264;
 - (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:98;
 - (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising eight
20 contiguous amino acids of SEQ ID NO:98;
 - (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
 - (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
 - (k) a polynucleotide that hybridizes under stringent conditions to any one of
25 the polynucleotides specified in (a)-(h); and
 - (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:97.
- 30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:97 from nucleotide 16 to nucleotide 492; the nucleotide sequence of the full-length protein coding sequence of clone BP163_1 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone BP163_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide
35 encodes the full-length or a mature protein encoded by the cDNA insert of clone BP163_1

deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ
5 ID NO:98, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:98.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:97 and SEQ ID NO:99.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X
SSC at 65 degrees C to a nucleotide sequence selected from the group consisting
15 of:

(aa) SEQ ID NO:97;

(ab) SEQ ID NO:99, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:99; and

(ac) the nucleotide sequence of the cDNA insert of clone
20 BP163_1 deposited with the ATCC under accession number 98264;

(ii) hybridizing said probe(s) to human genomic DNA in conditions
at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in
6X SSC at 65 degrees C to a nucleotide sequence selected from the group
consisting of:

(ba) SEQ ID NO:97;

(bb) SEQ ID NO:99, but excluding the poly(A) tail at the 3'
30 end of SEQ ID NO:99; and

(bc) the nucleotide sequence of the cDNA insert of clone
BP163_1 deposited with the ATCC under accession number 98264;

(ii) hybridizing said primer(s) to human genomic DNA in conditions
35 at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:97 and SEQ ID NO:99, and
 5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:97 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97, and extending contiguously from a nucleotide sequence corresponding to the 5' end of
 10 SEQ ID NO:97 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:97. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 16 to nucleotide 492, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 16 to nucleotide 492, to a nucleotide
 15 sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide 16 to nucleotide 492.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:98;
- 20 (b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BP163_1 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein
 25 comprises the amino acid sequence of SEQ ID NO:98. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:98, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment
 30 comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:98.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101
 35 from nucleotide 72 to nucleotide 569;

- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BZ16_3 deposited with the ATCC under accession number 98264;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BZ16_3 deposited with the ATCC under accession number 98264;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising 15 eight contiguous amino acids of SEQ ID NO:102;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:101.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:101 from nucleotide 72 to nucleotide 569; the nucleotide sequence of the full-length protein coding sequence of clone BZ16_3 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone BZ16_3 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide 30 encodes the full-length or a mature protein encoded by the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102 from amino acid 1 to amino acid 124. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a 35 fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment

preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:102, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:102.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:101 and SEQ ID NO:100.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:100;
 - (ab) SEQ ID NO:101, but excluding the poly(A) tail at the 3'
 - 15 end of SEQ ID NO:101; and
 - (ac) the nucleotide sequence of the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
 - 25 consisting of:
 - (ba) SEQ ID NO:100;
 - (bb) SEQ ID NO:101, but excluding the poly(A) tail at the 3'
 - end of SEQ ID NO:101; and
 - (bc) the nucleotide sequence of the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - 30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:100 and SEQ ID NO:101, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:100 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:101 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101 from nucleotide 72 to nucleotide 569, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:101 from nucleotide 72 to nucleotide 569, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:101 from nucleotide 72 to nucleotide 569.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
 - (b) the amino acid sequence of SEQ ID NO:102 from amino acid 1 to amino acid 124;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BZ16_3 deposited with the ATCC under accession number 98264;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:102 or the amino acid sequence of SEQ ID NO:102 from amino acid 1 to amino acid 124. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:102, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:102.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103 from nucleotide 405 to nucleotide 662;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103 from nucleotide 519 to nucleotide 662;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103 from nucleotide 1 to nucleotide 584;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CC182_1 deposited with the ATCC under accession number 98264;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CC182_1 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CC182_1 deposited with the ATCC under accession number 98264;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CC182_1 deposited with the ATCC under accession number 98264;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:103.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:103 from nucleotide 405 to nucleotide 662; the nucleotide sequence of SEQ ID NO:103 from nucleotide 519 to nucleotide 662; the nucleotide sequence of SEQ ID NO:103 from nucleotide 1 to nucleotide 584; the nucleotide sequence of the full-length protein coding sequence of clone

CC182_1 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone CC182_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CC182_1 deposited with the ATCC under
5 accession number 98264. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:104 from amino acid 1 to amino acid 60. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably
10 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:104.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
15 NO:103.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X
20 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and
 - (ab) the nucleotide sequence of the cDNA insert of clone
25 CC182_1 deposited with the ATCC under accession number 98264;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in
30 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:103, but excluding the poly(A) tail at the 3'
35 end of SEQ ID NO:103; and

- (bb) the nucleotide sequence of the cDNA insert of clone CC182_1 deposited with the ATCC under accession number 98264;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:103 to a nucleotide

10 sequence corresponding to the 3' end of SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 405 to nucleotide 662, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from nucleotide 405 to nucleotide

15 662, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 405 to nucleotide 662. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 519 to nucleotide 662, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from nucleotide 519

20 to nucleotide 662, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 519 to nucleotide 662. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 1 to nucleotide 584, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:103 from

25 nucleotide 1 to nucleotide 584, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 1 to nucleotide 584.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- 30 (b) the amino acid sequence of SEQ ID NO:104 from amino acid 1 to amino acid 60;
- (c) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CC182_1
- 35 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:104 or the amino acid sequence of SEQ ID NO:104 from amino acid 1 to amino acid 60. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID
 5 NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:104.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 311 to nucleotide 409;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 24 to nucleotide 414;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CG109_1 deposited with the ATCC under accession number 98264;
- 20 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CG109_1 deposited with the ATCC under accession number 98264;
- 25 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino
 30 acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h)
 35 or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:105.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:105 from nucleotide 311 to nucleotide 409; the nucleotide sequence of SEQ ID NO:105 from nucleotide 24 to nucleotide 414; the nucleotide sequence of the full-length protein coding sequence of clone CG109_1 deposited with the ATCC under accession number 98264; or the
10 nucleotide sequence of a mature protein coding sequence of clone CG109_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the
15 amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 11 to amino acid 20 of SEQ ID NO:106.

20 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:105.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- 25 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and
- 30 (ab) the nucleotide sequence of the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);

35 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and

(bb) the nucleotide sequence of the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105, and extending contiguously
15 from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:105 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 311 to nucleotide 409, and extending contiguously from a nucleotide sequence
20 corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 311 to nucleotide 409, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 311 to nucleotide 409. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 24 to nucleotide 414, and extending contiguously from a nucleotide
25 sequence corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 24 to nucleotide 414, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 24 to nucleotide 414.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:106;

(b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and

(c) the amino acid sequence encoded by the cDNA insert of clone CG109_1 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:106. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably
5 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 11 to amino acid 20 of SEQ ID NO:106.

In one embodiment, the present invention provides a composition comprising an isolated
10 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:108;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:108 from nucleotide 471 to nucleotide 611;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length
15 protein coding sequence of clone CJ397_1 deposited with the ATCC under accession number 98264;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CJ397_1 deposited with the ATCC under accession number 98264;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein
20 coding sequence of clone CJ397_1 deposited with the ATCC under accession number 98264;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CJ397_1 deposited with the ATCC under accession number 98264;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence
25 of SEQ ID NO:109;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:109 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:109;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f)
30 above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:108.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:108
5 from nucleotide 471 to nucleotide 611; the nucleotide sequence of the full-length protein coding sequence of clone CJ397_1 deposited with the ATCC under accession number 98264; or the nucleotide sequence of a mature protein coding sequence of clone CJ397_1 deposited with the ATCC under accession number 98264. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CJ397_1
10 deposited with the ATCC under accession number 98264. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:109 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:109, or a polynucleotide encoding a protein comprising a fragment of the amino acid
15 sequence of SEQ ID NO:109 having biological activity, the fragment comprising the amino acid sequence from amino acid 18 to amino acid 27 of SEQ ID NO:109.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:108, SEQ ID NO:107, and SEQ ID NO:110.

Further embodiments of the invention provide isolated polynucleotides produced
20 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 25 (aa) SEQ ID NO:107;
(ab) SEQ ID NO:108;
(ac) SEQ ID NO:110, but excluding the poly(A) tail at the 3' end of SEQ ID NO:110; and
(ad) the nucleotide sequence of the cDNA insert of clone
30 CJ397_1 deposited with the ATCC under accession number 98264;
(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
(iii) isolating the DNA polynucleotides detected with the probe(s);

and

35 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:107;
(bb) SEQ ID NO:108;
(bc) SEQ ID NO:110, but excluding the poly(A) tail at the 3' end of SEQ ID NO:110; and
(bd) the nucleotide sequence of the cDNA insert of clone CJ397_1 deposited with the ATCC under accession number 98264;
- 10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide

15 sequence corresponding to the cDNA sequences of SEQ ID NO:107, SEQ ID NO:108, and SEQ ID NO:110, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:107 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:110, but excluding the poly(A) tail at the 3' end of SEQ ID NO:110. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the

20 cDNA sequence of SEQ ID NO:108, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:108 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:108. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:108 from nucleotide 471 to nucleotide 611, and extending contiguously from a nucleotide sequence

25 corresponding to the 5' end of said sequence of SEQ ID NO:108 from nucleotide 471 to nucleotide 611, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:108 from nucleotide 471 to nucleotide 611.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:109;
(b) a fragment of the amino acid sequence of SEQ ID NO:109, the fragment comprising eight contiguous amino acids of SEQ ID NO:109; and
(c) the amino acid sequence encoded by the cDNA insert of clone CJ397_1 deposited with the ATCC under accession number 98264;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:109. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:109 having biological activity, the fragment preferably comprising eight (more preferably
5 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:109, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:109 having biological activity, the fragment comprising the amino acid sequence from amino acid 18 to amino acid 27 of SEQ ID NO:109.

In one embodiment, the present invention provides a composition comprising an isolated
10 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111 from nucleotide 141 to nucleotide 1532;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111
15 from nucleotide 204 to nucleotide 1532;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111 from nucleotide 78 to nucleotide 476;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM795_4 deposited with the ATCC under accession
20 number 98271;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM795_4 deposited with the ATCC under accession number
25 98271;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino
30 acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

5 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:111.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:111 from nucleotide 141 to nucleotide 1532; the nucleotide sequence of SEQ ID NO:111 from
10 nucleotide 204 to nucleotide 1532; the nucleotide sequence of SEQ ID NO:111 from nucleotide 78 to nucleotide 476; the nucleotide sequence of the full-length protein coding sequence of clone AM795_4 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone AM795_4 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a
15 mature protein encoded by the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:112 from amino acid 1 to amino acid 112. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
20 ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 227 to amino acid 236 of SEQ ID NO:112.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:111.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:
30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and

- (ab) the nucleotide sequence of the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 10 consisting of:
- (ba) SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111; and
- (bb) the nucleotide sequence of the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide
- 20 sequence corresponding to the cDNA sequence of SEQ ID NO:111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:111 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111
- 25 from nucleotide 141 to nucleotide 1532, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 141 to nucleotide 1532, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 141 to nucleotide 1532. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of
- 30 SEQ ID NO:111 from nucleotide 204 to nucleotide 1532, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 204 to nucleotide 1532, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 204 to nucleotide 1532. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence
- 35 corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 78 to nucleotide 476,

and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 78 to nucleotide 476, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 78 to nucleotide 476.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
- (b) the amino acid sequence of SEQ ID NO:112 from amino acid 1 to amino acid 112;
- 10 (c) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AM795_4 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein
15 comprises the amino acid sequence of SEQ ID NO:112 or the amino acid sequence of SEQ ID NO:112 from amino acid 1 to amino acid 112. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a protein
20 comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 227 to amino acid 236 of SEQ ID NO:112.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:114;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:114 from nucleotide 19 to nucleotide 262;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:114 from nucleotide 91 to nucleotide 262;
- 30 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AT340_1 deposited with the ATCC under accession number 98271;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AT340_1 deposited with the ATCC under accession number 98271;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:115;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:115 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:115;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:114.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:114 from nucleotide 19 to nucleotide 262; the nucleotide sequence of SEQ ID NO:114 from nucleotide 91 to nucleotide 262; the nucleotide sequence of the full-length protein coding sequence of clone AT340_1 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone AT340_1 deposited with the ATCC under accession
- 25 number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:115 from amino acid 1 to amino acid 66. In further preferred embodiments, the present invention provides
- 30 a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:115 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:115, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:115 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino
- 35 acid 44 of SEQ ID NO:115.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:114 and SEQ ID NO:113.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (aa) SEQ ID NO:113;
- (ab) SEQ ID NO:114, but excluding the poly(A) tail at the 3' end of SEQ ID NO:114; and
- (ac) the nucleotide sequence of the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271;
- 15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
- 20 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:113;
- (bb) SEQ ID NO:114, but excluding the poly(A) tail at the 3' end of SEQ ID NO:114; and
- 25 (bc) the nucleotide sequence of the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:113 and SEQ ID NO:114, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:113 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:114, but excluding

35 the poly(A) tail at the 3' end of SEQ ID NO:114. Also preferably the polynucleotide isolated

according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:114, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:114 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:114, but excluding the poly(A) tail at the 3' end of SEQ ID NO:114. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:114 from nucleotide 19 to nucleotide 262, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:114 from nucleotide 19 to nucleotide 262, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:114 from nucleotide 19 to nucleotide 262. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:114 from nucleotide 91 to nucleotide 262, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:114 from nucleotide 91 to nucleotide 262, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:114 from nucleotide 91 to nucleotide 262.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:115;
 - (b) the amino acid sequence of SEQ ID NO:115 from amino acid 1 to amino acid 66;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:115, the fragment comprising eight contiguous amino acids of SEQ ID NO:115; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone AT340_1 deposited with the ATCC under accession number 98271;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:115 or the amino acid sequence of SEQ ID NO:115 from amino acid 1 to amino acid 66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:115 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:115, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:115 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:115.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:116;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:116 from nucleotide 2 to nucleotide 601;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:116 from nucleotide 401 to nucleotide 601;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG132_1 deposited with the ATCC under accession number 98271;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG132_1 deposited with the ATCC under accession number 98271;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:117;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:117 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:117;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:116.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:116 from nucleotide 2 to nucleotide 601; the nucleotide sequence of SEQ ID NO:116 from nucleotide 401 to nucleotide 601; the nucleotide sequence of the full-length protein coding sequence of clone BG132_1 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone BG132_1 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a

mature protein encoded by the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:117 from amino acid 119 to amino acid 200. In further preferred embodiments, the present invention
5 provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:117 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:117, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:117 having biological activity, the fragment comprising the amino acid sequence from amino
10 acid 95 to amino acid 104 of SEQ ID NO:117.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:116 and SEQ ID NO:118.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:116;
 - 20 (ab) SEQ ID NO:118, but excluding the poly(A) tail at the 3' end of SEQ ID NO:118; and
 - (ac) the nucleotide sequence of the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions
25 at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
30 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:116;
 - (bb) SEQ ID NO:118, but excluding the poly(A) tail at the 3' end of SEQ ID NO:118; and

(bc) the nucleotide sequence of the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

5 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:116 and SEQ ID NO:118, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:116 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:118, but excluding the poly(A) tail at the 3' end of SEQ ID NO:118. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:116, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:116 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:116. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:116 from nucleotide 2 to nucleotide 601, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:116 from nucleotide 2 to nucleotide 601, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:116 from nucleotide 2 to nucleotide 601. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:116 from nucleotide 401 to nucleotide 601, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:116 from nucleotide 401 to nucleotide 601, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:116 from nucleotide 401 to nucleotide 601.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:117;

30 (b) the amino acid sequence of SEQ ID NO:117 from amino acid 119 to amino acid 200;

(c) a fragment of the amino acid sequence of SEQ ID NO:117, the fragment comprising eight contiguous amino acids of SEQ ID NO:117; and

(d) the amino acid sequence encoded by the cDNA insert of clone BG132_1 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:117 or the amino acid sequence of SEQ ID NO:117 from amino acid 119 to amino acid 200. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:117 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:117, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:117 having biological activity, the fragment comprising the amino acid sequence from amino acid 95 to amino acid 104 of SEQ ID NO:117.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:119;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:119 from nucleotide 225 to nucleotide 701;
- 15 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG219_2 deposited with the ATCC under accession number 98271;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG219_2 deposited with the ATCC under accession number 98271;
- 20 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG219_2 deposited with the ATCC under accession number 98271;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG219_2 deposited with the ATCC under accession number 98271;
- 25 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:120;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:120;
- 30 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of
- 35 the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:119.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:119 from nucleotide 225 to nucleotide 701; the nucleotide sequence of the full-length protein coding sequence of clone BG219_2 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone BG219_2 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG219_2 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:120 from amino acid 1 to amino acid 97. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:120, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:120.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:119.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BG219_2 deposited with the ATCC under accession number 98271;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119; and
- (bb) the nucleotide sequence of the cDNA insert of clone BG219_2 deposited with the ATCC under accession number 98271;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 10 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:119, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:119 to a nucleotide

15 sequence corresponding to the 3' end of SEQ ID NO:119, but excluding the poly(A) tail at the 3' end of SEQ ID NO:119. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:119 from nucleotide 225 to nucleotide 701, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:119 from nucleotide 225 to nucleotide

20 701, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:119 from nucleotide 225 to nucleotide 701.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:120;
- 25 (b) the amino acid sequence of SEQ ID NO:120 from amino acid 1 to amino acid 97;
- (c) a fragment of the amino acid sequence of SEQ ID NO:120, the fragment comprising eight contiguous amino acids of SEQ ID NO:120; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BG219_2
- 30 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:120 or the amino acid sequence of SEQ ID NO:120 from amino acid 1 to amino acid 97. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID

35 NO:120 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:120, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:120 having biological activity, the fragment comprising the amino acid sequence from amino acid 74 to amino acid 83 of SEQ ID NO:120.

- 5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121
10 from nucleotide 2115 to nucleotide 2510;
 - (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121
10 from nucleotide 1 to nucleotide 324;
 - (d) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone BG366_2 deposited with the ATCC under accession
number 98271;
 - 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone BG366_2 deposited with the ATCC under accession number 98271;
 - (f) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone BG366_2 deposited with the ATCC under accession number
98271;
 - 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone BG366_2 deposited with the ATCC under accession number 98271;
 - (h) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:122;
 - (i) a polynucleotide encoding a protein comprising a fragment of the amino
25 acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising
eight contiguous amino acids of SEQ ID NO:122;
 - (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g)
above;
 - (k) a polynucleotide which encodes a species homologue of the protein of (h)
30 or (i) above ;
 - (l) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i); and
 - (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
35 length of SEQ ID NO:121.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:121 from nucleotide 2115 to nucleotide 2510; the nucleotide sequence of SEQ ID NO:121 from nucleotide 1 to nucleotide 324; the nucleotide sequence of the full-length protein coding sequence of clone BG366_2 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone BG366_2 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG366_2 deposited with the ATCC under accession number 98271. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:122, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:122.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:121.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BG366_2 deposited with the ATCC under accession number 98271;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121; and

(bb) the nucleotide sequence of the cDNA insert of clone BG366_2 deposited with the ATCC under accession number 98271;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:121, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:121 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:121, but excluding the poly(A) tail at the 3' end of SEQ ID NO:121. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:121
15 from nucleotide 2115 to nucleotide 2510, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:121 from nucleotide 2115 to nucleotide 2510, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:121 from nucleotide 2115 to nucleotide 2510. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
20 sequence of SEQ ID NO:121 from nucleotide 1 to nucleotide 324, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:121 from nucleotide 1 to nucleotide 324, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:121 from nucleotide 1 to nucleotide 324.

In other embodiments, the present invention provides a composition comprising a protein,
25 wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:122;

(b) a fragment of the amino acid sequence of SEQ ID NO:122, the fragment comprising eight contiguous amino acids of SEQ ID NO:122; and

(c) the amino acid sequence encoded by the cDNA insert of clone BG366_2
30 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:122. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably comprising eight (more preferably
35 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:122, or a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:122.

- In one embodiment, the present invention provides a composition comprising an isolated
- 5 polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:123;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 215;
 - (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:123
10 from nucleotide 27 to nucleotide 181;
 - (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BV172_2 deposited with the ATCC under accession number 98271;
 - (e) a polynucleotide encoding the full-length protein encoded by the cDNA
15 insert of clone BV172_2 deposited with the ATCC under accession number 98271;
 - (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV172_2 deposited with the ATCC under accession number 98271;
 - (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
20 of clone BV172_2 deposited with the ATCC under accession number 98271;
 - (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:124;
 - (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising
25 eight contiguous amino acids of SEQ ID NO:124;
 - (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
 - (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
 - (l) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(i); and
 - (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:123.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 215; the nucleotide sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 181; the nucleotide sequence of the full-length protein coding sequence of clone BV172_2 deposited with the ATCC under accession number 98271; or the
5 nucleotide sequence of a mature protein coding sequence of clone BV172_2 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV172_2 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid
10 sequence of SEQ ID NO:124 from amino acid 1 to amino acid 51. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:124, or a polynucleotide encoding a protein comprising a fragment of the
15 amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:124.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:123.

Further embodiments of the invention provide isolated polynucleotides produced
20 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
25 (aa) SEQ ID NO:123, but excluding the poly(A) tail at the 3' end of SEQ ID NO:123; and
(ab) the nucleotide sequence of the cDNA insert of clone BV172_2 deposited with the ATCC under accession number 98271;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions
30 at least as stringent as 4X SSC at 50 degrees C; and
(iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:123, but excluding the poly(A) tail at the 3' end of SEQ ID NO:123; and

(bb) the nucleotide sequence of the cDNA insert of clone BV172_2 deposited with the ATCC under accession number 98271;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

10 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:123, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:123 to a nucleotide
15 sequence corresponding to the 3' end of SEQ ID NO:123, but excluding the poly(A) tail at the 3' end of SEQ ID NO:123. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 215, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide
20 215, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 215. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 181, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:123 from nucleotide 27 to
25 nucleotide 181, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:123 from nucleotide 27 to nucleotide 181.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:124;
30 (b) the amino acid sequence of SEQ ID NO:124 from amino acid 1 to amino acid 51;

(c) a fragment of the amino acid sequence of SEQ ID NO:124, the fragment comprising eight contiguous amino acids of SEQ ID NO:124; and

(d) the amino acid sequence encoded by the cDNA insert of clone BV172_2
35 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:124 or the amino acid sequence of SEQ ID NO:124 from amino acid 1 to amino acid 51. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:124, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:124 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:124.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125 from nucleotide 338 to nucleotide 409;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125 from nucleotide 362 to nucleotide 409;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:125 from nucleotide 270 to nucleotide 419;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CC247_10 deposited with the ATCC under accession number 98271;
- 20 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CC247_10 deposited with the ATCC under accession number 98271;
- 25 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:126;
- 30 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:126;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
- 35 above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

5 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:125.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:125 from nucleotide 338 to nucleotide 409; the nucleotide sequence of SEQ ID NO:125 from
10 nucleotide 362 to nucleotide 409; the nucleotide sequence of SEQ ID NO:125 from nucleotide 270 to nucleotide 419; the nucleotide sequence of the full-length protein coding sequence of clone CC247_10 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone CC247_10 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-
15 length or a mature protein encoded by the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:126, or a
20 polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising the amino acid sequence from amino acid 7 to amino acid 16 of SEQ ID NO:126.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:125.

25 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting
30 of:

(aa) SEQ ID NO:125, but excluding the poly(A) tail at the 3' end of SEQ ID NO:125; and

(ab) the nucleotide sequence of the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

5 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:125, but excluding the poly(A) tail at the 3' end of SEQ ID NO:125; and

(bb) the nucleotide sequence of the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

15 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:125 to a nucleotide
20 sequence corresponding to the 3' end of SEQ ID NO:125, but excluding the poly(A) tail at the 3' end of SEQ ID NO:125. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125 from nucleotide 338 to nucleotide 409, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:125 from nucleotide 338 to nucleotide
25 409, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:125 from nucleotide 338 to nucleotide 409. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125 from nucleotide 362 to nucleotide 409, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:125 from nucleotide 362
30 to nucleotide 409, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:125 from nucleotide 362 to nucleotide 409. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:125 from nucleotide 270 to nucleotide 419, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:125 from

nucleotide 270 to nucleotide 419, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:125 from nucleotide 270 to nucleotide 419.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 5 (a) the amino acid sequence of SEQ ID NO:126;
- (b) a fragment of the amino acid sequence of SEQ ID NO:126, the fragment comprising eight contiguous amino acids of SEQ ID NO:126; and
- (c) the amino acid sequence encoded by the cDNA insert of clone CC247_10 deposited with the ATCC under accession number 98271;
- 10 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:126. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:126, or a protein
- 15 comprising a fragment of the amino acid sequence of SEQ ID NO:126 having biological activity, the fragment comprising the amino acid sequence from amino acid 7 to amino acid 16 of SEQ ID NO:126.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 20 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127 from nucleotide 128 to nucleotide 1600;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127 from nucleotide 281 to nucleotide 1600;
- 25 (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127 from nucleotide 62 to nucleotide 373;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CI480_9 deposited with the ATCC under accession number 98271;
- 30 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CI480_9 deposited with the ATCC under accession number 98271;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CI480_9 deposited with the ATCC under accession number 98271;

- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CI480_9 deposited with the ATCC under accession number 98271;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:128;
- 5 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:128;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- 10 (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of
- 15 the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:127.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:127 from nucleotide 128 to nucleotide 1600; the nucleotide sequence of SEQ ID NO:127 from nucleotide 281 to nucleotide 1600; the nucleotide sequence of SEQ ID NO:127 from nucleotide

20 62 to nucleotide 373; the nucleotide sequence of the full-length protein coding sequence of clone CI480_9 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone CI480_9 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CI480_9 deposited with the ATCC under

25 accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:128 from amino acid 1 to amino acid 82. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment preferably comprising eight (more preferably

30 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:128, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:128.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID

35 NO:127.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:127, but excluding the poly(A) tail at the 3' end of SEQ ID NO:127; and

(ab) the nucleotide sequence of the cDNA insert of clone CI480_9 deposited with the ATCC under accession number 98271;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:127, but excluding the poly(A) tail at the 3' end of SEQ ID NO:127; and

(bb) the nucleotide sequence of the cDNA insert of clone CI480_9 deposited with the ATCC under accession number 98271;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:127 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:127, but excluding the poly(A) tail at the 3' end of SEQ ID NO:127. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127 from nucleotide 128 to nucleotide 1600, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:127 from nucleotide 128 to nucleotide 1600, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:127

from nucleotide 128 to nucleotide 1600. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127 from nucleotide 281 to nucleotide 1600, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:127 from nucleotide 281 to nucleotide 1600, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:127 from nucleotide 281 to nucleotide 1600. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:127 from nucleotide 62 to nucleotide 373, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:127 from nucleotide 62 to nucleotide 373, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:127 from nucleotide 62 to nucleotide 373.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:128;
- (b) the amino acid sequence of SEQ ID NO:128 from amino acid 1 to amino acid 82;
- (c) a fragment of the amino acid sequence of SEQ ID NO:128, the fragment comprising eight contiguous amino acids of SEQ ID NO:128; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CI480_9 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:128 or the amino acid sequence of SEQ ID NO:128 from amino acid 1 to amino acid 82. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:128, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising the amino acid sequence from amino acid 240 to amino acid 249 of SEQ ID NO:128.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide 488;
- 5 (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;
- 10 (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;
- 15 (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:130;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:130;
- 20 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- 25 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:129.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958; the nucleotide sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958; the nucleotide sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide 488; the nucleotide sequence of the full-length protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-length or a

30

35

mature protein encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:130 from amino acid 1 to amino acid 34. In further preferred embodiments, the present invention provides
5 a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:130, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising the amino acid sequence from amino acid 591 to
10 amino acid 600 of SEQ ID NO:130.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:129.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:129, but excluding the poly(A) tail at the 3'
20 end of SEQ ID NO:129; and
- (ab) the nucleotide sequence of the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
30 consisting of:
- (ba) SEQ ID NO:129, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:129; and
- (bb) the nucleotide sequence of the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).
- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:129 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:129, but excluding the poly(A) tail at the 3' end of SEQ ID NO:129. Also preferably the polynucleotide isolated according to the above
- 10 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958. Also preferably the polynucleotide isolated according
- 15 to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958. Also preferably the
- 20 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide 488, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide 488, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide
- 25 488.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:130;
- (b) the amino acid sequence of SEQ ID NO:130 from amino acid 1 to amino
- 30 acid 34;
- (c) a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:130 or the amino acid sequence of SEQ ID NO:130 from amino acid 1 to amino acid 34. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:130, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130 having biological activity, the fragment comprising the amino acid sequence from amino acid 591 to amino acid 600 of SEQ ID NO:130.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131 from nucleotide 914 to nucleotide 2353;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131 from nucleotide 1793 to nucleotide 2353;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:131 from nucleotide 1037 to nucleotide 1260;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CT748_2 deposited with the ATCC under accession number 98271;
- 20 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CT748_2 deposited with the ATCC under accession number 98271;
- 25 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:132;
- 30 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:132;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
- 35 above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

5 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:131.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:131 from nucleotide 914 to nucleotide 2353; the nucleotide sequence of SEQ ID NO:131 from
10 nucleotide 1793 to nucleotide 2353; the nucleotide sequence of SEQ ID NO:131 from nucleotide 1037 to nucleotide 1260; the nucleotide sequence of the full-length protein coding sequence of clone CT748_2 deposited with the ATCC under accession number 98271; or the nucleotide sequence of a mature protein coding sequence of clone CT748_2 deposited with the ATCC under accession number 98271. In other preferred embodiments, the polynucleotide encodes the full-
15 length or a mature protein encoded by the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:132 from amino acid 22 to amino acid 116. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid
20 sequence of SEQ ID NO:132 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:132, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising the amino acid sequence from amino acid 234 to amino acid 243 of SEQ ID NO:132.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:131.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131; and

- (ab) the nucleotide sequence of the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 10 consisting of:
- (ba) SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131; and
- (bb) the nucleotide sequence of the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide
- 20 sequence corresponding to the cDNA sequence of SEQ ID NO:131, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:131 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:131, but excluding the poly(A) tail at the 3' end of SEQ ID NO:131. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:131
- 25 from nucleotide 914 to nucleotide 2353, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:131 from nucleotide 914 to nucleotide 2353, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:131 from nucleotide 914 to nucleotide 2353. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of
- 30 SEQ ID NO:131 from nucleotide 1793 to nucleotide 2353, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:131 from nucleotide 1793 to nucleotide 2353, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:131 from nucleotide 1793 to nucleotide 2353. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence
- 35 corresponding to the cDNA sequence of SEQ ID NO:131 from nucleotide 1037 to nucleotide

1260, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:131 from nucleotide 1037 to nucleotide 1260, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:131 from nucleotide 1037 to nucleotide 1260.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:132;
- (b) the amino acid sequence of SEQ ID NO:132 from amino acid 22 to amino acid 116;
- 10 (c) a fragment of the amino acid sequence of SEQ ID NO:132, the fragment comprising eight contiguous amino acids of SEQ ID NO:132; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CT748_2 deposited with the ATCC under accession number 98271;

the protein being substantially free from other mammalian proteins. Preferably such protein
15 comprises the amino acid sequence of SEQ ID NO:132 or the amino acid sequence of SEQ ID NO:132 from amino acid 22 to amino acid 116. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:132, or a protein
20 comprising a fragment of the amino acid sequence of SEQ ID NO:132 having biological activity, the fragment comprising the amino acid sequence from amino acid 234 to amino acid 243 of SEQ ID NO:132.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133 from nucleotide 22 to nucleotide 462;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AJ1_1 deposited with the ATCC under accession
30 number 98278;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AJ1_1 deposited with the ATCC under accession number
35 98278;

-165-

- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:134;
- 5 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:134;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- 10 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:133.
- 15

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:133 from nucleotide 22 to nucleotide 462; the nucleotide sequence of the full-length protein coding sequence of clone AJ1_1 deposited with the ATCC under accession number 98278; or the

20 nucleotide sequence of a mature protein coding sequence of clone AJ1_1 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of

25 SEQ ID NO:134 from amino acid 52 to amino acid 147. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:134, or a polynucleotide encoding a protein comprising a fragment of the amino acid

30 sequence of SEQ ID NO:134 having biological activity, the fragment comprising the amino acid sequence from amino acid 68 to amino acid 77 of SEQ ID NO:134.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:133 and SEQ ID NO:135.

Further embodiments of the invention provide isolated polynucleotides produced

35 according to a process selected from the group consisting of:

-166-

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5

(aa) SEQ ID NO:133;

(ab) SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135; and

(ac) the nucleotide sequence of the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278;

10

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

15

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:133;

(bb) SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135; and

20

(bc) the nucleotide sequence of the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

25

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:133 and SEQ ID NO:135, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:133 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:135, but excluding the poly(A) tail at the 3' end of SEQ ID NO:135. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:133, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:133 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:133. Also preferably the polynucleotide isolated according to the above

30
35

process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:133 from nucleotide 22 to nucleotide 462, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:133 from nucleotide 22 to nucleotide 462, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:133 from nucleotide 22 to nucleotide 462.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:134;
 - (b) the amino acid sequence of SEQ ID NO:134 from amino acid 52 to amino acid 147;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:134, the fragment comprising eight contiguous amino acids of SEQ ID NO:134; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone AJ1_1 deposited with the ATCC under accession number 98278;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:134 or the amino acid sequence of SEQ ID NO:134 from amino acid 52 to amino acid 147. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:134, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising the amino acid sequence from amino acid 68 to amino acid 77 of SEQ ID NO:134.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:136;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:136 from nucleotide 7 to nucleotide 1647;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:136 from nucleotide 1 to nucleotide 305;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AQ73_3 deposited with the ATCC under accession number 98278;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AQ73_3 deposited with the ATCC under accession number 98278;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:137;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:137 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:137;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:136.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:136 from nucleotide 7 to nucleotide 1647; the nucleotide sequence of SEQ ID NO:136 from nucleotide 1 to nucleotide 305; the nucleotide sequence of the full-length protein coding sequence of clone AQ73_3 deposited with the ATCC under accession number 98278; or the nucleotide sequence of a mature protein coding sequence of clone AQ73_3 deposited with the ATCC under
- 25 accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:137 from amino acid 1 to amino acid 68. In further preferred embodiments, the present
- 30 invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:137 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:137, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:137 having biological activity, the fragment comprising the amino acid sequence from
- 35 amino acid 268 to amino acid 277 of SEQ ID NO:137.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:136.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (aa) SEQ ID NO:136, but excluding the poly(A) tail at the 3' end of SEQ ID NO:136; and
- (ab) the nucleotide sequence of the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 15 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 20 consisting of:
- (ba) SEQ ID NO:136, but excluding the poly(A) tail at the 3' end of SEQ ID NO:136; and
- (bb) the nucleotide sequence of the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278;
- 25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide

30 sequence corresponding to the cDNA sequence of SEQ ID NO:136, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:136 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:136, but excluding the poly(A) tail at the 3' end of SEQ ID NO:136. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:136

35 from nucleotide 7 to nucleotide 1647, and extending contiguously from a nucleotide sequence

corresponding to the 5' end of said sequence of SEQ ID NO:136 from nucleotide 7 to nucleotide 1647, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:136 from nucleotide 7 to nucleotide 1647. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ
5 ID NO:136 from nucleotide 1 to nucleotide 305, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:136 from nucleotide 1 to nucleotide 305, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:136 from nucleotide 1 to nucleotide 305.

In other embodiments, the present invention provides a composition comprising a protein,
10 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:137;
- (b) the amino acid sequence of SEQ ID NO:137 from amino acid 1 to amino acid 68;
- (c) a fragment of the amino acid sequence of SEQ ID NO:137, the fragment
15 comprising eight contiguous amino acids of SEQ ID NO:137; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AQ73_3 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:137 or the amino acid sequence of SEQ ID
20 NO:137 from amino acid 1 to amino acid 68. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:137 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:137, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:137 having biological activity,
25 the fragment comprising the amino acid sequence from amino acid 268 to amino acid 277 of SEQ ID NO:137.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:138;
- 30 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:138 from nucleotide 62 to nucleotide 757;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:138 from nucleotide 357 to nucleotide 703;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG142_1 deposited with the ATCC under accession number 98278;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG142_1 deposited with the ATCC under accession number 98278;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:139;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:139 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:139;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 20 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:138.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:138 from nucleotide 62 to nucleotide 757; the nucleotide sequence of SEQ ID NO:138 from nucleotide 357 to nucleotide 703; the nucleotide sequence of the full-length protein coding sequence of clone BG142_1 deposited with the ATCC under accession number 98278; or the nucleotide sequence of a mature protein coding sequence of clone BG142_1 deposited with the
- 30 ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:139 from amino acid 184 to amino acid 214. In further preferred
- 35 embodiments, the present invention provides a polynucleotide encoding a protein comprising a

-172-

fragment of the amino acid sequence of SEQ ID NO:139 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:139, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:139 having biological activity, the fragment comprising the amino acid sequence from amino acid 111 to amino acid 120 of SEQ ID NO:139.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:138.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 15 (aa) SEQ ID NO:138, but excluding the poly(A) tail at the 3' end of SEQ ID NO:138; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 25 (ba) SEQ ID NO:138, but excluding the poly(A) tail at the 3' end of SEQ ID NO:138; and
 - (bb) the nucleotide sequence of the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278;
 - 30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:138, and extending contiguously

from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:138 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:138, but excluding the poly(A) tail at the 3' end of SEQ ID NO:138. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:138
5 from nucleotide 62 to nucleotide 757, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:138 from nucleotide 62 to nucleotide 757, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:138 from nucleotide 62 to nucleotide 757. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ
10 ID NO:138 from nucleotide 357 to nucleotide 703, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:138 from nucleotide 357 to nucleotide 703, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:138 from nucleotide 357 to nucleotide 703.

In other embodiments, the present invention provides a composition comprising a protein,
15 wherein said protein comprises an amino acid sequence selected from the group consisting of:
(a) the amino acid sequence of SEQ ID NO:139;
(b) the amino acid sequence of SEQ ID NO:139 from amino acid 184 to amino acid 214;
(c) a fragment of the amino acid sequence of SEQ ID NO:139, the fragment
20 comprising eight contiguous amino acids of SEQ ID NO:139; and
(d) the amino acid sequence encoded by the cDNA insert of clone BG142_1 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:139 or the amino acid sequence of SEQ ID
25 NO:139 from amino acid 184 to amino acid 214. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:139 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:139, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:139 having biological activity,
30 the fragment comprising the amino acid sequence from amino acid 111 to amino acid 120 of SEQ ID NO:139.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:140;

-174-

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:140 from nucleotide 404 to nucleotide 535;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:140 from nucleotide 1 to nucleotide 666;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BV66_1 deposited with the ATCC under accession number 98278;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BV66_1 deposited with the ATCC under accession number 98278;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BV66_1 deposited with the ATCC under accession number 98278;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BV66_1 deposited with the ATCC under accession number 98278;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:141;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:141 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:141;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:140.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:140
- 30 from nucleotide 404 to nucleotide 535; the nucleotide sequence of SEQ ID NO:140 from nucleotide 1 to nucleotide 666; the nucleotide sequence of the full-length protein coding sequence of clone BV66_1 deposited with the ATCC under accession number 98278; or the nucleotide sequence of a mature protein coding sequence of clone BV66_1 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-
- 35 length or a mature protein encoded by the cDNA insert of clone BV66_1 deposited with the

ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:141 from amino acid 1 to amino acid 38. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid
5 sequence of SEQ ID NO:141 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:141, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:141 having biological activity, the fragment comprising the amino acid sequence from amino acid 17 to amino acid 26 of SEQ ID NO:141.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:140.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - 15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:140, but excluding the poly(A) tail at the 3' end of SEQ ID NO:140; and
 - 20 (ab) the nucleotide sequence of the cDNA insert of clone BV66_1 deposited with the ATCC under accession number 98278;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
 - 25 and
 - (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 30 (ba) SEQ ID NO:140, but excluding the poly(A) tail at the 3' end of SEQ ID NO:140; and
 - (bb) the nucleotide sequence of the cDNA insert of clone BV66_1 deposited with the ATCC under accession number 98278;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions
 - 35 at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:140, and extending contiguously
5 from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:140 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:140, but excluding the poly(A) tail at the 3' end of SEQ ID NO:140. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:140 from nucleotide 404 to nucleotide 535, and extending contiguously from a nucleotide sequence
10 corresponding to the 5' end of said sequence of SEQ ID NO:140 from nucleotide 404 to nucleotide 535, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:140 from nucleotide 404 to nucleotide 535. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:140 from nucleotide 1 to nucleotide 666, and extending contiguously from a nucleotide
15 sequence corresponding to the 5' end of said sequence of SEQ ID NO:140 from nucleotide 1 to nucleotide 666, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:140 from nucleotide 1 to nucleotide 666.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:141;
- (b) the amino acid sequence of SEQ ID NO:141 from amino acid 1 to amino acid 38;
- (c) a fragment of the amino acid sequence of SEQ ID NO:141, the fragment comprising eight contiguous amino acids of SEQ ID NO:141; and
- 25 (d) the amino acid sequence encoded by the cDNA insert of clone BV66_1 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:141 or the amino acid sequence of SEQ ID NO:141 from amino acid 1 to amino acid 38. In further preferred embodiments, the present
30 invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:141 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:141, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:141 having biological activity, the fragment comprising the amino acid sequence from amino acid 17 to amino acid 26 of SEQ
35 ID NO:141.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:142;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:142
5 from nucleotide 1204 to nucleotide 1389;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:142
from nucleotide 881 to nucleotide 1380;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone BV291_3 deposited with the ATCC under accession
10 number 98278;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone BV291_3 deposited with the ATCC under accession number 98278;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone BV291_3 deposited with the ATCC under accession number
15 98278;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone BV291_3 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:143;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino
20 acid sequence of SEQ ID NO:143 having biological activity, the fragment comprising
eight contiguous amino acids of SEQ ID NO:143;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g)
above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h)
25 or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
length of SEQ ID NO:142.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:142
from nucleotide 1204 to nucleotide 1389; the nucleotide sequence of SEQ ID NO:142 from
nucleotide 881 to nucleotide 1380; the nucleotide sequence of the full-length protein coding
35 sequence of clone BV291_3 deposited with the ATCC under accession number 98278; or the

nucleotide sequence of a mature protein coding sequence of clone BV291_3 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BV291_3 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, 5 the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:143 from amino acid 1 to amino acid 59. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:143 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino 10 acids of SEQ ID NO:143, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:143 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:143.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:142.

15 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 20 (aa) SEQ ID NO:142, but excluding the poly(A) tail at the 3' end of SEQ ID NO:142; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BV291_3 deposited with the ATCC under accession number 98278;
 - 25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - 30 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:142, but excluding the poly(A) tail at the 3' end of SEQ ID NO:142; and

-179-

- (bb) the nucleotide sequence of the cDNA insert of clone BV291_3 deposited with the ATCC under accession number 98278;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:142, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:142 to a nucleotide
10 sequence corresponding to the 3' end of SEQ ID NO:142, but excluding the poly(A) tail at the 3' end of SEQ ID NO:142. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:142 from nucleotide 1204 to nucleotide 1389, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:142 from nucleotide 1204 to
15 nucleotide 1389, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:142 from nucleotide 1204 to nucleotide 1389. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:142 from nucleotide 881 to nucleotide 1380, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:142 from
20 nucleotide 881 to nucleotide 1380, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:142 from nucleotide 881 to nucleotide 1380.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:143;
- 25 (b) the amino acid sequence of SEQ ID NO:143 from amino acid 1 to amino acid 59;
- (c) a fragment of the amino acid sequence of SEQ ID NO:143, the fragment comprising eight contiguous amino acids of SEQ ID NO:143; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BV291_3
30 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:143 or the amino acid sequence of SEQ ID NO:143 from amino acid 1 to amino acid 59. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID
35 NO:143 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:143, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:143 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:143.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:144;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:144 from nucleotide 189 to nucleotide 1115;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:144 from nucleotide 1 to nucleotide 451;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CK201_1 deposited with the ATCC under accession number 98278;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CK201_1 deposited with the ATCC under accession number 98278;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CK201_1 deposited with the ATCC under accession number 98278;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CK201_1 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:145;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:145 having biological activity, the fragment comprising
25 eight contiguous amino acids of SEQ ID NO:145;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
35 length of SEQ ID NO:144.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:144 from nucleotide 189 to nucleotide 1115; the nucleotide sequence of SEQ ID NO:144 from nucleotide 1 to nucleotide 451; the nucleotide sequence of the full-length protein coding sequence of clone CK201_1 deposited with the ATCC under accession number 98278; or the nucleotide
5 sequence of a mature protein coding sequence of clone CK201_1 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CK201_1 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID
10 NO:145 from amino acid 1 to amino acid 88. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:145 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:145, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
15 ID NO:145 having biological activity, the fragment comprising the amino acid sequence from amino acid 149 to amino acid 158 of SEQ ID NO:145.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:144.

Further embodiments of the invention provide isolated polynucleotides produced
20 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
25 (aa) SEQ ID NO:144, but excluding the poly(A) tail at the 3' end of SEQ ID NO:144; and
(ab) the nucleotide sequence of the cDNA insert of clone CK201_1 deposited with the ATCC under accession number 98278;
(ii) hybridizing said probe(s) to human genomic DNA in conditions
30 at least as stringent as 4X SSC at 50 degrees C; and
(iii) isolating the DNA polynucleotides detected with the probe(s);
and
- (b) a process comprising the steps of:

-182-

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:144, but excluding the poly(A) tail at the 3' end of SEQ ID NO:144; and

(bb) the nucleotide sequence of the cDNA insert of clone CK201_1 deposited with the ATCC under accession number 98278;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

10 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:144, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:144 to a nucleotide
15 sequence corresponding to the 3' end of SEQ ID NO:144, but excluding the poly(A) tail at the 3' end of SEQ ID NO:144. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:144 from nucleotide 189 to nucleotide 1115, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:144 from nucleotide 189 to nucleotide
20 1115, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:144 from nucleotide 189 to nucleotide 1115. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:144 from nucleotide 1 to nucleotide 451, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:144 from
25 nucleotide 1 to nucleotide 451, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:144 from nucleotide 1 to nucleotide 451.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:145;
30 (b) the amino acid sequence of SEQ ID NO:145 from amino acid 1 to amino acid 88;

(c) a fragment of the amino acid sequence of SEQ ID NO:145, the fragment comprising eight contiguous amino acids of SEQ ID NO:145; and

(d) the amino acid sequence encoded by the cDNA insert of clone CK201_1
35 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:145 or the amino acid sequence of SEQ ID NO:145 from amino acid 1 to amino acid 88. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:145 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:145, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:145 having biological activity, the fragment comprising the amino acid sequence from amino acid 149 to amino acid 158 of SEQ ID NO:145.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:146;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:146 from nucleotide 117 to nucleotide 923;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:146 from nucleotide 174 to nucleotide 923;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:146 from nucleotide 1 to nucleotide 316;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CQ331_2 deposited with the ATCC under accession number 98278;
- 20 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CQ331_2 deposited with the ATCC under accession number 98278;
- 25 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:147;
- 30 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:147 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:147;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
- 35 above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

5 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:146.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:146 from nucleotide 117 to nucleotide 923; the nucleotide sequence of SEQ ID NO:146 from
10 nucleotide 174 to nucleotide 923; the nucleotide sequence of SEQ ID NO:146 from nucleotide 1 to nucleotide 316; the nucleotide sequence of the full-length protein coding sequence of clone CQ331_2 deposited with the ATCC under accession number 98278; or the nucleotide sequence of a mature protein coding sequence of clone CQ331_2 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a
15 mature protein encoded by the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:147 from amino acid 1 to amino acid 57. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
20 ID NO:147 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:147, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:147 having biological activity, the fragment comprising the amino acid sequence from amino acid 129 to amino acid 138 of SEQ ID NO:147.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:146.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:
30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:146, but excluding the poly(A) tail at the 3' end of SEQ ID NO:146; and

- (ab) the nucleotide sequence of the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (ba) SEQ ID NO:146, but excluding the poly(A) tail at the 3' end of SEQ ID NO:146; and
- (bb) the nucleotide sequence of the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide
- 20 sequence corresponding to the cDNA sequence of SEQ ID NO:146, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:146 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:146, but excluding the poly(A) tail at the 3' end of SEQ ID NO:146. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:146
- 25 from nucleotide 117 to nucleotide 923, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:146 from nucleotide 117 to nucleotide 923, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:146 from nucleotide 117 to nucleotide 923. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ
- 30 ID NO:146 from nucleotide 174 to nucleotide 923, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:146 from nucleotide 174 to nucleotide 923, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:146 from nucleotide 174 to nucleotide 923. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
- 35 sequence of SEQ ID NO:146 from nucleotide 1 to nucleotide 316, and extending contiguously

-186-

from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:146 from nucleotide 1 to nucleotide 316, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:146 from nucleotide 1 to nucleotide 316.

In other embodiments, the present invention provides a composition comprising a protein,
5 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:147;
- (b) the amino acid sequence of SEQ ID NO:147 from amino acid 1 to amino acid 57;
- (c) a fragment of the amino acid sequence of SEQ ID NO:147, the fragment
10 comprising eight contiguous amino acids of SEQ ID NO:147; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CQ331_2 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:147 or the amino acid sequence of SEQ ID
15 NO:147 from amino acid 1 to amino acid 57. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:147 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:147, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:147 having biological activity,
20 the fragment comprising the amino acid sequence from amino acid 129 to amino acid 138 of SEQ ID NO:147.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:148;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:148 from nucleotide 223 to nucleotide 483;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:148 from nucleotide 22 to nucleotide 397;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
30 protein coding sequence of clone CT550_1 deposited with the ATCC under accession number 98278;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CT550_1 deposited with the ATCC under accession number 98278;
- 5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:149;
- 10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:149 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:149;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 15 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:148.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:148 from nucleotide 223 to nucleotide 483; the nucleotide sequence of SEQ ID NO:148 from nucleotide 22 to nucleotide 397; the nucleotide sequence of the full-length protein coding sequence of clone CT550_1 deposited with the ATCC under accession number 98278; or the nucleotide sequence of a mature protein coding sequence of clone CT550_1 deposited with the
- 25 ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:149 from amino acid 1 to amino acid 58. In further preferred
- 30 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:149 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:149, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:149 having biological activity, the fragment comprising the
- 35 amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:149.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:148.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:148, but excluding the poly(A) tail at the 3'
- 10 end of SEQ ID NO:148; and
- (ab) the nucleotide sequence of the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 15 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 20 consisting of:
- (ba) SEQ ID NO:148, but excluding the poly(A) tail at the 3' end of SEQ ID NO:148; and
- (bb) the nucleotide sequence of the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278;
- 25 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide

30 sequence corresponding to the cDNA sequence of SEQ ID NO:148, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:148 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:148, but excluding the poly(A) tail at the 3' end of SEQ ID NO:148. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:148

35 from nucleotide 223 to nucleotide 483, and extending contiguously from a nucleotide sequence

corresponding to the 5' end of said sequence of SEQ ID NO:148 from nucleotide 223 to nucleotide 483, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:148 from nucleotide 223 to nucleotide 483. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:148 from nucleotide 22 to nucleotide 397, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:148 from nucleotide 22 to nucleotide 397, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:148 from nucleotide 22 to nucleotide 397.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:149;
- (b) the amino acid sequence of SEQ ID NO:149 from amino acid 1 to amino acid 58;
- (c) a fragment of the amino acid sequence of SEQ ID NO:149, the fragment comprising eight contiguous amino acids of SEQ ID NO:149; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CT550_1 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:149 or the amino acid sequence of SEQ ID NO:149 from amino acid 1 to amino acid 58. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:149 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:149, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:149 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:149.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:150;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:150 from nucleotide 112 to nucleotide 969;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:150 from nucleotide 154 to nucleotide 969;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:150 from nucleotide 1 to nucleotide 423;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CT585_1 deposited with the ATCC under accession number 98278;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CT585_1 deposited with the ATCC under accession number 98278;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:151;
- 15 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:151 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:151;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:150.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:150 from nucleotide 112 to nucleotide 969; the nucleotide sequence of SEQ ID NO:150 from nucleotide 154 to nucleotide 969; the nucleotide sequence of SEQ ID NO:150 from nucleotide 1 to nucleotide 423; the nucleotide sequence of the full-length protein coding sequence of clone CT585_1 deposited with the ATCC under accession number 98278; or the nucleotide sequence
- 30 of a mature protein coding sequence of clone CT585_1 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:151 from
- 35 amino acid 1 to amino acid 104. In further preferred embodiments, the present invention provides

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:151 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:151, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:151 having biological activity, the fragment comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:151.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:150.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:150, but excluding the poly(A) tail at the 3' end of SEQ ID NO:150; and

- (ab) the nucleotide sequence of the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278;

- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:150, but excluding the poly(A) tail at the 3' end of SEQ ID NO:150; and

- (bb) the nucleotide sequence of the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and

- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:150, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:150 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:150, but excluding the poly(A) tail at the 3' end of SEQ ID NO:150. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:150 from nucleotide 112 to nucleotide 969, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:150 from nucleotide 112 to nucleotide 969, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:150 from nucleotide 112 to nucleotide 969. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:150 from nucleotide 154 to nucleotide 969, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:150 from nucleotide 154 to nucleotide 969, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:150 from nucleotide 154 to nucleotide 969. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:150 from nucleotide 1 to nucleotide 423, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:150 from nucleotide 1 to nucleotide 423, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:150 from nucleotide 1 to nucleotide 423.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:151;
 - (b) the amino acid sequence of SEQ ID NO:151 from amino acid 1 to amino acid 104;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:151, the fragment comprising eight contiguous amino acids of SEQ ID NO:151; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone CT585_1 deposited with the ATCC under accession number 98278;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:151 or the amino acid sequence of SEQ ID NO:151 from amino acid 1 to amino acid 104. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:151 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:151, or a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:151 having biological activity, the fragment comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:151.

In one embodiment, the present invention provides a composition comprising an isolated
5 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:152;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:152
from nucleotide 37 to nucleotide 2766;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:152
10 from nucleotide 243 to nucleotide 789;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone CT797_3 deposited with the ATCC under accession
number 98278;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
15 insert of clone CT797_3 deposited with the ATCC under accession number 98278;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone CT797_3 deposited with the ATCC under accession number
98278;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
20 of clone CT797_3 deposited with the ATCC under accession number 98278;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:153;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:153 having biological activity, the fragment comprising
25 eight contiguous amino acids of SEQ ID NO:153;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g)
above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h)
or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
length of SEQ ID NO:152.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:152 from nucleotide 37 to nucleotide 2766; the nucleotide sequence of SEQ ID NO:152 from nucleotide 243 to nucleotide 789; the nucleotide sequence of the full-length protein coding sequence of clone CT797_3 deposited with the ATCC under accession number 98278; or the
5 nucleotide sequence of a mature protein coding sequence of clone CT797_3 deposited with the ATCC under accession number 98278. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CT797_3 deposited with the ATCC under accession number 98278. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid
10 sequence of SEQ ID NO:153 from amino acid 75 to amino acid 251. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:153 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:153, or a polynucleotide encoding a protein comprising a fragment of the
15 amino acid sequence of SEQ ID NO:153 having biological activity, the fragment comprising the amino acid sequence from amino acid 450 to amino acid 459 of SEQ ID NO:153.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:152.

Further embodiments of the invention provide isolated polynucleotides produced
20 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
25 (aa) SEQ ID NO:152, but excluding the poly(A) tail at the 3' end of SEQ ID NO:152; and
(ab) the nucleotide sequence of the cDNA insert of clone CT797_3 deposited with the ATCC under accession number 98278;
(ii) hybridizing said probe(s) to human genomic DNA in conditions
30 at least as stringent as 4X SSC at 50 degrees C; and
(iii) isolating the DNA polynucleotides detected with the probe(s);
and
- (b) a process comprising the steps of:

-195-

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:152, but excluding the poly(A) tail at the 3' end of SEQ ID NO:152; and

(bb) the nucleotide sequence of the cDNA insert of clone CT797_3 deposited with the ATCC under accession number 98278;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

10 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:152, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:152 to a nucleotide
15 sequence corresponding to the 3' end of SEQ ID NO:152, but excluding the poly(A) tail at the 3' end of SEQ ID NO:152. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:152 from nucleotide 37 to nucleotide 2766, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:152 from nucleotide 37 to nucleotide
20 2766, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:152 from nucleotide 37 to nucleotide 2766. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:152 from nucleotide 243 to nucleotide 789, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:152 from nucleotide 243
25 to nucleotide 789, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:152 from nucleotide 243 to nucleotide 789.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:153;
30 (b) the amino acid sequence of SEQ ID NO:153 from amino acid 75 to amino acid 251;

(c) a fragment of the amino acid sequence of SEQ ID NO:153, the fragment comprising eight contiguous amino acids of SEQ ID NO:153; and

(d) the amino acid sequence encoded by the cDNA insert of clone CT797_3
35 deposited with the ATCC under accession number 98278;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:153 or the amino acid sequence of SEQ ID NO:153 from amino acid 75 to amino acid 251. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:153 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:153, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:153 having biological activity, the fragment comprising the amino acid sequence from amino acid 450 to amino acid 459 of SEQ ID NO:153.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:155;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:155 from nucleotide 41 to nucleotide 760;
- 15 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CB107_1 deposited with the ATCC under accession number 98279;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279;
- 20 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CB107_1 deposited with the ATCC under accession number 98279;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279;
- 25 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:156;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:156;
- 30 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k) a polynucleotide that hybridizes under stringent conditions to any one of
- 35 the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:155.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:155 from nucleotide 41 to nucleotide 760; the nucleotide sequence of the full-length protein coding sequence of clone CB107_1 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CB107_1 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:156 from amino acid 127 to amino acid 240. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:156, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:156.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:155, SEQ ID NO:154, and SEQ ID NO:157.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:154;
 - (ab) SEQ ID NO:155;
 - (ac) SEQ ID NO:157, but excluding the poly(A) tail at the 3' end of SEQ ID NO:157; and
 - (ad) the nucleotide sequence of the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:154;

(bb) SEQ ID NO:155;

(bc) SEQ ID NO:157, but excluding the poly(A) tail at the 3' end of SEQ ID NO:157; and

(bd) the nucleotide sequence of the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:154, SEQ ID NO:155, and SEQ ID NO:157, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:154 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:157, but excluding the poly(A) tail at the 3' end of SEQ ID NO:157. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:155, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:155 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:155. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:155 from nucleotide 41 to nucleotide 760, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:155 from nucleotide 41 to nucleotide 760, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:155 from nucleotide 41 to nucleotide 760.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:156;

(b) the amino acid sequence of SEQ ID NO:156 from amino acid 127 to amino acid 240;

- (c) a fragment of the amino acid sequence of SEQ ID NO:156, the fragment comprising eight contiguous amino acids of SEQ ID NO:156; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CB107_1 deposited with the ATCC under accession number 98279;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:156 or the amino acid sequence of SEQ ID NO:156 from amino acid 127 to amino acid 240. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment preferably comprising eight (more preferably
- 10 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:156, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:156 having biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:156.

In one embodiment, the present invention provides a composition comprising an isolated

15 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:158;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:158 from nucleotide 374 to nucleotide 1108;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:158
- 20 from nucleotide 500 to nucleotide 1108;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:158 from nucleotide 1 to nucleotide 387;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CG300_3 deposited with the ATCC under accession
- 25 number 98279;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CG300_3 deposited with the ATCC under accession number
- 30 98279;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:159;

(j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:159 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:159;

5 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

10 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:158.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:158 from nucleotide 374 to nucleotide 1108; the nucleotide sequence of SEQ ID NO:158 from
15 nucleotide 500 to nucleotide 1108; the nucleotide sequence of SEQ ID NO:158 from nucleotide 1 to nucleotide 387; the nucleotide sequence of the full-length protein coding sequence of clone CG300_3 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CG300_3 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a
20 mature protein encoded by the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:159 from amino acid 23 to amino acid 57. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
25 ID NO:159 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:159, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:159 having biological activity, the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:159.

30 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:158.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

-201-

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:158, but excluding the poly(A) tail at the 3' end of SEQ ID NO:158; and

(ab) the nucleotide sequence of the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:158, but excluding the poly(A) tail at the 3' end of SEQ ID NO:158; and

(bb) the nucleotide sequence of the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:158, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:158 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:158, but excluding the poly(A) tail at the 3' end of SEQ ID NO:158. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:158 from nucleotide 374 to nucleotide 1108, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:158 from nucleotide 374 to nucleotide 1108, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:158 from nucleotide 374 to nucleotide 1108. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:158 from nucleotide 500 to nucleotide 1108, and extending contiguously from a

nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:158 from nucleotide 500 to nucleotide 1108, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:158 from nucleotide 500 to nucleotide 1108. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence
5 corresponding to the cDNA sequence of SEQ ID NO:158 from nucleotide 1 to nucleotide 387, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:158 from nucleotide 1 to nucleotide 387, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:158 from nucleotide 1 to nucleotide 387.

In other embodiments, the present invention provides a composition comprising a protein,
10 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:159;
- (b) the amino acid sequence of SEQ ID NO:159 from amino acid 23 to amino acid 57;
- (c) a fragment of the amino acid sequence of SEQ ID NO:159, the fragment
15 comprising eight contiguous amino acids of SEQ ID NO:159; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CG300_3 deposited with the ATCC under accession number 98279;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:159 or the amino acid sequence of SEQ ID
20 NO:159 from amino acid 23 to amino acid 57. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:159 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:159, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:159 having biological activity,
25 the fragment comprising the amino acid sequence from amino acid 117 to amino acid 126 of SEQ ID NO:159.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:160;
- 30 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:160 from nucleotide 126 to nucleotide 3053;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:160 from nucleotide 180 to nucleotide 3053;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:160
35 from nucleotide 49 to nucleotide 382;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CJ145_1 deposited with the ATCC under accession number 98279;
- 5 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CJ145_1 deposited with the ATCC under accession number 98279;
- 10 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:161;
- 15 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:161 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:161;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 20 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:160.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:160 from nucleotide 126 to nucleotide 3053; the nucleotide sequence of SEQ ID NO:160 from nucleotide 180 to nucleotide 3053; the nucleotide sequence of SEQ ID NO:160 from nucleotide 49 to nucleotide 382; the nucleotide sequence of the full-length protein coding sequence of clone CJ145_1 deposited with the ATCC under accession number 98279; or the nucleotide sequence of
- 30 a mature protein coding sequence of clone CJ145_1 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:161 from
- 35 amino acid 1 to amino acid 87. In further preferred embodiments, the present invention provides

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:161 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:161, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:161 having
5 biological activity, the fragment comprising the amino acid sequence from amino acid 482 to amino acid 491 of SEQ ID NO:161.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:160.

Further embodiments of the invention provide isolated polynucleotides produced
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:160, but excluding the poly(A) tail at the 3' end of SEQ ID NO:160; and

(ab) the nucleotide sequence of the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:160, but excluding the poly(A) tail at the 3' end of SEQ ID NO:160; and

30 (bb) the nucleotide sequence of the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:160, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:160 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:160, but excluding the poly(A) tail at the 3' end of SEQ ID NO:160. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:160 from nucleotide 126 to nucleotide 3053, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:160 from nucleotide 126 to nucleotide 3053, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:160 from nucleotide 126 to nucleotide 3053. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:160 from nucleotide 180 to nucleotide 3053, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:160 from nucleotide 180 to nucleotide 3053, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:160 from nucleotide 180 to nucleotide 3053. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:160 from nucleotide 49 to nucleotide 382, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:160 from nucleotide 49 to nucleotide 382, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:160 from nucleotide 49 to nucleotide 382.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:161;
- (b) the amino acid sequence of SEQ ID NO:161 from amino acid 1 to amino acid 87;
- (c) a fragment of the amino acid sequence of SEQ ID NO:161, the fragment comprising eight contiguous amino acids of SEQ ID NO:161; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CJ145_1 deposited with the ATCC under accession number 98279;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:161 or the amino acid sequence of SEQ ID NO:161 from amino acid 1 to amino acid 87. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:161 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:161, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:161 having biological activity, the fragment comprising the amino acid sequence from amino acid 482 to amino acid 491 of SEQ ID NO:161.

- 5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:162;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:162
from nucleotide 40 to nucleotide 342;
 - 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:162
from nucleotide 127 to nucleotide 342;
 - (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:162
from nucleotide 11 to nucleotide 181;
 - (e) a polynucleotide comprising the nucleotide sequence of the full-length
15 protein coding sequence of clone CJ160_11 deposited with the ATCC under accession
number 98279;
 - (f) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone CJ160_11 deposited with the ATCC under accession number 98279;
 - (g) a polynucleotide comprising the nucleotide sequence of a mature protein
20 coding sequence of clone CJ160_11 deposited with the ATCC under accession number
98279;
 - (h) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone CJ160_11 deposited with the ATCC under accession number 98279;
 - (i) a polynucleotide encoding a protein comprising the amino acid sequence
25 of SEQ ID NO:163;
 - (j) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:163 having biological activity, the fragment comprising
eight contiguous amino acids of SEQ ID NO:163;
 - (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
30 above;
 - (l) a polynucleotide which encodes a species homologue of the protein of (i)
or (j) above ;
 - (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j); and

(n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:162.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:162 from nucleotide 40 to nucleotide 342; the nucleotide sequence of SEQ ID NO:162 from nucleotide 127 to nucleotide 342; the nucleotide sequence of SEQ ID NO:162 from nucleotide 11 to nucleotide 181; the nucleotide sequence of the full-length protein coding sequence of clone CJ160_11 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CJ160_11 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CJ160_11 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:163 from amino acid 7 to amino acid 48. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:163 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:163, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:163 having biological activity, the fragment comprising the amino acid sequence from amino acid 45 to amino acid 54 of SEQ ID NO:163.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:162.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:162, but excluding the poly(A) tail at the 3' end of SEQ ID NO:162; and
 - (ab) the nucleotide sequence of the cDNA insert of clone CJ160_11 deposited with the ATCC under accession number 98279;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:162, but excluding the poly(A) tail at the 3' end of SEQ ID NO:162; and
 - (bb) the nucleotide sequence of the cDNA insert of clone CJ160_11 deposited with the ATCC under accession number 98279;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - (iii) amplifying human DNA sequences; and
 - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:162, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:162 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:162, but excluding the poly(A) tail at the 3' end of SEQ ID NO:162. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:162 from nucleotide 40 to nucleotide 342, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:162 from nucleotide 40 to nucleotide 342, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:162 from nucleotide 40 to nucleotide 342. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:162 from nucleotide 127 to nucleotide 342, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:162 from nucleotide 127 to nucleotide 342, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:162 from nucleotide 127 to nucleotide 342. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:162 from nucleotide 11 to nucleotide 181, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:162 from nucleotide 11 to nucleotide 181, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:162 from nucleotide 11 to nucleotide 181.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:163;
- (b) the amino acid sequence of SEQ ID NO:163 from amino acid 7 to amino acid 48;
- (c) a fragment of the amino acid sequence of SEQ ID NO:163, the fragment comprising eight contiguous amino acids of SEQ ID NO:163; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CJ160_11 deposited with the ATCC under accession number 98279;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:163 or the amino acid sequence of SEQ ID NO:163 from amino acid 7 to amino acid 48. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:163 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:163, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:163 having biological activity, the fragment comprising the amino acid sequence from amino acid 45 to amino acid 54 of SEQ ID NO:163.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:164;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:164 from nucleotide 180 to nucleotide 467;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:164 from nucleotide 267 to nucleotide 467;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CO20_1 deposited with the ATCC under accession number 98279;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CO20_1 deposited with the ATCC under accession number 98279;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CO20_1 deposited with the ATCC under accession number 98279;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CO20_1 deposited with the ATCC under accession number 98279;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:165;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:165 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:165;
- 5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 10 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:164.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:164 from nucleotide 180 to nucleotide 467; the nucleotide sequence of SEQ ID NO:164 from
15 nucleotide 267 to nucleotide 467; the nucleotide sequence of the full-length protein coding sequence of clone CO20_1 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CO20_1 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CO20_1
20 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:165 from amino acid 1 to amino acid 37. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:165 having biological activity, the fragment
25 preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:165, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:165 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:165.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
30 NO:164 and SEQ ID NO:166.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:164;

(ab) SEQ ID NO:166, but excluding the poly(A) tail at the 3' end of SEQ ID NO:166; and

(ac) the nucleotide sequence of the cDNA insert of clone CO20_1 deposited with the ATCC under accession number 98279;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:164;

(bb) SEQ ID NO:166, but excluding the poly(A) tail at the 3' end of SEQ ID NO:166; and

(bc) the nucleotide sequence of the cDNA insert of clone CO20_1 deposited with the ATCC under accession number 98279;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:164 and SEQ ID NO:166, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:164 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:166, but excluding the poly(A) tail at the 3' end of SEQ ID NO:166. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:164, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:164 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:164. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:164

from nucleotide 180 to nucleotide 467, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:164 from nucleotide 180 to nucleotide 467, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:164 from nucleotide 180 to nucleotide 467. Also preferably the polynucleotide isolated according to
5 the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:164 from nucleotide 267 to nucleotide 467, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:164 from nucleotide 267 to nucleotide 467, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:164 from nucleotide 267 to nucleotide 467.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:165;
- (b) the amino acid sequence of SEQ ID NO:165 from amino acid 1 to amino acid 37;
- 15 (c) a fragment of the amino acid sequence of SEQ ID NO:165, the fragment comprising eight contiguous amino acids of SEQ ID NO:165; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CO20_1 deposited with the ATCC under accession number 98279;

the protein being substantially free from other mammalian proteins. Preferably such protein
20 comprises the amino acid sequence of SEQ ID NO:165 or the amino acid sequence of SEQ ID NO:165 from amino acid 1 to amino acid 37. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:165 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:165, or a protein
25 comprising a fragment of the amino acid sequence of SEQ ID NO:165 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:165.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:167;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:167 from nucleotide 176 to nucleotide 520;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:167 from nucleotide 317 to nucleotide 520;

- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:167 from nucleotide 118 to nucleotide 413;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CO223_3 deposited with the ATCC under accession number 98291;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CO223_3 deposited with the ATCC under accession number 98291;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CO223_3 deposited with the ATCC under accession number 98291;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CO223_3 deposited with the ATCC under accession number 98291;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:168;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:168 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:168;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:167.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:167 from nucleotide 176 to nucleotide 520; the nucleotide sequence of SEQ ID NO:167 from nucleotide 317 to nucleotide 520; the nucleotide sequence of SEQ ID NO:167 from nucleotide 118 to nucleotide 413; the nucleotide sequence of the full-length protein coding sequence of clone CO223_3 deposited with the ATCC under accession number 98291; or the nucleotide sequence of a mature protein coding sequence of clone CO223_3 deposited with the ATCC under accession number 98291. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CO223_3 deposited with the ATCC under accession number 98291. In yet other preferred embodiments, the present invention provides a

polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:168 from amino acid 1 to amino acid 80. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:168 having biological activity, the fragment preferably comprising eight (more preferably
5 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:168, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:168 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:168.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
10 NO:167.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X
15 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:167, but excluding the poly(A) tail at the 3' end of SEQ ID NO:167; and
 - (ab) the nucleotide sequence of the cDNA insert of clone
20 CO223_3 deposited with the ATCC under accession number 98291;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- 25 (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:167, but excluding the poly(A) tail at the 3'
30 end of SEQ ID NO:167; and
 - (bb) the nucleotide sequence of the cDNA insert of clone CO223_3 deposited with the ATCC under accession number 98291;
 - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
 - 35 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:167, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:167 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:167, but excluding the poly(A) tail at the 3' end of SEQ ID NO:167. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:167 from nucleotide 176 to nucleotide 520, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:167 from nucleotide 176 to nucleotide 520, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:167 from nucleotide 176 to nucleotide 520. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:167 from nucleotide 317 to nucleotide 520, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:167 from nucleotide 317 to nucleotide 520, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:167 from nucleotide 317 to nucleotide 520. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:167 from nucleotide 118 to nucleotide 413, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:167 from nucleotide 118 to nucleotide 413, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:167 from nucleotide 118 to nucleotide 413.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:168;
- (b) the amino acid sequence of SEQ ID NO:168 from amino acid 1 to amino acid 80;
- (c) a fragment of the amino acid sequence of SEQ ID NO:168, the fragment comprising eight contiguous amino acids of SEQ ID NO:168; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CO223_3 deposited with the ATCC under accession number 98291;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:168 or the amino acid sequence of SEQ ID NO:168 from amino acid 1 to amino acid 80. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:168 having biological activity, the fragment preferably comprising eight (more preferably

twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:168, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:168 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:168.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:169;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:169 from nucleotide 303 to nucleotide 542;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:169 from nucleotide 1 to nucleotide 435;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CO310_2 deposited with the ATCC under accession number 98279;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CO310_2 deposited with the ATCC under accession number 98279;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CO310_2 deposited with the ATCC under accession number 98279;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CO310_2 deposited with the ATCC under accession number 98279;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:170;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:170 having biological activity, the fragment comprising
25 eight contiguous amino acids of SEQ ID NO:170;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
35 length of SEQ ID NO:169.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:169 from nucleotide 303 to nucleotide 542; the nucleotide sequence of SEQ ID NO:169 from nucleotide 1 to nucleotide 435; the nucleotide sequence of the full-length protein coding sequence of clone CO310_2 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CO310_2 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CO310_2 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:170 from amino acid 1 to amino acid 44. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:170 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:170, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:170 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:170.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:169.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:169, but excluding the poly(A) tail at the 3' end of SEQ ID NO:169; and
 - (ab) the nucleotide sequence of the cDNA insert of clone CO310_2 deposited with the ATCC under accession number 98279;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:169, but excluding the poly(A) tail at the 3' end of SEQ ID NO:169; and
- (bb) the nucleotide sequence of the cDNA insert of clone CO310_2 deposited with the ATCC under accession number 98279;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 10 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:169, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:169 to a nucleotide

15 sequence corresponding to the 3' end of SEQ ID NO:169, but excluding the poly(A) tail at the 3' end of SEQ ID NO:169. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:169 from nucleotide 303 to nucleotide 542, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:169 from nucleotide 303 to nucleotide

20 542, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:169 from nucleotide 303 to nucleotide 542. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:169 from nucleotide 1 to nucleotide 435, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:169 from nucleotide 1 to

25 nucleotide 435, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:169 from nucleotide 1 to nucleotide 435.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:170;
- 30 (b) the amino acid sequence of SEQ ID NO:170 from amino acid 1 to amino acid 44;
- (c) a fragment of the amino acid sequence of SEQ ID NO:170, the fragment comprising eight contiguous amino acids of SEQ ID NO:170; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CO310_2
- 35 deposited with the ATCC under accession number 98279;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:170 or the amino acid sequence of SEQ ID NO:170 from amino acid 1 to amino acid 44. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:170 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:170, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:170 having biological activity, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:170.

10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:171;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:171 from nucleotide 40 to nucleotide 455;
- 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:171 from nucleotide 85 to nucleotide 455;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:171 from nucleotide 265 to nucleotide 515;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CP258_3 deposited with the ATCC under accession number 98279;
- 20 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CP258_3 deposited with the ATCC under accession number 98279;
- 25 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:172;
- 30 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:172 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:172;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
- 35 above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

5 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:171.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:171 from nucleotide 40 to nucleotide 455; the nucleotide sequence of SEQ ID NO:171 from nucleotide
10 85 to nucleotide 455; the nucleotide sequence of SEQ ID NO:171 from nucleotide 265 to nucleotide 515; the nucleotide sequence of the full-length protein coding sequence of clone CP258_3 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CP258_3 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a
15 mature protein encoded by the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:172 from amino acid 64 to amino acid 138. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence
20 of SEQ ID NO:172 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:172, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:172 having biological activity, the fragment comprising the amino acid sequence from amino acid 64 to amino acid 73 of SEQ ID NO:172.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:171.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- 30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (aa) SEQ ID NO:171, but excluding the poly(A) tail at the 3' end of SEQ ID NO:171; and

- (ab) the nucleotide sequence of the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (ba) SEQ ID NO:171, but excluding the poly(A) tail at the 3' end of SEQ ID NO:171; and
- (bb) the nucleotide sequence of the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide
- 20 sequence corresponding to the cDNA sequence of SEQ ID NO:171, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:171 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:171, but excluding the poly(A) tail at the 3' end of SEQ ID NO:171. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:171
- 25 from nucleotide 40 to nucleotide 455, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:171 from nucleotide 40 to nucleotide 455, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:171 from nucleotide 40 to nucleotide 455. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ
- 30 ID NO:171 from nucleotide 85 to nucleotide 455, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:171 from nucleotide 85 to nucleotide 455, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:171 from nucleotide 85 to nucleotide 455. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA
- 35 sequence of SEQ ID NO:171 from nucleotide 265 to nucleotide 515, and extending contiguously

from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:171 from nucleotide 265 to nucleotide 515, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:171 from nucleotide 265 to nucleotide 515.

In other embodiments, the present invention provides a composition comprising a protein,
5 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:172;
- (b) the amino acid sequence of SEQ ID NO:172 from amino acid 64 to amino acid 138;
- (c) a fragment of the amino acid sequence of SEQ ID NO:172, the fragment
10 comprising eight contiguous amino acids of SEQ ID NO:172; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CP258_3 deposited with the ATCC under accession number 98279;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:172 or the amino acid sequence of SEQ ID
15 NO:172 from amino acid 64 to amino acid 138. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:172 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:172, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:172 having biological activity,
20 the fragment comprising the amino acid sequence from amino acid 64 to amino acid 73 of SEQ ID NO:172.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:173;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:173 from nucleotide 105 to nucleotide 1007;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:173 from nucleotide 801 to nucleotide 1007;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:173
30 from nucleotide 1 to nucleotide 352;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CW1155_3 deposited with the ATCC under accession number 98279;
- 35 (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CW1155_3 deposited with the ATCC under accession number 98279;

- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CW1155_3 deposited with the ATCC under accession number 98279;
- 5 (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CW1155_3 deposited with the ATCC under accession number 98279;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:174;
- 10 (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:174 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:174;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 15 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:173.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:173 from nucleotide 105 to nucleotide 1007; the nucleotide sequence of SEQ ID NO:173 from nucleotide 801 to nucleotide 1007; the nucleotide sequence of SEQ ID NO:173 from nucleotide 1 to nucleotide 352; the nucleotide sequence of the full-length protein coding sequence of clone CW1155_3 deposited with the ATCC under accession number 98279; or the nucleotide sequence
- 25 of a mature protein coding sequence of clone CW1155_3 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CW1155_3 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID
- 30 NO:174 from amino acid 1 to amino acid 83. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:174 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:174, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:174 having biological activity, the fragment comprising the amino acid sequence from amino acid 145 to amino acid 154 of SEQ ID NO:174.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:173.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X
10 SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:173, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:173; and

(ab) the nucleotide sequence of the cDNA insert of clone
CW1155_3 deposited with the ATCC under accession number 98279;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

20 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:173, but excluding the poly(A) tail at the 3'
end of SEQ ID NO:173; and

25 (bb) the nucleotide sequence of the cDNA insert of clone CW1155_3 deposited with the ATCC under accession number 98279;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:173, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:173 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:173, but excluding the poly(A) tail at the 3'
35 end of SEQ ID NO:173. Also preferably the polynucleotide isolated according to the above

process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:173 from nucleotide 105 to nucleotide 1007, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:173 from nucleotide 105 to nucleotide 1007, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:173 from nucleotide 105 to nucleotide 1007. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:173 from nucleotide 801 to nucleotide 1007, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:173 from nucleotide 801 to nucleotide 1007, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:173 from nucleotide 801 to nucleotide 1007. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:173 from nucleotide 1 to nucleotide 352, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:173 from nucleotide 1 to nucleotide 352, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:173 from nucleotide 1 to nucleotide 352.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:174;
 - (b) the amino acid sequence of SEQ ID NO:174 from amino acid 1 to amino acid 83;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:174, the fragment comprising eight contiguous amino acids of SEQ ID NO:174; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone CW1155_3 deposited with the ATCC under accession number 98279;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:174 or the amino acid sequence of SEQ ID NO:174 from amino acid 1 to amino acid 83. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:174 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:174, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:174 having biological activity, the fragment comprising the amino acid sequence from amino acid 145 to amino acid 154 of SEQ ID NO:174.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:175;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:175 from nucleotide 11 to nucleotide 1699;
 - (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:175 from nucleotide 1682 to nucleotide 1699;
 - (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:175 from nucleotide 737 to nucleotide 1134;
 - (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CZ247_2 deposited with the ATCC under accession number 98279;
 - (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279;
 - (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CZ247_2 deposited with the ATCC under accession number 98279;
 - (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279;
 - (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:176;
 - (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:176 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:176;
 - (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
 - (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
 - (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
 - (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:175.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:175 from nucleotide 11 to nucleotide 1699; the nucleotide sequence of SEQ ID NO:175 from nucleotide 1682 to nucleotide 1699; the nucleotide sequence of SEQ ID NO:175 from nucleotide 737 to nucleotide 1134; the nucleotide sequence of the full-length protein coding sequence of

clone CZ247_2 deposited with the ATCC under accession number 98279; or the nucleotide sequence of a mature protein coding sequence of clone CZ247_2 deposited with the ATCC under accession number 98279. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:176 from amino acid 298 to amino acid 374. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:176 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:176, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:176 having biological activity, the fragment comprising the amino acid sequence from amino acid 276 to amino acid 285 of SEQ ID NO:176.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:175.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:175, but excluding the poly(A) tail at the 3' end of SEQ ID NO:175; and
 - (ab) the nucleotide sequence of the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (ba) SEQ ID NO:175, but excluding the poly(A) tail at the 3' end of SEQ ID NO:175; and

- (bb) the nucleotide sequence of the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:175, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:175 to a nucleotide
10 sequence corresponding to the 3' end of SEQ ID NO:175, but excluding the poly(A) tail at the 3' end of SEQ ID NO:175. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:175 from nucleotide 11 to nucleotide 1699, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:175 from nucleotide 11 to nucleotide
15 1699, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:175 from nucleotide 11 to nucleotide 1699. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:175 from nucleotide 1682 to nucleotide 1699, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:175 from
20 nucleotide 1682 to nucleotide 1699, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:175 from nucleotide 1682 to nucleotide 1699. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:175 from nucleotide 737 to nucleotide 1134, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said
25 sequence of SEQ ID NO:175 from nucleotide 737 to nucleotide 1134, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:175 from nucleotide 737 to nucleotide 1134.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:176;
- (b) the amino acid sequence of SEQ ID NO:176 from amino acid 298 to amino acid 374;
- (c) a fragment of the amino acid sequence of SEQ ID NO:176, the fragment comprising eight contiguous amino acids of SEQ ID NO:176; and

(d) the amino acid sequence encoded by the cDNA insert of clone CZ247_2 deposited with the ATCC under accession number 98279; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:176 or the amino acid sequence of SEQ ID NO:176 from amino acid 298 to amino acid 374. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:176 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:176, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:176 having biological activity, the fragment comprising the amino acid sequence from amino acid 276 to amino acid 285 of SEQ ID NO:176.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:177;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:177 from nucleotide 918 to nucleotide 1262;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:177 from nucleotide 999 to nucleotide 1262;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:177 from nucleotide 928 to nucleotide 1134;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM666_1 deposited with the ATCC under accession number 98292;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM666_1 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM666_1 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM666_1 deposited with the ATCC under accession number 98292;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:178;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:178 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:178;

- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- 5 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:177.
- 10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:177 from nucleotide 918 to nucleotide 1262; the nucleotide sequence of SEQ ID NO:177 from nucleotide 999 to nucleotide 1262; the nucleotide sequence of SEQ ID NO:177 from nucleotide 928 to nucleotide 1134; the nucleotide sequence of the full-length protein coding sequence of clone AM666_1 deposited with the ATCC under accession number 98292; or the nucleotide
- 15 sequence of a mature protein coding sequence of clone AM666_1 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM666_1 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID
- 20 NO:178 from amino acid 5 to amino acid 72. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:178 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:178, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
- 25 ID NO:178 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:178.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:177.

Further embodiments of the invention provide isolated polynucleotides produced

30 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

-231-

- (aa) SEQ ID NO:177, but excluding the poly(A) tail at the 3' end of SEQ ID NO:177; and
- (ab) the nucleotide sequence of the cDNA insert of clone AM666_1 deposited with the ATCC under accession number 98292;
- 5 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- 10 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:177, but excluding the poly(A) tail at the 3' end of SEQ ID NO:177; and
- 15 (bb) the nucleotide sequence of the cDNA insert of clone AM666_1 deposited with the ATCC under accession number 98292;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 20 (iv) isolating the polynucleotide products of step (b)(iii).
- Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:177, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:177 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:177, but excluding the poly(A) tail at the 3' end of SEQ ID NO:177. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:177 from nucleotide 918 to nucleotide 1262, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:177 from nucleotide 918 to nucleotide 1262, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:177 from nucleotide 918 to nucleotide 1262. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:177 from nucleotide 999 to nucleotide 1262, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:177 from nucleotide 999 to nucleotide 1262, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:177 from nucleotide 999 to nucleotide 1262. Also preferably the
- 35

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:177 from nucleotide 928 to nucleotide 1134, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:177 from nucleotide 928 to nucleotide 1134, to a nucleotide sequence
5 corresponding to the 3' end of said sequence of SEQ ID NO:177 from nucleotide 928 to nucleotide 1134.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:178;
- 10 (b) the amino acid sequence of SEQ ID NO:178 from amino acid 5 to amino acid 72;
- (c) a fragment of the amino acid sequence of SEQ ID NO:178, the fragment comprising eight contiguous amino acids of SEQ ID NO:178; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AM666_1
15 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:178 or the amino acid sequence of SEQ ID NO:178 from amino acid 5 to amino acid 72. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID
20 NO:178 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:178, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:178 having biological activity, the fragment comprising the amino acid sequence from amino acid 52 to amino acid 61 of SEQ ID NO:178.

25 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:179;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:179
from nucleotide 751 to nucleotide 906;
- 30 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:179
from nucleotide 829 to nucleotide 906;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:179
from nucleotide 556 to nucleotide 831;

- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BN387_3 deposited with the ATCC under accession number 98292;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BN387_3 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:180;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:180 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:180;
- (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:179.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:179 from nucleotide 751 to nucleotide 906; the nucleotide sequence of SEQ ID NO:179 from nucleotide 829 to nucleotide 906; the nucleotide sequence of SEQ ID NO:179 from nucleotide 556 to nucleotide 831; the nucleotide sequence of the full-length protein coding sequence of clone BN387_3 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone BN387_3 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:180 from amino acid 1 to amino acid 27. In further preferred embodiments, the present invention provides

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:180 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:180, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:180 having
5 biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:180.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:179.

Further embodiments of the invention provide isolated polynucleotides produced
10 according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:179, but excluding the poly(A) tail at the 3' end of SEQ ID NO:179; and

(ab) the nucleotide sequence of the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292;
(ii) hybridizing said probe(s) to human genomic DNA in conditions
20 at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);
and

- (b) a process comprising the steps of:
 - (i) preparing one or more polynucleotide primers that hybridize in
25 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:179, but excluding the poly(A) tail at the 3' end of SEQ ID NO:179; and

(bb) the nucleotide sequence of the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292;
30

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:179, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:179 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:179, but excluding the poly(A) tail at the 3' end of SEQ ID NO:179. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:179 from nucleotide 751 to nucleotide 906, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:179 from nucleotide 751 to nucleotide 906, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:179 from nucleotide 751 to nucleotide 906. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:179 from nucleotide 829 to nucleotide 906, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:179 from nucleotide 829 to nucleotide 906, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:179 from nucleotide 829 to nucleotide 906. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:179 from nucleotide 556 to nucleotide 831, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:179 from nucleotide 556 to nucleotide 831, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:179 from nucleotide 556 to nucleotide 831.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:180;
 - (b) the amino acid sequence of SEQ ID NO:180 from amino acid 1 to amino acid 27;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:180, the fragment comprising eight contiguous amino acids of SEQ ID NO:180; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BN387_3 deposited with the ATCC under accession number 98292;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:180 or the amino acid sequence of SEQ ID NO:180 from amino acid 1 to amino acid 27. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:180 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:180, or a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:180 having biological activity, the fragment comprising the amino acid sequence from amino acid 21 to amino acid 30 of SEQ ID NO:180.

In one embodiment, the present invention provides a composition comprising an isolated
5 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:181;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:181
from nucleotide 139 to nucleotide 765;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:181
10 from nucleotide 1 to nucleotide 416;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone BQ135_2 deposited with the ATCC under accession
number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
15 insert of clone BQ135_2 deposited with the ATCC under accession number 98292;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone BQ135_2 deposited with the ATCC under accession number
98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
20 of clone BQ135_2 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:182;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:182 having biological activity, the fragment comprising
25 eight contiguous amino acids of SEQ ID NO:182;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g)
above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h)
or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the
length of SEQ ID NO:181.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:181 from nucleotide 139 to nucleotide 765; the nucleotide sequence of SEQ ID NO:181 from nucleotide 1 to nucleotide 416; the nucleotide sequence of the full-length protein coding sequence of clone BQ135_2 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone BQ135_2 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BQ135_2 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:182 from amino acid 1 to amino acid 93. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:182 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:182, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:182 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:182.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:181.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - (aa) SEQ ID NO:181, but excluding the poly(A) tail at the 3' end of SEQ ID NO:181; and
 - (ab) the nucleotide sequence of the cDNA insert of clone BQ135_2 deposited with the ATCC under accession number 98292;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- 5 (ba) SEQ ID NO:181, but excluding the poly(A) tail at the 3' end of SEQ ID NO:181; and
- (bb) the nucleotide sequence of the cDNA insert of clone BQ135_2 deposited with the ATCC under accession number 98292;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 10 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:181, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:181 to a nucleotide

15 sequence corresponding to the 3' end of SEQ ID NO:181, but excluding the poly(A) tail at the 3' end of SEQ ID NO:181. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:181 from nucleotide 139 to nucleotide 765, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:181 from nucleotide 139 to nucleotide

20 765, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:181 from nucleotide 139 to nucleotide 765. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:181 from nucleotide 1 to nucleotide 416, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:181 from nucleotide 1 to

25 nucleotide 416, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:181 from nucleotide 1 to nucleotide 416.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:182;
- 30 (b) the amino acid sequence of SEQ ID NO:182 from amino acid 1 to amino acid 93;
- (c) a fragment of the amino acid sequence of SEQ ID NO:182, the fragment comprising eight contiguous amino acids of SEQ ID NO:182; and
- (d) the amino acid sequence encoded by the cDNA insert of clone BQ135_2
- 35 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:182 or the amino acid sequence of SEQ ID NO:182 from amino acid 1 to amino acid 93. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:182 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:182, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:182 having biological activity, the fragment comprising the amino acid sequence from amino acid 99 to amino acid 108 of SEQ ID NO:182.

- 10 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:183;
 - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:183 from nucleotide 214 to nucleotide 714;
 - 15 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:183 from nucleotide 151 to nucleotide 531;
 - (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CR678_1 deposited with the ATCC under accession number 98292;
 - 20 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CR678_1 deposited with the ATCC under accession number 98292;
 - (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CR678_1 deposited with the ATCC under accession number 98292;
 - 25 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CR678_1 deposited with the ATCC under accession number 98292;
 - (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:184;
 - (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:184 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:184;
 - 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
 - (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
 - 35

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:183.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:183 from nucleotide 214 to nucleotide 714; the nucleotide sequence of SEQ ID NO:183 from nucleotide 151 to nucleotide 531; the nucleotide sequence of the full-length protein coding sequence of clone CR678_1 deposited with the ATCC under accession number 98292; or the
10 nucleotide sequence of a mature protein coding sequence of clone CR678_1 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CR678_1 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid
15 sequence of SEQ ID NO:184 from amino acid 1 to amino acid 106. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:184 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:184, or a polynucleotide encoding a protein comprising a fragment of the
20 amino acid sequence of SEQ ID NO:184 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:184.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:183.

Further embodiments of the invention provide isolated polynucleotides produced
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:183, but excluding the poly(A) tail at the 3' end of SEQ ID NO:183; and

(ab) the nucleotide sequence of the cDNA insert of clone CR678_1 deposited with the ATCC under accession number 98292;

(ii) hybridizing said probe(s) to human genomic DNA in conditions
35 at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
- 5 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:183, but excluding the poly(A) tail at the 3' end of SEQ ID NO:183; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 10 CR678_1 deposited with the ATCC under accession number 98292;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 15 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:183, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:183 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:183, but excluding the poly(A) tail at the 3' end of SEQ ID NO:183. Also preferably the polynucleotide isolated according to the above
- 20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:183 from nucleotide 214 to nucleotide 714, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:183 from nucleotide 214 to nucleotide 714, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:183 from nucleotide 214 to nucleotide 714. Also preferably the polynucleotide isolated according to
- 25 the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:183 from nucleotide 151 to nucleotide 531, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:183 from nucleotide 151 to nucleotide 531, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:183 from nucleotide 151 to nucleotide 531.
- 30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:184;
- (b) the amino acid sequence of SEQ ID NO:184 from amino acid 1 to amino acid 106;

- (c) a fragment of the amino acid sequence of SEQ ID NO:184, the fragment comprising eight contiguous amino acids of SEQ ID NO:184; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CR678_1 deposited with the ATCC under accession number 98292;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:184 or the amino acid sequence of SEQ ID NO:184 from amino acid 1 to amino acid 106. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:184 having biological activity, the fragment preferably comprising eight (more preferably
- 10 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:184, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:184 having biological activity, the fragment comprising the amino acid sequence from amino acid 78 to amino acid 87 of SEQ ID NO:184.
- In one embodiment, the present invention provides a composition comprising an isolated
- 15 polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:185;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:185 from nucleotide 116 to nucleotide 4498;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:185
- 20 from nucleotide 1221 to nucleotide 1711;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CW420_2 deposited with the ATCC under accession number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA
- 25 insert of clone CW420_2 deposited with the ATCC under accession number 98292;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CW420_2 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
- 30 of clone CW420_2 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:186;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:186 having biological activity, the fragment comprising
- 35 eight contiguous amino acids of SEQ ID NO:186;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:185.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:185 from nucleotide 116 to nucleotide 4498; the nucleotide sequence of SEQ ID NO:185 from nucleotide 1221 to nucleotide 1711; the nucleotide sequence of the full-length protein coding sequence of clone CW420_2 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone CW420_2 deposited with the
15 ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CW420_2 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:186 from amino acid 370 to amino acid 532. In further preferred
20 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:186 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:186, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:186 having biological activity, the fragment comprising the
25 amino acid sequence from amino acid 725 to amino acid 734 of SEQ ID NO:186.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:185.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

30 (a) a process comprising the steps of:
(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

35 (aa) SEQ ID NO:185, but excluding the poly(A) tail at the 3' end of SEQ ID NO:185; and

- (ab) the nucleotide sequence of the cDNA insert of clone CW420_2 deposited with the ATCC under accession number 98292;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group
- 10 consisting of:
- (ba) SEQ ID NO:185, but excluding the poly(A) tail at the 3' end of SEQ ID NO:185; and
- (bb) the nucleotide sequence of the cDNA insert of clone CW420_2 deposited with the ATCC under accession number 98292;
- 15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide

20 sequence corresponding to the cDNA sequence of SEQ ID NO:185, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:185 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:185, but excluding the poly(A) tail at the 3' end of SEQ ID NO:185. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:185

25 from nucleotide 116 to nucleotide 4498, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:185 from nucleotide 116 to nucleotide 4498, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:185 from nucleotide 116 to nucleotide 4498. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of

30 SEQ ID NO:185 from nucleotide 1221 to nucleotide 1711, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:185 from nucleotide 1221 to nucleotide 1711, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:185 from nucleotide 1221 to nucleotide 1711.

In other embodiments, the present invention provides a composition comprising a protein,

35 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:186;
 - (b) the amino acid sequence of SEQ ID NO:186 from amino acid 370 to amino acid 532;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:186, the fragment comprising eight contiguous amino acids of SEQ ID NO:186; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone CW420_2 deposited with the ATCC under accession number 98292;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:186 or the amino acid sequence of SEQ ID NO:186 from amino acid 370 to amino acid 532. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:186 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:186, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:186 having biological activity, the fragment comprising the amino acid sequence from amino acid 725 to amino acid 734 of SEQ ID NO:186.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:187;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:187 from nucleotide 119 to nucleotide 2176;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:187 from nucleotide 1 to nucleotide 529;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone CW795_2 deposited with the ATCC under accession number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CW795_2 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:188;

- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:188 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:188;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:187.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:187 from nucleotide 119 to nucleotide 2176; the nucleotide sequence of SEQ ID NO:187 from nucleotide 1 to nucleotide 529; the nucleotide sequence of the full-length protein coding sequence of clone CW795_2 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone CW795_2 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:188 from amino acid 1 to amino acid 137. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:188 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:188, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:188 having biological activity, the fragment comprising the amino acid sequence from amino acid 338 to amino acid 347 of SEQ ID NO:188.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:187.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (aa) SEQ ID NO:187, but excluding the poly(A) tail at the 3' end of SEQ ID NO:187; and

(ab) the nucleotide sequence of the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

10 (iii) isolating the DNA polynucleotides detected with the probe(s);
and

(b) a process comprising the steps of:

15 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:187, but excluding the poly(A) tail at the 3' end of SEQ ID NO:187; and

(bb) the nucleotide sequence of the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292;

20 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

25 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:187, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:187 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:187, but excluding the poly(A) tail at the 3' end of SEQ ID NO:187. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:187
30 from nucleotide 119 to nucleotide 2176, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:187 from nucleotide 119 to nucleotide 2176, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:187 from nucleotide 119 to nucleotide 2176. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of
35 SEQ ID NO:187 from nucleotide 1 to nucleotide 529, and extending contiguously from a

nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:187 from nucleotide 1 to nucleotide 529, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:187 from nucleotide 1 to nucleotide 529.

In other embodiments, the present invention provides a composition comprising a protein,
5 wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:188;
- (b) the amino acid sequence of SEQ ID NO:188 from amino acid 1 to amino acid 137;
- (c) a fragment of the amino acid sequence of SEQ ID NO:188, the fragment
10 comprising eight contiguous amino acids of SEQ ID NO:188; and
- (d) the amino acid sequence encoded by the cDNA insert of clone CW795_2 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:188 or the amino acid sequence of SEQ ID
15 NO:188 from amino acid 1 to amino acid 137. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:188 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:188, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:188 having biological activity,
20 the fragment comprising the amino acid sequence from amino acid 338 to amino acid 347 of SEQ ID NO:188.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:189;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:189 from nucleotide 401 to nucleotide 589;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:189 from nucleotide 258 to nucleotide 627;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
30 protein coding sequence of clone CW823_3 deposited with the ATCC under accession number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone CW823_3 deposited with the ATCC under accession number 98292;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone CW823_3 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert
5 of clone CW823_3 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:190;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:190 having biological activity, the fragment comprising
10 eight contiguous amino acids of SEQ ID NO:190;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of
15 the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:189.
- 20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:189 from nucleotide 401 to nucleotide 589; the nucleotide sequence of SEQ ID NO:189 from nucleotide 258 to nucleotide 627; the nucleotide sequence of the full-length protein coding sequence of clone CW823_3 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone CW823_3 deposited with the
25 ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone CW823_3 deposited with the ATCC under accession number 98292. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:190 having biological activity, the fragment preferably
30 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:190, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:190 having biological activity, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:190.
- Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID
35 NO:189.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:189, but excluding the poly(A) tail at the 3' end of SEQ ID NO:189; and

(ab) the nucleotide sequence of the cDNA insert of clone CW823_3 deposited with the ATCC under accession number 98292;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:189, but excluding the poly(A) tail at the 3' end of SEQ ID NO:189; and

(bb) the nucleotide sequence of the cDNA insert of clone CW823_3 deposited with the ATCC under accession number 98292;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:189, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:189 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:189, but excluding the poly(A) tail at the 3' end of SEQ ID NO:189. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:189 from nucleotide 401 to nucleotide 589, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:189 from nucleotide 401 to nucleotide 589, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:189

from nucleotide 401 to nucleotide 589. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:189 from nucleotide 258 to nucleotide 627, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:189 from nucleotide 258 to nucleotide 627, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:189 from nucleotide 258 to nucleotide 627.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:190;
- 10 (b) a fragment of the amino acid sequence of SEQ ID NO:190, the fragment comprising eight contiguous amino acids of SEQ ID NO:190; and
- (c) the amino acid sequence encoded by the cDNA insert of clone CW823_3 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein
15 comprises the amino acid sequence of SEQ ID NO:190. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:190 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:190, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:190 having biological activity,
20 the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:190.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:191;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:191 from nucleotide 548 to nucleotide 868;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:191 from nucleotide 590 to nucleotide 868;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length
30 protein coding sequence of clone DF989_3 deposited with the ATCC under accession number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone DF989_3 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:192;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:192 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:192;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:191.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:191 from nucleotide 548 to nucleotide 868; the nucleotide sequence of SEQ ID NO:191 from nucleotide 590 to nucleotide 868; the nucleotide sequence of the full-length protein coding sequence of clone DF989_3 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone DF989_3 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:192 from amino acid 75 to amino acid 107. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:192 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:192, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:192 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:192.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:191 and SEQ ID NO:193.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (aa) SEQ ID NO:191;
- (ab) SEQ ID NO:193, but excluding the poly(A) tail at the 3' end of SEQ ID NO:193; and
- (ac) the nucleotide sequence of the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292;
- 15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that hybridize in
- 20 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:191;
- (bb) SEQ ID NO:193, but excluding the poly(A) tail at the 3' end of SEQ ID NO:193; and
- 25 (bc) the nucleotide sequence of the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- 30 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequences of SEQ ID NO:191 and SEQ ID NO:193, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:191 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:193, but excluding

35 the poly(A) tail at the 3' end of SEQ ID NO:193. Also preferably the polynucleotide isolated

according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:191, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:191 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:191. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:191 from nucleotide 548 to nucleotide 868, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:191 from nucleotide 548 to nucleotide 868, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:191 from nucleotide 548 to nucleotide 868. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:191 from nucleotide 590 to nucleotide 868, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:191 from nucleotide 590 to nucleotide 868, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:191 from nucleotide 590 to nucleotide 868.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:192;
- (b) the amino acid sequence of SEQ ID NO:192 from amino acid 75 to amino acid 107;
- (c) a fragment of the amino acid sequence of SEQ ID NO:192, the fragment comprising eight contiguous amino acids of SEQ ID NO:192; and
- (d) the amino acid sequence encoded by the cDNA insert of clone DF989_3 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:192 or the amino acid sequence of SEQ ID NO:192 from amino acid 75 to amino acid 107. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:192 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:192, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:192 having biological activity, the fragment comprising the amino acid sequence from amino acid 48 to amino acid 57 of SEQ ID NO:192.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:194;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:194 from nucleotide 251 to nucleotide 787;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:194 from nucleotide 371 to nucleotide 787;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone DL162_1 deposited with the ATCC under accession number 98292;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone DL162_1 deposited with the ATCC under accession number 98292;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone DL162_1 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone DL162_1 deposited with the ATCC under accession number 98292;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:195;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:195 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:195;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:194.
- Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:194
- 30 from nucleotide 251 to nucleotide 787; the nucleotide sequence of SEQ ID NO:194 from nucleotide 371 to nucleotide 787; the nucleotide sequence of the full-length protein coding sequence of clone DL162_1 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone DL162_1 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide
- 35 encodes the full-length or a mature protein encoded by the cDNA insert of clone DL162_1

deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:195 from amino acid 38 to amino acid 170. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a
5 fragment of the amino acid sequence of SEQ ID NO:195 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:195, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:195 having biological activity, the fragment comprising the amino acid sequence from amino acid 84 to amino acid 93 of SEQ ID NO:195.

10 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:194.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:
15 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:194, but excluding the poly(A) tail at the 3' end of SEQ ID NO:194; and

20 (ab) the nucleotide sequence of the cDNA insert of clone DL162_1 deposited with the ATCC under accession number 98292;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

25 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:194, but excluding the poly(A) tail at the 3' end of SEQ ID NO:194; and

(bb) the nucleotide sequence of the cDNA insert of clone DL162_1 deposited with the ATCC under accession number 98292;

35 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:194, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:194 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:194, but excluding the poly(A) tail at the 3' end of SEQ ID NO:194. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:194 from nucleotide 251 to nucleotide 787, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:194 from nucleotide 251 to nucleotide 787, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:194 from nucleotide 251 to nucleotide 787. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:194 from nucleotide 371 to nucleotide 787, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:194 from nucleotide 371 to nucleotide 787, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:194 from nucleotide 371 to nucleotide 787.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:195;
- (b) the amino acid sequence of SEQ ID NO:195 from amino acid 38 to amino acid 170;
- (c) a fragment of the amino acid sequence of SEQ ID NO:195, the fragment comprising eight contiguous amino acids of SEQ ID NO:195; and
- (d) the amino acid sequence encoded by the cDNA insert of clone DL162_1 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:195 or the amino acid sequence of SEQ ID NO:195 from amino acid 38 to amino acid 170. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:195 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:195, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:195 having biological activity, the fragment comprising the amino acid sequence from amino acid 84 to amino acid 93 of SEQ ID NO:195.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:196;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:196
5 from nucleotide 121 to nucleotide 3345;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:196
from nucleotide 160 to nucleotide 3345;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:196
from nucleotide 2592 to nucleotide 3318;
- 10 (e) a polynucleotide comprising the nucleotide sequence of the full-length
protein coding sequence of clone DL162_2 deposited with the ATCC under accession
number 98292;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA
insert of clone DL162_2 deposited with the ATCC under accession number 98292;
- 15 (g) a polynucleotide comprising the nucleotide sequence of a mature protein
coding sequence of clone DL162_2 deposited with the ATCC under accession number
98292;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert
of clone DL162_2 deposited with the ATCC under accession number 98292;
- 20 (i) a polynucleotide encoding a protein comprising the amino acid sequence
of SEQ ID NO:197;
- (j) a polynucleotide encoding a protein comprising a fragment of the amino
acid sequence of SEQ ID NO:197 having biological activity, the fragment comprising
eight contiguous amino acids of SEQ ID NO:197;
- 25 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h)
above;
- (l) a polynucleotide which encodes a species homologue of the protein of (i)
or (j) above ;
- (m) a polynucleotide that hybridizes under stringent conditions to any one of
30 the polynucleotides specified in (a)-(j); and
- (n) a polynucleotide that hybridizes under stringent conditions to any one of
the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the
length of SEQ ID NO:196.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:196
35 from nucleotide 121 to nucleotide 3345; the nucleotide sequence of SEQ ID NO:196 from

nucleotide 160 to nucleotide 3345; the nucleotide sequence of SEQ ID NO:196 from nucleotide 2592 to nucleotide 3318; the nucleotide sequence of the full-length protein coding sequence of clone DL162_2 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone DL162_2 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone DL162_2 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:197 from amino acid 860 to amino acid 1066. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:197 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:197, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:197 having biological activity, the fragment comprising the amino acid sequence from amino acid 532 to amino acid 541 of SEQ ID NO:197.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:196.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 20 (a) a process comprising the steps of:
 - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
 - 25 (aa) SEQ ID NO:196, but excluding the poly(A) tail at the 3' end of SEQ ID NO:196; and
 - (ab) the nucleotide sequence of the cDNA insert of clone DL162_2 deposited with the ATCC under accession number 98292;
 - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
 - 30 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
 - 35 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:196, but excluding the poly(A) tail at the 3' end of SEQ ID NO:196; and

(bb) the nucleotide sequence of the cDNA insert of clone DL162_2 deposited with the ATCC under accession number 98292;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide
10 sequence corresponding to the cDNA sequence of SEQ ID NO:196, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:196 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:196, but excluding the poly(A) tail at the 3' end of SEQ ID NO:196. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:196
15 from nucleotide 121 to nucleotide 3345, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:196 from nucleotide 121 to nucleotide 3345, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:196 from nucleotide 121 to nucleotide 3345. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of
20 SEQ ID NO:196 from nucleotide 160 to nucleotide 3345, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:196 from nucleotide 160 to nucleotide 3345, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:196 from nucleotide 160 to nucleotide 3345. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence
25 corresponding to the cDNA sequence of SEQ ID NO:196 from nucleotide 2592 to nucleotide 3318, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:196 from nucleotide 2592 to nucleotide 3318, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:196 from nucleotide 2592 to nucleotide 3318.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:197;

(b) the amino acid sequence of SEQ ID NO:197 from amino acid 860 to amino acid 1066;

- (c) a fragment of the amino acid sequence of SEQ ID NO:197, the fragment comprising eight contiguous amino acids of SEQ ID NO:197; and
- (d) the amino acid sequence encoded by the cDNA insert of clone DL162_2 deposited with the ATCC under accession number 98292;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:197 or the amino acid sequence of SEQ ID NO:197 from amino acid 860 to amino acid 1066. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:197 having biological activity, the fragment preferably comprising eight (more preferably
- 10 twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:197, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:197 having biological activity, the fragment comprising the amino acid sequence from amino acid 532 to amino acid 541 of SEQ ID NO:197.

In one embodiment, the present invention provides a composition comprising an isolated

15 polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:198;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:198 from nucleotide 117 to nucleotide 2600;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:198
- 20 from nucleotide 2130 to nucleotide 2600;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:198 from nucleotide 1 to nucleotide 506;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone EC172_1 deposited with the ATCC under accession
- 25 number 98292;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292;
- (g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone EC172_1 deposited with the ATCC under accession number
- 30 98292;
- (h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292;
- (i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:199;

(j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:199 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:199;

5 (k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

10 (n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:198.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:198 from nucleotide 117 to nucleotide 2600; the nucleotide sequence of SEQ ID NO:198 from
15 nucleotide 2130 to nucleotide 2600; the nucleotide sequence of SEQ ID NO:198 from nucleotide 1 to nucleotide 506; the nucleotide sequence of the full-length protein coding sequence of clone EC172_1 deposited with the ATCC under accession number 98292; or the nucleotide sequence of a mature protein coding sequence of clone EC172_1 deposited with the ATCC under accession number 98292. In other preferred embodiments, the polynucleotide encodes the full-length or a
20 mature protein encoded by the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:199 from amino acid 1 to amino acid 130. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ
25 ID NO:199 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:199, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:199 having biological activity, the fragment comprising the amino acid sequence from amino acid 409 to amino acid 418 of SEQ ID NO:199.

30 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:198.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:198, but excluding the poly(A) tail at the 3' end of SEQ ID NO:198; and

(ab) the nucleotide sequence of the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:198, but excluding the poly(A) tail at the 3' end of SEQ ID NO:198; and

(bb) the nucleotide sequence of the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:198, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:198 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:198, but excluding the poly(A) tail at the 3' end of SEQ ID NO:198. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:198 from nucleotide 117 to nucleotide 2600, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:198 from nucleotide 117 to nucleotide 2600, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:198 from nucleotide 117 to nucleotide 2600. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:198 from nucleotide 2130 to nucleotide 2600, and extending contiguously from a

nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:198 from nucleotide 2130 to nucleotide 2600, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:198 from nucleotide 2130 to nucleotide 2600. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence
5 corresponding to the cDNA sequence of SEQ ID NO:198 from nucleotide 1 to nucleotide 506, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:198 from nucleotide 1 to nucleotide 506, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:198 from nucleotide 1 to nucleotide 506.

- In other embodiments, the present invention provides a composition comprising a protein,
10 wherein said protein comprises an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:199;
 - (b) the amino acid sequence of SEQ ID NO:199 from amino acid 1 to amino acid 130;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:199, the fragment
15 comprising eight contiguous amino acids of SEQ ID NO:199; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone EC172_1 deposited with the ATCC under accession number 98292;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:199 or the amino acid sequence of SEQ ID
20 NO:199 from amino acid 1 to amino acid 130. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:199 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:199, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:199 having biological activity,
25 the fragment comprising the amino acid sequence from amino acid 409 to amino acid 418 of SEQ ID NO:199.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with such polynucleotide compositions. Also provided by the
30 present invention are organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein.

Processes are also provided for producing a protein, which comprise:

- (a) growing a culture of the host cell transformed with such polynucleotide compositions in a suitable culture medium; and
35
- (b) purifying the protein from the culture.

The protein produced according to such methods is also provided by the present invention.

Protein compositions of the present invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

- 5 Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

10 BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are schematic representations of the pED6 and pNOTs vectors, respectively, used for deposit of clones disclosed herein.

DETAILED DESCRIPTION

15 ISOLATED PROTEINS AND POLYNUCLEOTIDES

Nucleotide and amino acid sequences, as presently determined, are reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone in accordance with known methods. The predicted amino acid sequence (both full-length and mature forms) can then be
20 determined from such nucleotide sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have determined to be the reading frame best identifiable with sequence information available at the time of filing.

- 25 As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported
30 across the membrane of the endoplasmic reticulum.

Clone "AX65_22"

A polynucleotide of the present invention has been identified as clone "AX65_22". AX65_22 was isolated from a human adult testes cDNA library using methods which are selective
35 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding

a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AX65_22 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AX65_22 protein").

The nucleotide sequence of the 5' portion of AX65_22 as presently determined is reported in SEQ ID NO:1. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:2. The predicted amino acid sequence of the AX65_22 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2. Amino acids 8 to 20 of SEQ ID NO:2 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AX65_22 protein. Additional nucleotide sequence from the 3' portion of AX65_22, including a poly(A) tail, is reported in SEQ ID NO:3.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AX65_22 should be approximately 3500 bp.

The nucleotide sequence disclosed herein for AX65_22 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AX65_22 demonstrated at least some similarity with sequences identified as T08476 (Eukaryotic expression vector pAPEX-3p) and U46493 (Cloning vector pFlp recombinase gene, complete cds). The predicted AX65_22 protein demonstrated at least some homology with sequences identified as J01969 (DNA polymerase [Human adenovirus type 5]), R07640 (Deduced protein sequence of p170-2 comprising T4), and X57205 (fibroblast growth factor receptor [Homo sapiens]). Based upon sequence similarity, AX65_22 proteins and each similar protein or peptide may share at least some activity.

Clone "BD335_14"

A polynucleotide of the present invention has been identified as clone "BD335_14". BD335_14 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BD335_14 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BD335_14 protein").

The nucleotide sequence of BD335_14 as presently determined is reported in SEQ ID NO:4, and includes a poly(A) tail. What applicants presently believe to be the proper reading

frame and the predicted amino acid sequence of the BD335_14 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:5.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BD335_14 should be approximately 3000 bp.

- 5 The nucleotide sequence disclosed herein for BD335_14 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. The predicted BD335_14 protein demonstrated at least some similarity with sequences identified as U83511 (APXL [Homo sapiens]). Based upon sequence similarity, BD335_14 proteins and each homologous protein or peptide may share at least some activity. The
- 10 TopPredII computer program predicts three potential transmembrane domains within the BD335_14 protein sequence, one centered around amino acid 80, another around amino acid 320, and a third around amino acid 700 of SEQ ID NO:5.

Clone "BG241_1"

- 15 A polynucleotide of the present invention has been identified as clone "BG241_1". BG241_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG241_1 is a full-length clone, including the entire coding sequence of
- 20 a secreted protein (also referred to herein as "BG241_1 protein").

- The nucleotide sequence of the 5' portion of BG241_1 as presently determined is reported in SEQ ID NO:6. An additional internal nucleotide sequence from BG241_1 as presently determined is reported in SEQ ID NO:7. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:8.
- 25 Additional nucleotide sequence from the 3' portion of BG241_1, including a poly(A) tail, is reported in SEQ ID NO:9.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG241_1 should be approximately 800 bp.

- The nucleotide sequence disclosed herein for BG241_1 was searched against the GenBank
- 30 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BG241_1 demonstrated at least some similarity with sequences identified as AJ082187 (ox75f01.x1 Soares_NhHMPu_S1 Homo sapiens cDNA clone IMAGE 1662169 3' similar to contains element MSR1 repetitive element; mRNA sequence), W38781 (zb27g08.r1 Soares parathyroid tumor NbHPA Homo sapiens), and Y12781 (Homo sapiens
- 35 mRNA for transducin (beta) like 1 protein). The predicted BG241_1 protein demonstrated at

least some similarity to sequences identified as Y12781 (transducin (beta) like 1 protein [Homo sapiens]) and other beta-transducin-like proteins (see GenBank accession numbers L28125 and T86738). Based upon sequence similarity, BG241_1 proteins and each similar protein or peptide may share at least some activity.

5

Clone "BL187_4"

A polynucleotide of the present invention has been identified as clone "BL187_4". BL187_4 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
10 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BL187_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BL187_4 protein").

The nucleotide sequence of BL187_4 as presently determined is reported in SEQ ID NO:10, and includes a poly(A) tail. What applicants presently believe to be the proper reading
15 frame and the predicted amino acid sequence of the BL187_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:11. Amino acids 17 to 29 of SEQ ID NO:11 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal
20 sequence not be separated from the remainder of the BL187_4 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BL187_4 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for BL187_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
25 protocols. BL187_4 demonstrated at least some similarity with sequences identified as AA476210 (zw35g01.s1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 771312 3', mRNA sequence), AA868505 (ak43b04.s1 Soares testis NHT Homo sapiens cDNA clone IMAGE 1408687 3' similar to SW PLZF_HUMAN Q05516 ZINC FINGER PROTEIN PLZF; mRNA sequence), AA927876 (om18b09.s1 Soares NFL T GBC S1
30 Homo sapiens cDNA clone IMAGE:1541369 3', mRNA sequence), AD000671 (Homo sapiens DNA from chromosome 19-cosmid f24109 containing HRX2, genomic sequence), H48938 (EST0010 Homo sapiens cDNA clone HTN-6-15), and Z63958 (H.sapiens CpG DNA, clone 93d10, forward read cpg93d10.ft1a). The predicted amino acid sequence disclosed herein for BL187_4 was searched against the GenPept and GeneSeq amino acid

sequence databases using the BLASTX search protocol. The predicted BL187_4 protein demonstrated at least some similarity to sequences identified as R95242 (HIC-1 polypeptide), Z19002 (kruppel-like zinc finger protein [Homo sapiens]), and a number of other zinc-finger proteins. Based upon sequence similarity, BL187_4 proteins and each similar protein or peptide may share at least some activity. Motifs analysis indicates the presence of two zinc-finger (C2H2 type) domains centered around amino acids 375 and 430 of SEQ ID NO:11, respectively. The TopPredII computer program predicts two potential transmembrane domains within the BL187_4 protein sequence, one centered around amino acid 30 and another around amino acid 260 of SEQ ID NO:11. BL187_4 protein appears to be a novel secreted or membrane-associated zinc-finger protein.

Clone "BL249_18"

A polynucleotide of the present invention has been identified as clone "BL249_18". BL249_18 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BL249_18 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BL249_18 protein").

The nucleotide sequence of BL249_18 as presently determined is reported in SEQ ID NO:12, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BL249_18 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:13. Amino acids 32 to 44 of SEQ ID NO:13 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 45. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BL249_18 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BL249_18 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for BL249_18 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BL249_18 demonstrated at least some similarity with sequences identified as AA034864 (mi53f01.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 467257 5'), AA115100 (zl02h12.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 491207 3'), AA219365 (zr04c06.r1 Stratagene NT2 neuronal precursor 937230 Homo sapiens cDNA clone 650506 5'), AA399095 (zt59b06.r1 Soares

testis NHT Homo sapiens cDNA clone 726611 5'), R82633 (yj20a05.s1 Homo sapiens cDNA clone 149264 3'), and T22047 (Human gene signature HUMGS03590). The predicted amino acid sequence disclosed herein for BL249_18 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted

5 BL249_18 protein demonstrated at least some similarity to sequences identified as AC003673 (unknown protein (AAC09020.1) [Arabidopsis thaliana]) and Z98598 (hypothetical protein [Schizosaccharomyces pombe]). Based upon sequence similarity, BL249_18 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the BL249_18 protein sequence, one

10 centered around amino acid 45 and another around amino acid 680 of SEQ ID NO:13.

Clone "BO71_1"

A polynucleotide of the present invention has been identified as clone "BO71_1". BO71_1 was isolated from a human adult retina cDNA library using methods which are selective

15 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BO71_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BO71_1 protein").

The nucleotide sequence of the 5' portion of BO71_1 as presently determined is reported

20 in SEQ ID NO:14. An additional internal nucleotide sequence from BO71_1 as presently determined is reported in SEQ ID NO:15. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:16. Additional nucleotide sequence from the 3' portion of BO71_1, including a poly(A) tail, is reported in SEQ ID NO:17.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BO71_1 should be approximately 2000 bp.

The nucleotide sequence disclosed herein for BO71_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BO71_1 demonstrated at least some similarity with sequences identified as X86809

30 (H.sapiens mRNA for major astrocytic phosphoprotein PEA-15). The predicted amino acid sequence disclosed herein for BO71_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BO71_1 protein demonstrated at least some similarity to sequences identified as U06144 (cellular disintegrin-related protein [Mus musculus]). Based upon sequence similarity, BO71_1

35 proteins and each similar protein or peptide may share at least some activity.

Clone "BO365_2"

A polynucleotide of the present invention has been identified as clone "BO365_2". BO365_2 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
5 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BO365_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BO365_2 protein").

The nucleotide sequence of BO365_2 as presently determined is reported in SEQ ID NO:18, and includes a poly(A) tail. What applicants presently believe to be the proper reading
10 frame and the predicted amino acid sequence of the BO365_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:19.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BO365_2 should be approximately 2800 bp.

The nucleotide sequence disclosed herein for BO365_2 was searched against the GenBank
15 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BO365_2 demonstrated at least some similarity with sequences identified as D63876 (Human mRNA for KIAA0154 gene, partial cds) and Z83844 (Human DNA sequence *** SEQUENCING IN PROGRESS *** from clone 37E16; HTGS phase 1). The predicted amino acid sequence disclosed herein for BO365_2 was searched against the GenPept and
20 GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BO365_2 protein demonstrated at least some similarity to sequences identified as D10250 (alpha-fetoprotein enhancer binding protein [Homo sapiens]), D63876 (KIAA0154 gene product is related to mouse gamma adaptin [Homo sapiens]), and R23962 (AFP-1). Based upon sequence similarity, BO365_2 proteins and each similar protein or peptide may share at least
25 some activity. The TopPredII computer program predicts three potential transmembrane domains within the BO365_2 protein sequence, centered around amino acids 70, 140, and 180 of SEQ ID NO:19, respectively.

Clone "BV51_1"

A polynucleotide of the present invention has been identified as clone "BV51_1". BV51_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
30 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV51_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV51_1 protein").
35

The nucleotide sequence of the 5' portion of BV51_1 as presently determined is reported in SEQ ID NO:20. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:21. The predicted amino acid sequence of the BV51_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:21. Additional
5 nucleotide sequence from the 3' portion of BV51_1, including a poly(A) tail, is reported in SEQ ID NO:22.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV51_1 should be approximately 970 bp.

The nucleotide sequence disclosed herein for BV51_1 was searched against the GenBank
10 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV51_1 demonstrated at least some similarity with sequences identified as AB012130 (Homo sapiens SBC2 mRNA for sodium bicarbonate cotransporter2, complete cds) and U46493 (Cloning vector pFlp recombinase gene, complete cds). The predicted amino acid
15 sequence disclosed herein for BV51_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BV51_1 protein demonstrated at least some similarity to sequences identified as AB01213 (sodium bicarbonate cotransporter2 [Homo sapiens]). Based upon sequence similarity, BV51_1 proteins and each similar protein or peptide may share at least some activity.

20 Clone "BV140_3"

A polynucleotide of the present invention has been identified as clone "BV140_3". BV140_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence
25 of the encoded protein. BV140_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV140_3 protein").

The nucleotide sequence of the 5' portion of BV140_3 as presently determined is reported in SEQ ID NO:23. An additional internal nucleotide sequence from BV140_3 as presently determined is reported in SEQ ID NO:24. What applicants believe is the proper reading frame
30 and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:25. Additional nucleotide sequence from the 3' portion of BV140_3, including a poly(A) tail, is reported in SEQ ID NO:26.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV140_3 should be approximately 3500 bp.

The nucleotide sequence disclosed herein for BV140_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV140_3 demonstrated at least some similarity with sequences identified as H72799 (yu07d10.r1 Homo sapiens cDNA clone 233107 5') and T94057 (ye33g08.r1 Homo sapiens cDNA clone 119582 5'). Based upon sequence similarity, BV140_3 proteins and each similar protein or peptide may share at least some activity.

Clone "BV141_2"

A polynucleotide of the present invention has been identified as clone "BV141_2". BV141_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV141_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV141_2 protein").

The nucleotide sequence of BV141_2 as presently determined is reported in SEQ ID NO:27, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BV141_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV141_2 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for BV141_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV141_2 demonstrated at least some similarity with sequences identified as L26860 (Mus musculus (C6e) heavy chain immunoglobulin variable region gene). Based upon sequence similarity, BV141_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the BV141_2 protein sequence, one centered around amino acid 34 and another around amino acid 65 of SEQ ID NO:28. The nucleotide sequence of BV141_2 indicates that it may contain one or more of the following repetitive element(s): L1 repeat.

Clone "CC194_4"

A polynucleotide of the present invention has been identified as clone "CC194_4". CC194_4 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. CC194_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CC194_4 protein").

The nucleotide sequence of the 5' portion of CC194_4 as presently determined is reported in SEQ ID NO:29. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:30. The predicted amino acid sequence of the CC194_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:30. Amino acids 88 to 100 of SEQ ID NO:30 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 101. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CC194_4 protein. Additional nucleotide sequence from the 3' portion of CC194_4, including a poly(A) tail, is reported in SEQ ID NO:31.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CC194_4 should be approximately 3300 bp.

The nucleotide sequence disclosed herein for CC194_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CC194_4 demonstrated at least some similarity with sequences identified as AA722214 (zh20f10.s1 Soares pineal gland N3HPG Homo sapiens cDNA clone 412651 3', mRNA sequence), H11476 (ym10h08.s1 Homo sapiens cDNA clone 47781 3'), H11581 (ym10h08.r1 Soares infant brain 1NIB Homo sapiens cDNA clone IMAGE:47781 5' similar to SP:C36E8.3 CE00911; mRNA sequence), H23044 (ym51d07.r1 Homo sapiens cDNA clone 52058 5' similar to SP:C36E8.3 CE00911), N93789 (zb64g05.s1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 308408 3'), and W54544 (mc99a01.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus). The predicted amino acid sequence disclosed herein for CC194_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CC194_4 protein demonstrated at least some similarity to sequences identified as M38561 (CAD [Homo sapiens]). Based upon sequence similarity, CC194_4 proteins and each similar protein or peptide may share at least some activity.

30

Clone "DA136_11"

A polynucleotide of the present invention has been identified as clone "DA136_11". DA136_11 was isolated from a human adult placenta cDNA library using methods which are

35

selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. DA136_11 includes at least a portion of the coding sequence of a secreted protein (also referred to herein as "DA136_11 protein").

5 The nucleotide sequence of DA136_11 as presently determined is reported in SEQ ID NO:32, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the DA136_11 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:33.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone DA136_11 should be approximately 3800 bp.

 The nucleotide sequence disclosed herein for DA136_11 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. DA136_11 demonstrated at least some similarity with sequences identified as AA523414 (ng30a07.s1 NCI_CGAP_Co3 Homo sapiens cDNA clone 936276), H89334
15 (yw25h09.r1 Homo sapiens cDNA clone 253313 5'), R59925 (yh11b12.s1 Homo sapiens cDNA clone 42891 3), T66165 (Human interleukin-12 receptor alpha chain NR4 DNA), Y09328 (H.sapiens mRNA for IL13 receptor alpha-1 chain), and Y10659 (H.sapiens IL-13Ra mRNA). The predicted amino acid sequence disclosed herein for DA136_11 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX
20 search protocol. The predicted DA136_11 protein demonstrated at least some similarity to sequences identified as L08960 (cell adhesion molecule [Gallus gallus]), M34083 (lactogen receptor precursor [Rattus norvegicus]), M59941 (GM-CSF receptor beta chain [Homo sapiens]), W09822 (Human interleukin-12 receptor alpha chain NR4), X61178 (interleukin-5 receptor type 3 [Homo sapiens]), and Y10659 (IL-13Ra [Homo sapiens]).
25 Based upon sequence similarity, DA136_11 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the DA136_11 protein sequence, centered around amino acid 215 of SEQ ID NO:33.

30 Clone "AR415_4"

 A polynucleotide of the present invention has been identified as clone "AR415_4". AR415_4 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. AR415_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AR415_4 protein").

The nucleotide sequence of AR415_4 as presently determined is reported in SEQ ID NO:34, and includes a poly(A) tail. What applicants presently believe to be the proper reading
 5 frame and the predicted amino acid sequence of the AR415_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:35. Amino acids 14 to 26 of SEQ ID NO:35 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal
 10 sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AR415_4 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AR415_4 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for AR415_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
 15 protocols. AR415_4 demonstrated at least some similarity with sequences identified as AA100799 (zm26d01.s1 Stratagene pancreas (#937208) Homo sapiens cDNA clone 526753 3'), AA100852 (zm26d01.r1 Stratagene pancreas (#937208) Homo sapiens cDNA clone 526753 5' similar to SW CO02_HUMAN P19075 TUMOR-ASSOCIATED ANTIGEN CO-029), AA146605 (zo35c09.r1 Stratagene colon (#937204) Homo sapiens cDNA clone 588880 5' similar to
 20 SW:CO02_HUMAN P19075 TUMOR-ASSOCIATED ANTIGEN CO-029), AA224847 (nc33c12.s1 NCI CGAP Pr2 Homo sapiens cDNA clone 4079 similar to SW:CO02_HUMAN P19075 TUMOR-ASSOCIATED ANTIGEN CO-029), AA225191 (nc21h08.s1 NCI CGAP Pr1 Homo sapiens cDNA clone 2968), AA593864 (nn19f08.s1 NCI CGAP Co12 Homo sapiens cDNA clone IMAGE:1084359), D26483 (Mouse mRNA for PE31/TALLA), M33680 (Human
 25 26-kDa cell surface protein TAPA-1 mRNA, complete cds), T14726 (Human CD53 antigen cDNA), and T23814 (Human gene signature HUMGS05723). The predicted amino acid sequence disclosed herein for AR415_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AR415_4 protein demonstrated at least some sequence similarity with sequences identified as D29808 (TALLA-1
 30 [Homo sapiens]), M35252 (tumor-associated antigen [Homo sapiens]), and R22360 (CO-029 tumour associated antigen protein). Based upon sequence similarity, AR415_4 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the AR415_4 protein sequence centered around amino acid 100 of SEQ ID NO:35.

Clone "AS63_29"

A polynucleotide of the present invention has been identified as clone "AS63_29". AS63_29 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
5 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AS63_29 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AS63_29 protein").

The nucleotide sequence of the 5' portion of AS63_29 as presently determined is reported in SEQ ID NO:36. What applicants presently believe is the proper reading frame for the coding
10 region is indicated in SEQ ID NO:37. The predicted amino acid sequence of the AS63_29 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:37. Amino acids 28 to 40 of SEQ ID NO:37 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 41. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted
15 leader/signal sequence not be separated from the remainder of the AS63_29 protein. Additional nucleotide sequence from the 3' portion of AS63_29, including a poly(A) tail, is reported in SEQ ID NO:38.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AS63_29 should be approximately 1700 bp.

20 The nucleotide sequence disclosed herein for AS63_29 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AS63_29 demonstrated at least some similarity with sequences identified as L26877 (Mus musculus (B20c) heavy chain immunoglobulin variable region gene), T09146 (EST07039 Homo sapiens cDNA clone HIBBP68 5' end), T23466 (seq3050 Homo sapiens cDNA clone
25 Hy18-Ch13-Charon40-cDNA-100 3'), and W55739 (ma35f05.r1 Life Tech mouse brain Mus musculus cDNA clone 312705 5'). The predicted amino acid sequence disclosed herein for AS63_29 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AS63_29 protein demonstrated at least some sequence similarity with sequences identified as R04032 (Full length T4 encoded by plasmid
30 pBG381). Based upon sequence similarity, AS63_29 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the AS63_29 protein sequence, near the amino terminus.

Clone "AY304_14"

A polynucleotide of the present invention has been identified as clone "AY304_14". AY304_14 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AY304_14 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AY304_14 protein").

The nucleotide sequence of AY304_14 as presently determined is reported in SEQ ID NO:39, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the AY304_14 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:40.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AY304_14 should be approximately 2200 bp.

The nucleotide sequence disclosed herein for AY304_14 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AY304_14 demonstrated at least some similarity with sequences identified as AA127688 (zk92f05.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 490305 3'), AA179609 (zp49g11.r1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 612836 5'), AA276253 (vc40f05.r1 Barstead MPLRB1 Mus musculus cDNA clone 777057 5'), H15545 (ym27d04.s1 Homo sapiens cDNA clone 49495 3' similar to contains PTR5 repetitive element), L08441 (Human autonomously replicating sequence (ARS) mRNA), N34949 (yy49h09.s1 Homo sapiens cDNA clone 276929 3'), R48594 (yj65d07.s1 Homo sapiens cDNA clone 153613 3'), T21160 (Human gene signature HUMGS02466), U43284 (Cloning vector phGFP-S65T, complete sequence, green fluorescent protein (gfp) gene, complete cds), and Z45151 (H. sapiens partial cDNA sequence; clone c-2hh04). The predicted amino acid sequence disclosed herein for AY304_14 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AY304_14 protein demonstrated at least some sequence similarity with sequences identified as D86984 (similar to yeast adenylate cyclase (S56776) [Homo sapiens]), J01415 (cytochrome oxidase subunit 3 [Homo sapiens]), V00662 (cytochrome oxidase III [Homo sapiens]), and X68948 (envelope glycoprotein [Spleen focus-forming virus]). Based upon sequence similarity, AY304_14 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the AY304_14 protein sequence, one centered around amino acid 81 and another around amino acid 120 of SEQ ID NO:40.

35 Clone "BG160_1"

A polynucleotide of the present invention has been identified as clone "BG160_1". BG160_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG160_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BG160_1 protein").

The nucleotide sequence of BG160_1 as presently determined is reported in SEQ ID NO:41, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BG160_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:42. Amino acids 588 to 600 of SEQ ID NO:42 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 601. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BG160_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG160_1 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for BG160_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BG160_1 demonstrated at least some similarity with sequences identified as A60021 (tropomyosin-related protein, neuronal - rat ;contains element MER27 repetitive element), AA081525 (zn20e02.r1 Stratagene neuroepithelium NT2RAMI 937234 Homo sapiens cDNA clone 547994 5'), AA092565 (l15773.seq.F Fetal heart, Lambda ZAP Express Homo sapiens cDNA 5'), D56138 (Human fetal brain cDNA 5'-end GEN-416H11), D61090 (Human fetal brain cDNA 5'-end GEN-155A07), D61184 (Human fetal brain cDNA 5'-end GEN-165A01), L10335 (Homo sapiens neuro-endocrine-specific protein C (NSP) mRNA, complete cds), N21304 (yx53f07.s1 Homo sapiens cDNA clone 265477 3' similar to SP:A60021 A60021 TROPOMYOSIN-RELATED PROTEIN, NEURONAL), and W95814 (ze07f11.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 358317 5' similar to PIR:A60021). The predicted amino acid sequence disclosed herein for BG160_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BG160_1 protein demonstrated at least some sequence similarity with sequences identified as L10334 (neuroendocrine-specific protein B [Homo sapiens]), L10335 (neuroendocrine-specific protein C [Homo sapiens]). Based upon sequence similarity, BG160_1 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer

program predicts three potential transmembrane domains within the BG160_1 protein sequence, centered around amino acids 84, 484, and 595 of SEQ ID NO:42, respectively.

BG160_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 110 kDa was detected in membrane fractions using SDS polyacrylamide gel
5 electrophoresis.

Clone "BO432_4"

A polynucleotide of the present invention has been identified as clone "BO432_4". BO432_4 was isolated from a human adult retina cDNA library using methods which are selective
10 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BO432_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BO432_4 protein").

The nucleotide sequence of the 5' portion of BO432_4 as presently determined is reported
15 in SEQ ID NO:43. An additional internal nucleotide sequence from BO432_4 as presently determined is reported in SEQ ID NO:44. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:45. Additional nucleotide sequence from the 3' portion of BO432_4, including a poly(A) tail, is reported in SEQ ID NO:46.

20 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BO432_4 should be approximately 1700 bp.

The nucleotide sequence disclosed herein for BO432_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BO432_4 demonstrated at least some similarity with sequences identified as
25 AA283626 (zt15e09.s1 Soares NbHTGBC Homo sapiens cDNA clone 713224 3'), AA406486 (zv12g02.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 753458 5' similar to WP F35G2.2 CE05809 E.COLIYCACLKE), AA570446 (nk62c12.s1 NCI_CGAP_Sch1 Homo sapiens cDNA clone IMAGE:1018102), N55855 (J3389F Homo sapiens cDNA clone J3389 5'), Q10613 (Rianodin receptor gene), T62691 (yc70d10.r1 Homo sapiens cDNA clone 86035 5'), and W90766
30 (zh79h04.s1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 418327 3'). The predicted amino acid sequence disclosed herein for BO432_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BO432_4 protein demonstrated at least some sequence similarity with sequences identified as Z69637 (F35G2.2 [Caenorhabditis elegans]). Based upon sequence similarity, BO432_4 proteins
35 and each homologous protein or peptide may share at least some activity. The TopPredII

computer program predicts a potential transmembrane domain at the amino terminus of the BO432_4 protein sequence. The BO432_4 protein may also contain the bacterial lysR family signature, a motif found in bacterial transcriptional regulators and which is possibly indicative of a helix-turn-helix structure.

5

Clone "BO538_2"

A polynucleotide of the present invention has been identified as clone "BO538_2". BO538_2 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
10 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BO538_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BO538_2 protein").

The nucleotide sequence of the 5' portion of BO538_2 as presently determined is reported in SEQ ID NO:47. What applicants presently believe is the proper reading frame for the coding
15 region is indicated in SEQ ID NO:48. The predicted amino acid sequence of the BO538_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Additional nucleotide sequence from the 3' portion of BO538_2, including a poly(A) tail, is reported in SEQ ID NO:49.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
20 BO538_2 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for BO538_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BO538_2 demonstrated at least some similarity with sequences identified as AA503100 (ne44h01.s1 NCI_CGAP_Co3 Homo sapiens cDNA clone 900241), R44035
25 (yg21g09.s1 Homo sapiens cDNA clone 33167 3'), T21630 (Human gene signature HUMGS03066), and W64854 (me06d12.r1 Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone 386711 5' similar to PIR S40989 S40989 hypothetical protein F55H2.6 - Caenorhabditis elegans). The predicted amino acid sequence disclosed herein for BO538_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX
30 search protocol. The predicted BO538_2 protein demonstrated at least some sequence similarity with sequences identified as M60525 (nerve growth factor inducible protein [Rattus norvegicus]), R28916 (Type III procollagen), and Z27080 (F55H2.6 [Caenorhabditis elegans]). Based upon sequence similarity, BO538_2 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains
35 within the BO538_2 protein sequence.

Clone "BR595_4"

A polynucleotide of the present invention has been identified as clone "BR595_4". BR595_4 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
5 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BR595_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BR595_4 protein").

The nucleotide sequence of the 5' portion of BR595_4 as presently determined is reported in SEQ ID NO:50. What applicants presently believe is the proper reading frame for the coding
10 region is indicated in SEQ ID NO:51. The predicted amino acid sequence of the BR595_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:51. Additional nucleotide sequence from the 3' portion of BR595_4, including a poly(A) tail, is reported in SEQ ID NO:52.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
15 BR595_4 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for BR595_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BR595_4 demonstrated at least some similarity with sequences identified as AA443742 (zw95b02.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 784683 3'),
20 AA600820 (np45b08.s1 NCI_CGAP_Br1.1 Homo sapiens cDNA clone IMAGE:1129239), T19410 (Human gene signature HUMGS00435), W87465 (zh67c04.s1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 417126 3'), and Z33587 (H. sapiens partial cDNA sequence; clone HEA89P; single read). Based upon sequence similarity, BR595_4 proteins and each homologous protein or peptide may share at least some activity.

25

Clone "CI490_2"

A polynucleotide of the present invention has been identified as clone "CI490_2". CI490_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
30 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CI490_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CI490_2 protein").

The nucleotide sequence of CI490_2 as presently determined is reported in SEQ ID NO:53, and includes a poly(A) tail. What applicants presently believe to be the proper reading
35 frame and the predicted amino acid sequence of the CI490_2 protein corresponding to the

foregoing nucleotide sequence is reported in SEQ ID NO:54. Amino acids 64 to 76 of SEQ ID NO:54 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 77. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CI490_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CI490_2 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for CI490_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CI490_2 demonstrated at least some similarity with sequences identified as H30751 (yo79a04.r1 Homo sapiens cDNA clone 184110 5'), H49766 (yo24f01.r1 Homo sapiens cDNA clone 178873 5' similar to SP:S19586 N-METHYL-D-ASPARTATE RECEPTOR GLUTAMATE-BINDING CHAIN), H51158 (yo32d04.r1 Homo sapiens cDNA clone 179623 5'), R85211 (yo41d11.s1 Homo sapiens cDNA clone 180501 3' similar to SP S19586 N-METHYL-D-ASPARTATE RECEPTOR GLUTAMATE-BINDING CHAIN), S19586 (N-METHYL-D-ASPARTATE RECEPTOR GLUTAMATE-BINDING CHAIN), S61973 (NMDA receptor glutamate-binding subunit [rat, mRNA]), T01031 (Human leucine zipper protein-kinase cDNA sequence), and W56893 (zc01g05.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 321080 5' similar to PIR S19586 S19586 N-methyl-D-aspartate receptor glutamate-binding chain - rat). The predicted amino acid sequence disclosed herein for CI490_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CI490_2 protein demonstrated at least some sequence similarity with sequences identified as S61973 (NMDA receptor glutamate-binding subunit [Rattus sp.]) and U08020 (collagen pro-alpha-1 type I chain [Mus musculus]). Based upon sequence similarity, CI490_2 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts six potential transmembrane domains within the CI490_2 protein sequence, with the most amino-terminal transmembrane domain centered around amino acid 77 of SEQ ID NO:54.

30 Clone "CI522_1"

A polynucleotide of the present invention has been identified as clone "CI522_1". CI522_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. CI522_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CI522_1 protein").

The nucleotide sequence of the 5' portion of CI522_1 as presently determined is reported in SEQ ID NO:55. What applicants presently believe is the proper reading frame for the coding
5 region is indicated in SEQ ID NO:56. The predicted amino acid sequence of the CI522_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56. Amino acids 7 to 19 of SEQ ID NO:56 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted
10 leader/signal sequence not be separated from the remainder of the CI522_1 protein. Additional nucleotide sequence from the 3' portion of CI522_1, including a poly(A) tail, is reported in SEQ ID NO:57.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CI522_1 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for CI522_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CI522_1 demonstrated at least some similarity with sequences identified as AA028557 (mi18g05.r1 Soares mouse p3NMF19.5 Mus musculus cDNA clone 463928 5'), H32238 (EST107136 Rattus sp. cDNA 5' end), T33525 (EST58140 Homo sapiens cDNA 5' end similar
20 to None), U66468 (Human cell growth regulator CGR11 mRNA, complete cds), and X00525 (Mouse 28S ribosomal RNA). The predicted amino acid sequence disclosed herein for CI522_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CI522_1 protein demonstrated at least some sequence similarity with sequences identified as U66468 (cell growth regulator CGR11 [Homo sapiens]).
25 Based upon sequence similarity, CI522_1 proteins and each homologous protein or peptide may share at least some activity.

Clone "CN238_1"

A polynucleotide of the present invention has been identified as clone "CN238_1".
30 CN238_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CN238_1 includes at least a portion of the coding sequence of a secreted protein (also referred to herein as "CN238_1 protein").

The nucleotide sequence of CN238_1 as presently determined is reported in SEQ ID NO:58, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CN238_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:59.

- 5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CN238_1 should be approximately 2190 bp.

The nucleotide sequence disclosed herein for CN238_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CN238_1 demonstrated at least some similarity with sequences identified as
10 AA044097 (zk51b02.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 486315 5'),
AA044287 (zk51b02.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 486315 3'),
AA045440 (zk67c03.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 487876 3'),
AA143007 (zl48f01.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 505177 5'),
D51196 (Human fetal brain cDNA 3'-end GEN-016G05), D60310 (Human fetal brain cDNA
15 3'-end GEN-098A09), N69344 (yz43e04.s1 Homo sapiens cDNA clone 285822 3' similar to
gb:K00558 TUBULIN ALPHA-1 CHAIN (HUMAN)), W22250 (64B8 Human retina cDNA
Tsp509I-cleaved sublibrary Homo), and X01703 (Human gene for alpha-tubulin (b alpha 1)). The
predicted amino acid sequence disclosed herein for CN238_1 was searched against the GenPept
and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted
20 CN238_1 protein demonstrated at least some sequence similarity with sequences identified as
K00557 (alpha-tubulin [Homo sapiens]) and U51583 (zinc finger homeodomain enhancer-binding
protein-1 [Rattus norvegicus]). Based upon sequence similarity, CN238_1 proteins and each
homologous protein or peptide may share at least some activity.

25 Clone "CO390_1"

A polynucleotide of the present invention has been identified as clone "CO390_1". CO390_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence
30 of the encoded protein. CO390_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CO390_1 protein").

The nucleotide sequence of CO390_1 as presently determined is reported in SEQ ID NO:60, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CO390_1 protein corresponding to the
35 foregoing nucleotide sequence is reported in SEQ ID NO:61.

-286-

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CO390_1 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for CO390_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CO390_1 demonstrated at least some similarity with sequences identified as H84353 (y85a11.r1 Homo sapiens cDNA clone 249500 5'), L35532 (Pan troglodytes Alu repeat region), N80616 (Genomic clone encoding SAP(Phe)), R53922 (y103h10.s1 Homo sapiens cDNA clone 138211 3' similar to contains Alu repetitive element; contains TAR1 repetitive element), X75335 (H.sapiens Alu insertion in COL3A1 gene), X95882 (R.norvegicus mRNA for ATP ligand gated ion channel), and Y09561 (H.sapiens mRNA for P2X7 receptor). The predicted amino acid sequence disclosed herein for CO390_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CO390_1 protein demonstrated at least some sequence similarity with sequences identified as U45448 (P2x1 receptor [Homo sapiens]), W04216 (Rat superior cervical ganglion p2x receptor), X83688 (ATP receptor [Homo sapiens]), X95882 (P2X7 gene product [Rattus norvegicus]), and Y09561 (ATP receptor [Homo sapiens]). Based upon sequence similarity, CO390_1 proteins and each homologous protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the CO390_1 protein sequence, centered around amino acid 249 of SEQ ID NO:61. The nucleotide sequence of CO390_1 may contain an Alu repetitive element.

CO390_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 75 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

Clone "AJ20_2"

A polynucleotide of the present invention has been identified as clone "AJ20_2". AJ20_2 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AJ20_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AJ20_2 protein").

The nucleotide sequence of the 5' portion of AJ20_2 as presently determined is reported in SEQ ID NO:62. What applicants presently believe is the proper reading frame for the coding

region is indicated in SEQ ID NO:63. The predicted amino acid sequence of the AJ20_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:63. Amino acids 8 to 20 of SEQ ID NO:63 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AJ20_2 protein. Additional nucleotide sequence from the 3' portion of AJ20_2, including a poly(A) tail, is reported in SEQ ID NO:64.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AJ20_2 should be approximately 850 bp.

The nucleotide sequence disclosed herein for AJ20_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

Clone "AR440_1"

A polynucleotide of the present invention has been identified as clone "AR440_1". AR440_1 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AR440_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AR440_1 protein").

The partial nucleotide sequence of AR440_1, including its 3' end and a poly(A) tail, as presently determined is reported in SEQ ID NO:66. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:67. The predicted amino acid sequence of the AR440_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:67. Additional nucleotide sequence from the 5' portion of AR440_1 is reported in SEQ ID NO:65.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AR440_1 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for AR440_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database. The nucleotide sequence of AR440_1 indicates that it may contain an Alu repetitive element.

Clone "AS164_1"

A polynucleotide of the present invention has been identified as clone "AS164_1". AS164_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AS164_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AS164_1 protein").

The nucleotide sequence of AS164_1 as presently determined is reported in SEQ ID NO:68, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the AS164_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:69.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AS164_1 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for AS164_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AS164_1 demonstrated at least some similarity with sequences identified as H24668 (yl40h10.r1 Homo sapiens cDNA clone 160771 5'), N29757 (yw90h10.s1 Homo sapiens cDNA clone 259555 3'), T62184 (yb96d08.r1 Homo sapiens cDNA clone 79023 5'), Z69706 (Human DNA sequence from cosmid COS12 from a contig from the tip of the short arm of chromosome 16, spanning 2Mb of 16p13.3. Contains ESTs, Flanking sequences of 3' alpha globin H), and Z69890 (Human DNA sequence from cosmid RJ14 from a contig from the tip of the short arm of chromosome 16, spanning 2Mb of 16p13.3. Contains ESTs and CpG island). The predicted amino acid sequence disclosed herein for AS164_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AS164_1 protein demonstrated at least some similarity to sequences identified as A20359_1 (ryanodine receptor gene product [Homo sapiens]) and U78866 (putative arginine-aspartate-rich RNA binding protein [Arabidopsis thaliana]). Based upon sequence similarity, AS164_1 proteins and each similar protein or peptide may share at least some activity. The predicted AS164_1 protein sequence also contains repeated Asp-Arg RNA-binding motifs.

30 Clone "AX8_1"

A polynucleotide of the present invention has been identified as clone "AX8_1". AX8_1 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. AX8_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AX8_1 protein").

The nucleotide sequence of AX8_1 as presently determined is reported in SEQ ID NO:70, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the AX8_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:71. Amino acids 106 to 118 of SEQ ID NO:71 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 119. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AX8_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AX8_1 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for AX8_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database. The TopPredII computer program predicts three potential transmembrane domains within the AX8_1 protein sequence, centered around amino acids 111, 144, and 182 of SEQ ID NO:71, respectively.

AX8_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 35 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "BD176_3"

A polynucleotide of the present invention has been identified as clone "BD176_3". BD176_3 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BD176_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BD176_3 protein").

The nucleotide sequence of the 5' portion of BD176_3 as presently determined is reported in SEQ ID NO:72. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:73. The predicted amino acid sequence of the BD176_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:73. Amino acids 2 to 14 of SEQ ID NO:73 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted

leader/signal sequence not be separated from the remainder of the BD176_3 protein. Additional nucleotide sequence from the 3' portion of BD176_3, including a poly(A) tail, is reported in SEQ ID NO:74.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
5 BD176_3 should be approximately 1300 bp.

The nucleotide sequence disclosed herein for BD176_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BD176_3 demonstrated at least some similarity with sequences identified as AA029679 (ze94g10.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 366690 5'),
10 D45913 (Mouse NLRR-1 mRNA for leucine-rich-repeat protein, complete cds), R55610 (yg88h08.r1 Homo sapiens cDNA clone 40606 5'), and T07640 (EST05530 Homo sapiens cDNA clone HFBEM16). The predicted amino acid sequence disclosed herein for BD176_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BD176_3 protein demonstrated at least some similarity to
15 sequences identified as D45913 (leucine-rich-repeat protein [Mus musculus]) and M59472 (asparagine-rich antigen Pfa55-6 [Plasmodium falciparum]). Based upon sequence similarity, BD176_3 proteins and each similar protein or peptide may share at least some activity.

Clone "BD339_1"

20 A polynucleotide of the present invention has been identified as clone "BD339_1". BD339_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BD339_1 is a full-length clone, including the entire coding
25 sequence of a secreted protein (also referred to herein as "BD339_1 protein").

The nucleotide sequence of BD339_1 as presently determined is reported in SEQ ID NO:75, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BD339_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76.

30 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BD339_1 should be approximately 650 bp.

The nucleotide sequence disclosed herein for BD339_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BD339_1 demonstrated at least some similarity with sequences identified as H82422
35 (yu80d08.s1 Homo sapiens cDNA clone 240111 3), N62058 (EST53c05 Homo sapiens cDNA

clone), U21730 Human 5'-nucleotidase (CD73)), W01979 (za30h09.r1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 294113 5'), and W02015 (za32b11.r1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 294237 5'). Based upon sequence similarity, BD339_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII
5 computer program predicts three potential transmembrane domains within the BD339_1 protein sequence, centered around amino acids 14, 46, and 76 of SEQ ID NO:76, respectively.

Clone "BD427_1"

A polynucleotide of the present invention has been identified as clone "BD427_1".
10 BD427_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BD427_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BD427_1 protein").

15 The nucleotide sequence of BD427_1 as presently determined is reported in SEQ ID NO:77, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BD427_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
20 BD427_1 should be approximately 1810 bp.

The nucleotide sequence disclosed herein for BD427_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BD427_1 demonstrated at least some similarity with sequences identified as BD427_1 demonstrated at least some similarity with sequences identified as AA027122 (zk04a03.r1 Soares
25 pregnant uterus NbHPU Homo sapiens cDNA clone 469516 5'), N24735 (yx56b02.s1 Homo sapiens cDNA clone 265707 3'), and W84644 (zd91a06.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 356818 5'). Based upon sequence similarity, BD427_1 proteins and each similar protein or peptide may share at least some activity.

Clone "BL229_22"

30 A polynucleotide of the present invention has been identified as clone "BL229_22". BL229_22 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino

acid sequence of the encoded protein. BL229_22 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BL229_22 protein").

The nucleotide sequence of the 5' portion of BL229_22 as presently determined is reported in SEQ ID NO:79. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:80. The predicted amino acid sequence of the BL229_22 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:80. Additional nucleotide sequence from the 3' portion of BL229_22, including a poly(A) tail, is reported in SEQ ID NO:81.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BL229_22 should be approximately 870 bp.

The nucleotide sequence disclosed herein for BL229_22 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

15 Clone "BV123_16"

A polynucleotide of the present invention has been identified as clone "BV123_16". BV123_16 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV123_16 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV123_16 protein").

The nucleotide sequence of BV123_16 as presently determined is reported in SEQ ID NO:82, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BV123_16 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:83.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV123_16 should be approximately 1080 bp.

The nucleotide sequence disclosed herein for BV123_16 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV123_16 demonstrated at least some similarity with sequences identified as H29610 (ym61e03.s1 Homo sapiens cDNA clone 52653 3'), H52374 (yq81b12.r1 Homo sapiens cDNA clone 202175 5'), H66213 (yu16h10.s1 Homo sapiens cDNA), L08092 (Homo sapiens dystrophin (DMD) gene, intron 7, transposon-like sequence), L35670 (Homo sapiens (subclone H8 10_g5 from P1 35 H5 C8) DNA sequence), M62716 (Human CSP-B gene flanking sequence), N46985 (yy83a05.s1 Homo sapiens cDNA clone 280112 3'), R94603 (yq38a04.s1 Homo sapiens

cDNA clone 198030 3'), U91321 (Human chromosome 16p13 BAC clone CIT987SK-363E6, complete sequence), and Z82200 (Human DNA sequence from clone J333E231). Based upon sequence similarity, BV123_16 proteins and each similar protein or peptide may share at least some activity.

5

Clone "CH377_1"

A polynucleotide of the present invention has been identified as clone "CH377_1". CH377_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CH377_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CH377_1 protein").

The nucleotide sequence of CH377_1 as presently determined is reported in SEQ ID NO:84, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CH377_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:85. Amino acids 5 to 17 of SEQ ID NO:85 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CH377_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CH377_1 should be approximately 570 bp.

The nucleotide sequence disclosed herein for CH377_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CH377_1 demonstrated at least some similarity with sequences identified as AA507382 (nh73b01.s1 NCI_CGAP_Br1.1 Homo sapiens cDNA clone IMAGE 964105) and N70479 (za74f12.s1 Homo sapiens cDNA clone 298319 3'). Based upon sequence similarity, CH377_1 proteins and each similar protein or peptide may share at least some activity.

30

Clone "BD441_1"

A polynucleotide of the present invention has been identified as clone "BD441_1". BD441_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino

35

acid sequence of the encoded protein. BD441_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BD441_1 protein").

The nucleotide sequence of the 5' portion of BD441_1 as presently determined is reported in SEQ ID NO:86. An additional internal nucleotide sequence from BD441_1 as presently
5 determined is reported in SEQ ID NO:87. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:88. Additional nucleotide sequence from the 3' portion of BD441_1, including a poly(A) tail, is reported in SEQ ID NO:89.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
10 BD441_1 should be approximately 2400 bp.

The predicted amino acid sequence disclosed herein for BD441_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BD441_1 protein demonstrated at least some similarity to sequences identified as X61615 (leukemia inhibitory factor receptor [Homo sapiens]). Based upon sequence similarity,
15 BD441_1 proteins and each similar protein or peptide may share at least some activity.

Clone "BD441_2"

A polynucleotide of the present invention has been identified as clone "BD441_2". BD441_2 was isolated from a human fetal kidney cDNA library using methods which are
20 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BD441_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BD441_2 protein").

The nucleotide sequence of the 5' portion of BD441_2 as presently determined is reported
25 in SEQ ID NO:90. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:91. The predicted amino acid sequence of the BD441_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:91. Additional nucleotide sequence from the 3' portion of BD441_2, including a poly(A) tail, is reported in SEQ ID NO:92.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
30 BD441_2 should be approximately 1200 bp.

The predicted amino acid sequence disclosed herein for BD441_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BD441_2 protein demonstrated at least some similarity to sequences identified as

X61615 (leukemia inhibitory factor receptor [Homo sapiens]). Based upon sequence similarity, BD441_2 proteins and each similar protein or peptide may share at least some activity.

Clone "BG102_3"

5 A polynucleotide of the present invention has been identified as clone "BG102_3". BG102_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG102_3 is a full-length clone, including the entire coding sequence of
10 a secreted protein (also referred to herein as "BG102_3 protein").

The nucleotide sequence of BG102_3 as presently determined is reported in SEQ ID NO:93, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BG102_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:94. Amino acids 11 to 23 of SEQ ID
15 NO:94 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BG102_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
20 BG102_3 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for BG102_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BG102_3 demonstrated at least some similarity with sequences identified as AC002078 (Human BAC clone RG111H14 from 7q22, complete sequence), L11910 (Human
25 retinoblastoma susceptibility gene exons 1-27, complete cds), U62317 (Chromosome 22q13 BAC Clone CIT987SK-384D8 complete sequence), Z54147 (Human DNA sequence from cosmid L129H7, Huntington's Disease Region, chromosome 4p16.3 contains CpG island), Z75747 (Human DNA sequence from cosmid U96H1, between markers DXS366 and DXS87 on chromosome X *), and Z80899 (Human DNA sequence from cosmid F1121 on chromosome 6).

30 The predicted amino acid sequence disclosed herein for BG102_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BG102_3 protein demonstrated at least some similarity to sequences identified as M13100 (unknown protein [Rattus norvegicus]) and U15647 (reverse transcriptase [Mus musculus]). Based upon sequence similarity, BG102_3 proteins and each similar protein or

peptide may share at least some activity. The nucleotide sequence of BG102_3 indicates that it may contain an L1 repetitive element.

BG102_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 55 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

Clone "BK158_1"

A polynucleotide of the present invention has been identified as clone "BK158_1". BK158_1 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BK158_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BK158_1 protein").

The nucleotide sequence of BK158_1 as presently determined is reported in SEQ ID NO:95, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BK158_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:96.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BK158_1 should be approximately 1150 bp.

The nucleotide sequence disclosed herein for BK158_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BK158_1 demonstrated at least some similarity with sequences identified as N39195 (yv26e08.s1 Homo sapiens cDNA clone 243878 3') and N45263 (yv26e08.r1 Homo sapiens cDNA clone 243878 5'). Based upon sequence similarity, BK158_1 proteins and each similar protein or peptide may share at least some activity.

BK158_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 28 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "BP163_1"

A polynucleotide of the present invention has been identified as clone "BP163_1". BP163_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. BP163_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BP163_1 protein").

The nucleotide sequence of the 5' portion of BP163_1 as presently determined is reported in SEQ ID NO:97. What applicants presently believe is the proper reading frame for the coding
5 region is indicated in SEQ ID NO:98. The predicted amino acid sequence of the BP163_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:98. Additional nucleotide sequence from the 3' portion of BP163_1, including a poly(A) tail, is reported in SEQ ID NO:99.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
10 BP163_1 should be approximately 1240 bp.

The nucleotide sequence disclosed herein for BP163_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BP163_1 demonstrated at least some similarity with sequences identified as AA187086 (zp58h06.r1 Stratagene endothelial cell 937223 Homo sapiens cDNA clone 624443 5' similar to
15 TR G285943 G285943 ORF, COMPLETE CDS), AA301506 (EST14475 Testis tumor Homo sapiens cDNA 5' end similar to hypothetical protein (GB D14659)), D14659 (Human mRNA for KIAA0103 gene, complete cds), and W57328 (ma26d10.r1 Life Tech mouse brain Mus musculus cDNA clone). The predicted amino acid sequence disclosed herein for BP163_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search
20 protocol. The predicted BP163_1 protein demonstrated at least some similarity to sequences identified as D14659 (KIAA0103 [Homo sapiens]). Based upon sequence similarity, BP163_1 proteins and each similar protein or peptide may share at least some activity.

Clone "BZ16_3"

25 A polynucleotide of the present invention has been identified as clone "BZ16_3". BZ16_3 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BZ16_3 is a full-length clone, including the entire coding sequence of a
30 secreted protein (also referred to herein as "BZ16_3 protein").

The partial nucleotide sequence of BZ16_3, including its 3' end and a poly(A) tail, as presently determined is reported in SEQ ID NO:101. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:102. The predicted amino acid sequence of the BZ16_3 protein corresponding to the foregoing nucleotide sequence is

reported in SEQ ID NO:102. Additional nucleotide sequence from the 5' portion of BZ16_3 is reported in SEQ ID NO:100.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BZ16_3 should be approximately 2120 bp.

5 The nucleotide sequence disclosed herein for BZ16_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BZ16_3 demonstrated at least some similarity with sequences identified as F06886 (H. sapiens partial cDNA sequence; clone c-1nf02), F06870 (H. sapiens partial cDNA sequence; clone c-1nc11), N53511 (yz26b08.s1 Homo sapiens cDNA clone 284151 3'), T65313 (yc79g12.s1
10 Homo sapiens cDNA clone 22132 3'), U00084 (Haemophilus influenzae), W44815 (zc21d01.s1 Soares senescent fibroblasts NbHSF Homo sapiens cDNA clone 322945 3'), and Z49128 (Caenorhabditis elegans cosmid M03C11). The predicted amino acid sequence disclosed herein for BZ16_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BZ16_3 protein demonstrated at least some
15 similarity to sequences identified as D26185 (cell division protein [Bacillus subtilis]), L46096 (HEAH11465_1 cell division protein [Haemophilus influenzae]), and Z49128 (CEM03C11_5 M03C11.5 [Caenorhabditis elegans]). The BZ16_3 protein demonstrated at least some similarity to ATP-dependent proteases such as ftsH. Based upon sequence similarity, BZ16_3 proteins and each similar protein or peptide may share at least some activity.

20

Clone "CC182_1"

A polynucleotide of the present invention has been identified as clone "CC182_1". CC182_1 was isolated from a human adult brain cDNA library using methods which are selective
25 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CC182_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CC182_1 protein").

The nucleotide sequence of CC182_1 as presently determined is reported in SEQ ID
30 NO:103, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CC182_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:104. Amino acids 26 to 38 of SEQ ID NO:104 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 39. Due to the hydrophobic nature of the predicted leader/signal

sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CC182_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CC182_1 should be approximately 1600 bp.

5 The nucleotide sequence disclosed herein for CC182_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CC182_1 demonstrated at least some similarity with sequences identified as H61159 (yu37f08.s1 Homo sapiens cDNA clone 236007 3' similar to contains L1 repetitive element), L09709 (Human lysosomal-associated membrane glycoprotein-2 (LAMP2) gene, 5' end of CDS
10 and flanking region), W44797 (zb98e10.s1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 320874 3' similar to contains Alu repetitive element), and X62167 (H.sapiens mRNA for P2 protein of peripheral myelin). Based upon sequence similarity, CC182_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of CC182_1 indicates that it may contain an L1 repetitive element and/or a MER42C repetitive element.

15

Clone "CG109_1"

A polynucleotide of the present invention has been identified as clone "CG109_1". CG109_1 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
20 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CG109_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CG109_1 protein").

The nucleotide sequence of CG109_1 as presently determined is reported in SEQ ID NO:105, and includes a poly(A) tail. What applicants presently believe to be the proper reading
25 frame and the predicted amino acid sequence of the CG109_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:106.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CG109_1 should be approximately 600 bp.

The nucleotide sequence disclosed herein for CG109_1 was searched against the GenBank
30 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

Clone "CJ397_1"

A polynucleotide of the present invention has been identified as clone "CJ397_1".
35 CJ397_1 was isolated from a human fetal brain cDNA library using methods which are selective

for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CJ397_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CJ397_1 protein").

5 The nucleotide sequence of the 5' portion of CJ397_1 as presently determined is reported in SEQ ID NO:107. An additional internal nucleotide sequence from CJ397_1 as presently determined is reported in SEQ ID NO:108. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:109. Additional nucleotide sequence from the 3' portion of CJ397_1, including a poly(A)
10 tail, is reported in SEQ ID NO:110.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CJ397_1 should be approximately 1900 bp.

The nucleotide sequence disclosed herein for CJ397_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
15 protocols. CJ397_1 demonstrated at least some similarity with sequences identified as H18685 (yn52b08.s1 Homo sapiens cDNA clone 172023 3'), H46001 (yo13f06.s1 Homo sapiens cDNA clone 177827 3'), and T77612 (yc91f06.r1 Homo sapiens cDNA clone 23298 5'). Based upon sequence similarity, CJ397_1 proteins and each similar protein or peptide may share at least some activity.

20

Clone "AM795_4"

A polynucleotide of the present invention has been identified as clone "AM795_4". AM795_4 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified
25 as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AM795_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AM795_4 protein").

The nucleotide sequence of AM795_4 as presently determined is reported in SEQ ID NO:111, and includes a poly(A) tail. What applicants presently believe to be the proper reading
30 frame and the predicted amino acid sequence of the AM795_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:112. Amino acids 9 to 21 of SEQ ID NO:112 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 22. Amino acids 138 to 150 of SEQ ID NO:112 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid
35 151. Due to the hydrophobic nature of the predicted and the possible leader/signal sequences,

each of such sequences may act as a transmembrane domain should that leader/signal sequence not be separated from the remainder of the AM795_4 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM795_4 should be approximately 1900 bp.

5 The nucleotide sequence disclosed herein for AM795_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AM795_4 demonstrated at least some similarity with sequences identified as AF002700 (Homo sapiens GDNF family receptor alpha 2 (GFRalpha2) mRNA, complete cds), H05619 (yl70a10.s1 Homo sapiens cDNA clone 43207 3'), U46493 (Cloning vector pFlp recombina

10 se gene, complete cds), U59486 (Rattus norvegicus GDNF receptor alpha mRNA, complete cds), V00248 (Human Ret ligand retL2 cDNA), W73633 (zd55h01.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344593 3', mRNA sequence), and W73681 (zd55h01.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344593 5', mRNA sequence). The predicted amino acid sequence disclosed herein for AM795_4 was searched against the GenPept

15 and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AM795_4 protein demonstrated at least some similarity to sequences identified as AF002700 (GDNF family receptor alpha 2 (GFRalpha2) [Homo sapiens]), U59486 (GDNF receptor alpha [Rattus norvegicus]), and W37460 (Human Ret ligand retL2 cDNA). A receptor complex comprised of TrnR1 (GDNFR alpha) and Ret was found to be capable of mediating both GDNF

20 and NTN signaling. The receptor called TrnR2, identified based on homology to TrnR1, is 48% identical to TrnR1 and is encoded by a gene located on the short arm of chromosome 8. TrnR2 is attached to the cell surface via a GPI-linkage, and can mediate both NTN and GDNF signaling through Ret *in vitro* (Baloh *et al.*, 1997, *Neuron* 18(5): 793-802, which is incorporated by reference herein). Based upon sequence similarity, AM795_4 proteins and each similar protein

25 or peptide may share at least some activity.

AM795_4 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 76 kDa was detected in conditioned media fractions using SDS polyacrylamide gel electrophoresis.

30 Clone "AT340_1"

A polynucleotide of the present invention has been identified as clone "AT340_1". AT340_1 was isolated from a human adult blood (lymphocytes and dendritic cells treated with mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or

35 transmembrane protein on the basis of computer analysis of the amino acid sequence of the

encoded protein. AT340_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AT340_1 protein").

The partial nucleotide sequence of AT340_1, including its 3' end and a poly(A) tail, as presently determined is reported in SEQ ID NO:114. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:115. The predicted amino acid sequence of the AT340_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:115. Amino acids 12 to 24 of SEQ ID NO:115 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AT340_1 protein. Additional nucleotide sequence from the 5' portion of AT340_1 is reported in SEQ ID NO:113.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AT340_1 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for AT340_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AT340_1 demonstrated at least some similarity with sequences identified as AA039343 (zk39g04.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 485238 3'), R68951 (yi43g06.r1 Homo sapiens cDNA clone 142042 5' similar to SP:C35D10.1 CE01190), R77532 (yi76c01.r1 Homo sapiens cDNA), R92619 (yq04a04.r1 Homo sapiens cDNA clone 195918 5' similar to SP:C35D10.1 CE01190), and W60997 (zc99f09.s1 Pancreatic Islet Homo sapiens cDNA clone 339305 3'). The predicted amino acid sequence disclosed herein for AT340_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AT340_1 protein demonstrated at least some similarity to sequences identified as U21324 (similar to *S. cerevisiae* hypothetical protein YKL166 [*Caenorhabditis elegans*]). Based upon sequence similarity, AT340_1 proteins and each similar protein or peptide may share at least some activity.

Clone "BG132_1"

A polynucleotide of the present invention has been identified as clone "BG132_1". BG132_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG132_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BG132_1 protein").

The nucleotide sequence of the 5' portion of BG132_1 as presently determined is reported in SEQ ID NO:116. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:117. The predicted amino acid sequence of the BG132_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:117.

- 5 Amino acids 121 to 133 of SEQ ID NO:117 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 134. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BG132_1 protein. Additional nucleotide sequence from the 3' portion of BG132_1, including a poly(A) tail, is reported in SEQ ID NO:118.
- 10

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG132_1 should be approximately 2000 bp.

- The nucleotide sequence disclosed herein for BG132_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BG132_1 demonstrated at least some similarity with sequences identified as
- 15 AA078587 (7P05H12 Chromosome 7 Placental cDNA Library Homo sapiens cDNA clone 7P05H12), H14301 (ym63c04.r1 Homo sapiens cDNA clone 163590 5' similar to gb:U03642_cds1 PROBABLE G PROTEIN-COUPLED RECEPTOR APJ (HUMAN)), L09249 (putative G-protein coupled receptor, rhodopsin family), S79811 (adrenomedullin receptor [rats, lung, mRNA]), T36034 (rchd523 gene differentially expressed in cardiovascular disease), U58828 (Human IL8-related receptor (DRY12) mRNA, complete cds), and Y08162 (H.sapiens mRNA for heptahelix receptor). The predicted amino acid sequence disclosed herein for BG132_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BG132_1 protein demonstrated at least some similarity to
- 20 sequences identified as L06109 (G protein-coupled receptor [Gallus gallus]), L34339 (galanin receptor [Homo sapiens]), U30290 (galanin receptor GALR1 [Rattus norvegicus]), U58828 (IL8-related receptor [Homo sapiens]), W03739 (rchd523 gene product (G protein-coupled receptor)), X98510 (G protein-coupled receptor [Homo sapiens]), and Y08162 (heptahelix receptor [Homo sapiens]). Based upon sequence similarity, BG132_1 proteins and each similar
- 25 protein or peptide may share at least some activity.
- 30

Clone "BG219_2"

- A polynucleotide of the present invention has been identified as clone "BG219_2". BG219_2 was isolated from a human adult brain cDNA library using methods which are selective
- 35 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding

a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG219_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BG219_2 protein").

5 The nucleotide sequence of BG219_2 as presently determined is reported in SEQ ID NO:119, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BG219_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:120.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG219_2 should be approximately 700 bp.

10 The nucleotide sequence disclosed herein for BG219_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BG219_2 demonstrated at least some similarity with sequences identified as AA210695 (zr88b05.s1 Soares NbHTGBC Homo sapiens cDNA clone 682737 3'), C01459 (HUMGS0008450, Human Gene Signature, 3'-directed cDNA sequence), N22628 (EST49p115
15 Homo sapiens cDNA clone 49p115), and T26211 (Human gene signature HUMGS08450). Based upon sequence similarity, BG219_2 proteins and each similar protein or peptide may share at least some activity.

Clone "BG366_2"

20 A polynucleotide of the present invention has been identified as clone "BG366_2". BG366_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BG366_2 is a full-length clone, including the entire coding sequence of
25 a secreted protein (also referred to herein as "BG366_2 protein").

The nucleotide sequence of BG366_2 as presently determined is reported in SEQ ID NO:121, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BG366_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:122. The amino acid sequence of
30 another protein that could be encoded by BG366_2 is reported in SEQ ID NO:283.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG366_2 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for BG366_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
35 protocols. BG366_2 demonstrated at least some similarity with sequences identified as N39453

(yy49h03.s1 Homo sapiens cDNA clone 276917 3'). Based upon sequence similarity, BG366_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the BG366_2 protein sequence centered around amino acid 92 of SEQ ID NO:122.

5

Clone "BV172_2"

A polynucleotide of the present invention has been identified as clone "BV172_2". BV172_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
10 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV172_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV172_2 protein").

The nucleotide sequence of BV172_2 as presently determined is reported in SEQ ID NO:123, and includes a poly(A) tail. What applicants presently believe to be the proper reading
15 frame and the predicted amino acid sequence of the BV172_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:124.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV172_2 should be approximately 1650 bp.

The nucleotide sequence disclosed herein for BV172_2 was searched against the GenBank
20 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV172_2 demonstrated at least some similarity with sequences identified as No significant hits were found in the database. The TopPredII computer program predicts a potential transmembrane domain within the BV172_2 protein sequence centered around amino acid 19 of SEQ ID NO:124. The nucleotide sequence of BV172_2 indicates that it may contain one or more
25 of the following types of repetitive elements: an element similar to chicken CR1, human L1, Mer33.

30 Clone "CC247_10"

A polynucleotide of the present invention has been identified as clone "CC247_10". CC247_10 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino

acid sequence of the encoded protein. CC247_10 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CC247_10 protein").

The nucleotide sequence of CC247_10 as presently determined is reported in SEQ ID NO:125, and includes a poly(A) tail. What applicants presently believe to be the proper reading
5 frame and the predicted amino acid sequence of the CC247_10 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:126. Amino acids 1 to 8 of SEQ ID NO:126 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 9. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be
10 separated from the remainder of the CC247_10 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CC247_10 should be approximately 550 bp.

The nucleotide sequence disclosed herein for CC247_10 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA
15 search protocols. CC247_10 demonstrated at least some similarity with sequences identified as AA291226 (zs47d03.r1 NCI_CGAP_GCB1 Homo sapiens cDNA clone 700613 5'), T05738 (EST03627 Homo sapiens cDNA clone HFBDF64), W51195 (ma14b04.r1 Life Tech mouse brain Mus musculus cDNA clone 304495 5'), and W93640 (zd95d09.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 357233 3'). The predicted amino acid sequence disclosed herein for
20 CC247_10 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CC247_10 protein demonstrated at least some similarity to sequences identified as M62424 (thrombin receptor [Homo sapiens]). The predicted CC247_10 protein is highly hydrophobic. Based upon sequence similarity, CC247_10 proteins and each similar protein or peptide may share at least some activity.

25

Clone "CI480_9"

A polynucleotide of the present invention has been identified as clone "CI480_9". CI480_9 was isolated from a human adult brain cDNA library using methods which are selective
30 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CI480_9 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CI480_9 protein").

The nucleotide sequence of CI480_9 as presently determined is reported in SEQ ID
35 NO:127, and includes a poly(A) tail. What applicants presently believe to be the proper reading

frame and the predicted amino acid sequence of the CI480_9 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:128. Amino acids 39 to 51 of SEQ ID NO:128 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 52. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CI480_9 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CI480_9 should be approximately 1940 bp.

The nucleotide sequence disclosed herein for CI480_9 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CI480_9 demonstrated at least some similarity with sequences identified as N99342 (IMAGE:20093 Homo sapiens cDNA clone 20093), R89725 (ym99d09.r1 Homo sapiens cDNA clone 167057 5'), and U60644 (Human HU-K4 mRNA, complete cds). The predicted amino acid sequence disclosed herein for CI480_9 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CI480_9 protein demonstrated at least some similarity to sequences identified as U60644 (HU-K4 [Homo sapiens]). Based upon sequence similarity, CI480_9 proteins and each similar protein or peptide may share at least some activity.

CI480_9 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 63 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "CO722_1"

A polynucleotide of the present invention has been identified as clone "CO722_1". CO722_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CO722_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CO722_1 protein").

The nucleotide sequence of CO722_1 as presently determined is reported in SEQ ID NO:129, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CO722_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:130. Amino acids 17 to 29 of SEQ ID NO:130 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal

sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CO722_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CO722_1 should be approximately 6800 bp.

5 The nucleotide sequence disclosed herein for CO722_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CO722_1 demonstrated at least some similarity with sequences identified as AA186616 (zp71a08.s1 Stratagene endothelial cell 937223 Homo sapiens cDNA clone 625622 3' similar to contains Alu repetitive element), H10376 (ym08a03.s1 Homo sapiens cDNA clone
10 47067 3'), N86013 (J5997F Fetal heart, Lambda ZAP Express Homo sapiens cDNA), U55258 (Human hBRAVO/Nr-CAM precursor (hBRAVO/ Nr-CAM) gene, complete cds), W19770 (zb39d01.r1 Soares parathyroid tumor NbHPA Homo sapiens), W31608 (zb91d09.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone), and X58482 (Chicken mRNA for neuronal transmembrane protein Nr-CAM, ng-CAM related). The predicted amino acid sequence
15 disclosed herein for CO722_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CO722_1 protein demonstrated at least some similarity to sequences identified as AB002341 (KIAA0343 [Homo sapiens]) and X58482 (Nr-CAM protein [Gallus gallus]). Based upon sequence similarity, CO722_1 proteins and each similar protein or peptide may share at least some activity. The
20 TopPredII computer program predicts two potential transmembrane domains within the CO722_1 protein sequence, centered around amino acids 610 and 1070 of SEQ ID NO:130, respectively.

CO722_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 160 kDa was detected in conditioned media fractions using SDS polyacrylamide gel electrophoresis.

25

Clone "CT748_2"

A polynucleotide of the present invention has been identified as clone "CT748_2". CT748_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
30 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CT748_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CT748_2 protein").

The nucleotide sequence of CT748_2 as presently determined is reported in SEQ ID NO:131, and includes a poly(A) tail. What applicants presently believe to be the proper reading
35 frame and the predicted amino acid sequence of the CT748_2 protein corresponding to the

foregoing nucleotide sequence is reported in SEQ ID NO:132. Amino acids 281 to 293 of SEQ ID NO:132 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 294. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CT748_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CT748_2 should be approximately 5500 bp.

The nucleotide sequence disclosed herein for CT748_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CT748_2 demonstrated at least some similarity with sequences identified as T48063 (yb24f03.s1 Homo sapiens cDNA clone 72125 3') and X54175 (Human specific Alu element (HS C4N2) DNA). The predicted amino acid sequence disclosed herein for CT748_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CT748_2 protein demonstrated at least some similarity to sequences identified as Z36714 (cyclin F [Homo sapiens]). Based upon sequence similarity, CT748_2 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of CT748_2 indicates that it may contain an Alu repetitive element.

Clone "AJ1_1"

A polynucleotide of the present invention has been identified as clone "AJ1_1". AJ1_1 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AJ1_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AJ1_1 protein").

The nucleotide sequence of the 5' portion of AJ1_1 as presently determined is reported in SEQ ID NO:133. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:134. The predicted amino acid sequence of the AJ1_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:134. Additional nucleotide sequence from the 3' portion of AJ1_1, including a poly(A) tail, is reported in SEQ ID NO:135.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AJ1_1 should be approximately 925 bp.

The predicted amino acid sequence disclosed herein for AJ1_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The

predicted AJ1_1 protein demonstrated at least some similarity to sequences identified as U39060 (GRIP1 [Mus musculus]). Based upon sequence similarity, AJ1_1 proteins and each similar protein or peptide may share at least some activity.

5 Clone "AQ73_3"

A polynucleotide of the present invention has been identified as clone "AQ73_3". AQ73_3 was isolated from a human adult ovary (PA-1 teratocarcinoma, untreated tissue pooled with retinoic-acid-treated and activin-treated tissue) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified
10 as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. AQ73_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AQ73_3 protein").

The nucleotide sequence of AQ73_3 as presently determined is reported in SEQ ID NO:136, and includes a poly(A) tail. What applicants presently believe to be the proper reading
15 frame and the predicted amino acid sequence of the AQ73_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:137.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AQ73_3 should be approximately 2800 bp.

The nucleotide sequence disclosed herein for AQ73_3 was searched against the GenBank
20 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. AQ73_3 demonstrated at least some similarity with sequences identified as AA514474 (nf57g01.s1 NCI_CGAP_Co3 Homo sapiens cDNA clone 924048), T47520 (Human hepatoma-derived growth factor (HDGF-2) cDNA), W24708 (zb62e08.r1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 308198 5'), and W45513 (zc27g08.s1 Soares senescent
25 fibroblasts NbHSF Homo sapiens cDNA clone 323582 3'). The predicted amino acid sequence disclosed herein for AQ73_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted AQ73_3 protein demonstrated at least some similarity to sequences identified as D16431 (hepatoma-derived GF [Homo sapiens]), D63707 (mouse hepatoma derived growth factor (HDGF) [Mus musculus]), R66727 (Human
30 hepatoma derived growth factor), U18997 (ORF_f299 [Escherichia coli]), U97193 (similar to S. cerevisiae SIR2 (SP P06700) and mouse hepatoma derived growth factor HDGF (NID g945418) [Caenorhabditis elegans]), and W09404 (Human hepatoma-derived growth factor (HDGF-2)). Based upon sequence similarity, AQ73_3 proteins and each similar protein or peptide may share at least some activity.

AQ73_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 67 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5 Clone "BG142_1"

A polynucleotide of the present invention has been identified as clone "BG142_1". BG142_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence
10 of the encoded protein. BG142_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BG142_1 protein").

The nucleotide sequence of BG142_1 as presently determined is reported in SEQ ID NO:138, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BG142_1 protein corresponding to the
15 foregoing nucleotide sequence is reported in SEQ ID NO:139.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG142_1 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for BG142_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
20 protocols. BG142_1 demonstrated at least some similarity with sequences identified as AA170261 (ms87h11.r1 Soares mouse 3NbMS Mus musculus cDNA clone 618597 5' similar to TR E245601 E245601 G-RICH BOX-BINDING PROTEIN), L04282 (Human CACCC box-binding protein mRNA, complete cds), N27696 (yx51h12.r1 Homo sapiens cDNA clone 265319 5'), W96110 (ze09a11.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone
25 358460 5'), and W96111 (ze09a11.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 358460 3'). The predicted amino acid sequence disclosed herein for BG142_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BG142_1 protein demonstrated at least some similarity to sequences identified as U80078 (transcription factor BFCOL1 [Mus musculus]) and X98096 (G-rich
30 box-binding protein [Mus musculus]). Based upon sequence similarity, BG142_1 proteins and each similar protein or peptide may share at least some activity.

Clone "BV66_1"

A polynucleotide of the present invention has been identified as clone "BV66_1".
35 BV66_1 was isolated from a human adult brain cDNA library using methods which are selective

for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV66_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV66_1 protein").

5 The nucleotide sequence of BV66_1 as presently determined is reported in SEQ ID NO:140, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BV66_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:141.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV66_1 should be approximately 870 bp.

 The nucleotide sequence disclosed herein for BV66_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database. The nucleotide sequence of BV66_1 indicates that it may contain a TAAA1 simple repeat element.

15

Clone "BV291_3"

 A polynucleotide of the present invention has been identified as clone "BV291_3". BV291_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
20 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BV291_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BV291_3 protein").

 The nucleotide sequence of BV291_3 as presently determined is reported in SEQ ID NO:142, and includes a poly(A) tail. What applicants presently believe to be the proper reading
25 frame and the predicted amino acid sequence of the BV291_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:143.

 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BV291_3 should be approximately 2000 bp.

 The nucleotide sequence disclosed herein for BV291_3 was searched against the GenBank
30 and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BV291_3 demonstrated at least some similarity with sequences identified as H10954 (ym06e09.r1 Homo sapiens cDNA clone 47034 5'), H10955 (ym06e09.s1 Homo sapiens cDNA clone 47034 3'), N25300 (yw52c10.s1 Homo sapiens cDNA clone 255858 3'), T25940 (Human gene signature HUMGS08173), T68890 (yc30g11.s1 Homo sapiens cDNA clone 82244 3'),
35 T78286 (yc99a08.r1 Homo sapiens cDNA clone 24033 5'), Z39987 (H. sapiens partial cDNA

sequence; clone c-1oh05), and Z47073 (*Caenorhabditis elegans* cosmid ZC506). The predicted amino acid sequence disclosed herein for BV291_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted BV291_3 protein demonstrated at least some similarity to sequences identified as X02155 (BTTGR_1 thyroglobulin [*Bos taurus*]). Based upon sequence similarity, BV291_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the BV291_3 protein sequence centered around amino acid 48 of SEQ ID NO:143.

BV291_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 30 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

Clone "CK201_1"

A polynucleotide of the present invention has been identified as clone "CK201_1". CK201_1 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CK201_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CK201_1 protein").

The nucleotide sequence of CK201_1 as presently determined is reported in SEQ ID NO:144, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CK201_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:145.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CK201_1 should be approximately 1080 bp.

The nucleotide sequence disclosed herein for CK201_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CK201_1 demonstrated at least some similarity with sequences identified as AA129133 (zo09h12.s1 Stratagene neuroepithelium NT2RAMI 937234 *Homo sapiens* cDNA clone 567239 3' similar to contains Alu repetitive element), D81444 (Human fetal brain cDNA 5'-end GEN-164G10), R36326 (yg69h09.r1 *Homo sapiens* cDNA clone 38821 5'), T08553 (Oncogene R-ras mutant cDNA (exons 2-6)), T31595 (Probe (BLUR13) for Alu repeat sequence), X03273 (Human Alu-family cluster 5' of alpha(1)-acid glycoprotein gene), and X69907 (*H.sapiens* gene for mitochondrial ATP synthase c subunit). The predicted amino acid sequence disclosed herein for CK201_1 was searched against the GenPept and GeneSeq amino acid

sequence databases using the BLASTX search protocol. The predicted CK201_1 protein demonstrated at least some similarity to sequences identified as D21827 (major surface glycoprotein [Pneumocystis carinii]). Based upon sequence similarity, CK201_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of
5 CK201_1 indicates that it may contain an Alu repetitive element.

CK201_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 40 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

10 Clone "CQ331_2"

A polynucleotide of the present invention has been identified as clone "CQ331_2". CQ331_2 was isolated from a human adult heart cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence
15 of the encoded protein. CQ331_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CQ331_2 protein").

The nucleotide sequence of CQ331_2 as presently determined is reported in SEQ ID NO:146, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CQ331_2 protein corresponding to the
20 foregoing nucleotide sequence is reported in SEQ ID NO:147. Amino acids 7 to 19 of SEQ ID NO:147 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CQ331_2 protein.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CQ331_2 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for CQ331_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CQ331_2 demonstrated at least some similarity with sequences identified as J03766
30 (Canine cardiac calsequestrin mRNA, complete cds), L29766 (Homo sapiens epoxide hydrolase (EPHX) gene, complete cds), N83601 (KK1173F Homo sapiens cDNA clone KK1173 5' similar to CALSEQUESTRIN (CARDIAC)), T99646 (ye73f12.s1 Homo sapiens cDNA clone 123407 3' similar to contains Alu repetitive element;contains PTR5 repetitive element), and W76326 (zd60d04.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 345031 5' similar to
35 contains Alu repetitive element). The predicted amino acid sequence disclosed herein for

CQ331_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CQ331_2 protein demonstrated at least some similarity to sequences identified as J03766 (DOGCAL_1 Canine cardiac calsequestrin mRNA, complete cds [Canis canis]) and X55040 (calsequestrin [Oryctolagus cuniculus]). Based upon
5 sequence similarity, CQ331_2 proteins and each similar protein or peptide may share at least some activity.

Clone "CT550_1"

A polynucleotide of the present invention has been identified as clone "CT550_1".
10 CT550_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CT550_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CT550_1 protein").

15 The nucleotide sequence of CT550_1 as presently determined is reported in SEQ ID NO:148, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CT550_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:149.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
20 CT550_1 should be approximately 1070 bp.

The nucleotide sequence disclosed herein for CT550_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database. The TopPredII computer program predicts a potential transmembrane domain within the CT550_1 protein sequence centered around amino
25 acid 25 of SEQ ID NO:149.

CT550_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 7 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

30 Clone "CT585_1"

A polynucleotide of the present invention has been identified as clone "CT585_1". CT585_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence

of the encoded protein. CT585_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CT585_1 protein").

The nucleotide sequence of CT585_1 as presently determined is reported in SEQ ID NO:150, and includes a poly(A) tail. What applicants presently believe to be the proper reading
5 frame and the predicted amino acid sequence of the CT585_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:151. Amino acids 2 to 14 of SEQ ID NO:151 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal
10 sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CT585_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CT585_1 should be approximately 2710 bp.

The nucleotide sequence disclosed herein for CT585_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
15 protocols. CT585_1 demonstrated at least some similarity with sequences identified as AA069442 (zf74b02.s1 Soares pineal gland N3HPG Homo sapiens cDNA clone 382635 3'), L38961 (Homo sapiens putative transmembrane protein (B5) mRNA, complete cds), N34932 (yy49b10.s1 Homo sapiens cDNA clone 276859 3'), N60101 (TgESTzy11f10.r1 Toxoplasma gondii cDNA clone tgzy11f10.r1 5'), and U13019 (Caenorhabditis elegans cosmid T12A2). The predicted amino acid
20 sequence disclosed herein for CT585_1 was searched against the GenPept, GeneSeq, and SwissProt amino acid sequence databases using the BLASTX search protocol. The predicted CT585_1 protein demonstrated at least some similarity to sequences identified as L34260 (transmembrane protein [Mus musculus]), L38961 (transmembrane protein [Homo sapiens]), P46975 (Caenorhabditis elegans oligosaccharyl transferase stt3 [Caenorhabditis elegans]), and
25 U13019 (Caenorhabditis elegans cosmid T12A2 [Caenorhabditis elegans]). Based upon sequence similarity, CT585_1 proteins and each similar protein or peptide may share at least some activity.

Clone "CT797_3"

A polynucleotide of the present invention has been identified as clone "CT797_3".
30 CT797_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CT797_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CT797_3 protein").

The nucleotide sequence of CT797_3 as presently determined is reported in SEQ ID NO:152, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CT797_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:153.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CT797_3 should be approximately 3300 bp.

 The nucleotide sequence disclosed herein for CT797_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CT797_3 demonstrated at least some similarity with sequences identified as AA573847
10 (nk08d06.s1 NCI_CGAP_Co2 Homo sapiens cDNA clone IMAGE:1012907). The predicted amino acid sequence disclosed herein for CT797_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted CT797_3 protein demonstrated at least some similarity to sequences identified as U18309 (chromokinesin [Gallus gallus]) and Z82271 (T01G1.1 [Caenorhabditis elegans]). Based upon
15 sequence similarity, CT797_3 proteins and each similar protein or peptide may share at least some activity.

 CT797_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 80 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

20

Clone "CB107_1"

 A polynucleotide of the present invention has been identified as clone "CB107_1". CB107_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
25 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CB107_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CB107_1 protein").

 The nucleotide sequence of the 5' portion of CB107_1 as presently determined is reported in SEQ ID NO:154. An additional internal nucleotide sequence from CB107_1 as presently
30 determined is reported in SEQ ID NO:155. What applicants believe is the proper reading frame and the predicted amino acid sequence encoded by such internal sequence is reported in SEQ ID NO:156. Additional nucleotide sequence from the 3' portion of CB107_1, including a poly(A) tail, is reported in SEQ ID NO:157.

 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
35 CB107_1 should be approximately 3300 bp.

The nucleotide sequence disclosed herein for CB107_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CB107_1 demonstrated at least some similarity with sequences identified as AA121485 (zn80a02.s1 Stratagene lung carcinoma 937218 Homo sapiens cDNA clone 564458 3'),
5 AA428192 (zw51b08.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773559 3'),
D83018 (Human mRNA for nel-related protein 2, complete cds), F10919 (H. sapiens partial
cDNA sequence; clone c-3lg01), H15375 (ym28d09.r1 Homo sapiens cDNA clone 49527 5'
similar to SP A54105 A54105 FIBRILLIN-2 PRECURSOR), U48245 (Rattus norvegicus protein
kinase C-binding protein Nel mRNA, complete cds), U59230 (Mus musculus mel (MEL91)
10 mRNA, complete cds), and W28387 (46c5 Human retina cDNA randomly primed sublibrary
Homo sapiens cDNA). The predicted amino acid sequence disclosed herein for CB107_1 was
searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX
search protocol. The predicted CB107_1 protein demonstrated at least some similarity to
sequences identified as D83018 (nel-related protein 2 [Homo sapiens]), R05222 (Antigen
15 GX5401FL encoded by Eimeria tenella genomic DNA), R79964 (Connective tissue growth
factor), U48245 (RNU48245_1 protein kinase C-binding protein Nel [Rattus norvegicus]), and
U59230 (mel [Mus musculus]). Based upon sequence similarity, CB107_1 proteins and each
similar protein or peptide may share at least some activity.

20

Clone "CG300_3"

A polynucleotide of the present invention has been identified as clone "CG300_3".
CG300_3 was isolated from a human adult testes cDNA library using methods which are selective
for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
25 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence
of the encoded protein. CG300_3 is a full-length clone, including the entire coding sequence of
a secreted protein (also referred to herein as "CG300_3 protein").

The nucleotide sequence of CG300_3 as presently determined is reported in SEQ ID
NO:158, and includes a poly(A) tail. What applicants presently believe to be the proper reading
30 frame and the predicted amino acid sequence of the CG300_3 protein corresponding to the
foregoing nucleotide sequence is reported in SEQ ID NO:159. Amino acids 30 to 42 of SEQ ID
NO:159 are a predicted leader/signal sequence, with the predicted mature amino acid sequence
beginning at amino acid 43. Due to the hydrophobic nature of the predicted leader/signal
sequence, it is likely to act as a transmembrane domain should the predicted leader/signal
35 sequence not be separated from the remainder of the CG300_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CG300_3 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for CG300_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CG300_3 demonstrated at least some similarity with sequences identified as N40185 (yy44d08.s1 Homo sapiens cDNA clone 276399 3') and W01791 (za72d06.r1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 298091 5'). Based upon sequence similarity, CG300_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four potential transmembrane domains within the CG300_3 protein sequence, centered around amino acids 34, 98, 151, and 179 of SEQ ID NO:159, respectively.

CG300_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 29 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "CJ145_1"

A polynucleotide of the present invention has been identified as clone "CJ145_1". CJ145_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CJ145_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CJ145_1 protein").

The nucleotide sequence of CJ145_1 as presently determined is reported in SEQ ID NO:160, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CJ145_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:161. Amino acids 6 to 18 of SEQ ID NO:161 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 19. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CJ145_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CJ145_1 should be approximately 3600 bp.

The nucleotide sequence disclosed herein for CJ145_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search

protocols. CJ145_1 demonstrated at least some similarity with sequences identified as R43655 (yc86b04.s1 Homo sapiens cDNA clone 22829 3'), R50995 (yg63f06.s1 Homo sapiens cDNA clone 37377 3' similar to contains MER22 repetitive element), and W92748 (zd92h03.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 356981 3'). Based upon sequence similarity,

5 CJ145_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of CJ145_1 indicates that it may contain a CA simple repeat element.

Clone "CJ160_11"

A polynucleotide of the present invention has been identified as clone "CJ160_11".

10 CJ160_11 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CJ160_11 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CJ160_11 protein").

15 The nucleotide sequence of CJ160_11 as presently determined is reported in SEQ ID NO:162, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CJ160_11 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:163. Amino acids 17 to 29 of SEQ ID NO:163 are a predicted leader/signal sequence, with the predicted mature amino acid sequence

20 beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CJ160_11 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CJ160_11 should be approximately 1700 bp.

25 The nucleotide sequence disclosed herein for CJ160_11 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CJ160_11 demonstrated at least some similarity with sequences identified as AA024511 (ze76e04.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 364926 3') and AC000074 (00884; HTGS phase 3, complete sequence). Based upon sequence similarity,

30 CJ160_11 proteins and each similar protein or peptide may share at least some activity.

CJ160_11 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 96 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

35 Clone "CO20_1"

A polynucleotide of the present invention has been identified as clone "CO20_1". CO20_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CO20_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CO20_1 protein").

The nucleotide sequence of the 5' portion of CO20_1 as presently determined is reported in SEQ ID NO:164. What applicants presently believe is the proper reading frame for the coding region is indicated in SEQ ID NO:165. The predicted amino acid sequence of the CO20_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:165. Amino acids 17 to 29 of SEQ ID NO:165 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CO20_1 protein. Additional nucleotide sequence from the 3' portion of CO20_1, including a poly(A) tail, is reported in SEQ ID NO:166.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CO20_1 should be approximately 2400 bp.

The nucleotide sequence disclosed herein for CO20_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CO20_1 demonstrated at least some similarity with sequences identified as AA045770 (zl68b10.s1 Stratagene colon (#937204) Homo sapiens cDNA clone 509755 3' similar to SW:R13A_HUMAN P40429 60S RIBOSOMAL PROTEIN L13A), AA070899 (zm66c01.s1 Stratagene neuroepithelium (#937231) Homo sapiens cDNA clone 530592 3' similar to contains Alu repetitive element), AA325205 (EST28155 Cerebellum II Homo sapiens cDNA 5' end), N22253 (yw36a08.s1 Homo sapiens cDNA clone 254294 3' similar to SP S29539 S29539 BASIC PROTEIN, 23K), R01933 (ye85g07.s1 Homo sapiens cDNA clone 124572 3' similar to SP:S29539 S29539 BASIC PROTEIN, 23K), R12008 (yf51f04.r1 Homo sapiens cDNA clone 25456 5'), R39848 (yf51f04.s1 Homo sapiens cDNA clone 25456 3' similar to contains Alu repetitive element;contains PTR5 repetitive element), R56565 (yg91c12.r1 Homo sapiens cDNA clone 40891 5'), T19487 (Human gene signature HUMGS00543), T30988 (EST25695 Homo sapiens cDNA 5' end similar to None), U37026 (Rattus norvegicus brain sodium channel beta 2 subunit (SCNB2) mRNA, complete cds), and X56932 (H.sapiens mRNA for 23 kD highly basic protein). The predicted amino acid sequence disclosed herein for CO20_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol.

The predicted CO20_1 protein demonstrated at least some similarity to sequences identified as U37026 (sodium channel beta 2 subunit [Rattus norvegicus]), U58658 (unknown [Homo sapiens]), and X56932 (23 kD highly basic protein [Homo sapiens]). The sodium channel beta 2 subunit is a glycoprotein with an extracellular domain containing an immunoglobulin-like fold with
5 similarity to the neural cell adhesion molecule contactin. Based upon sequence similarity, CO20_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of CO20_1 indicates that it may contain an Alu repetitive element.

Clone "CO223_3"

10 A polynucleotide of the present invention has been identified as clone "CO223_3". CO223_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CO223_3 is a full-length clone, including the entire coding sequence of
15 a secreted protein (also referred to herein as "CO223_3 protein").

The nucleotide sequence of CO223_3 as presently determined is reported in SEQ ID NO:167, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CO223_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:168. Amino acids 35 to 47 of SEQ ID
20 NO:168 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 48. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CO223_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone
25 CO223_3 should be approximately 700 bp.

The nucleotide sequence disclosed herein for CO223_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CO223_3 demonstrated at least some similarity with sequences identified as AA004498 (zh87b06.r1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 428243
30 5' similar to gb M62505 C5A ANAPHYLATOXIN CHEMOTACTIC RECEPTOR (HUMAN);contains L1.t1 L1 repetitive element) and U47924 (Human chromosome 12p13 gene cluster, surface antigen CD4 (CD4), A, B, G-protein beta-3 subunit (GNB3), isopeptidase T (ISOT) and triosephosphate isomerase (TPI) genes, complete cds). Based upon sequence similarity, CO223_3 proteins and each similar protein or peptide may share at least some activity.

The 3' end of the CO223_3 polynucleotide sequence contains a 54-bp sequence that is repeated three times in the clone; these repeats begin at positions 314, 368, and 422 of SEQ ID NO:167 and encode amino acids 47 to 64, 65 to 82, and 83 to 99 of SEQ ID NO:168, respectively.

5 Clone "CO310_2"

A polynucleotide of the present invention has been identified as clone "CO310_2". CO310_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CO310_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CO310_2 protein").

The nucleotide sequence of CO310_2 as presently determined is reported in SEQ ID NO:169, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CO310_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:170.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CO310_2 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for CO310_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database. The nucleotide sequence of CO310_2 indicates that it may contain an L1 repetitive element.

Clone "CP258_3"

A polynucleotide of the present invention has been identified as clone "CP258_3". CP258_3 was isolated from a human adult salivary gland cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CP258_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CP258_3 protein").

The nucleotide sequence of CP258_3 as presently determined is reported in SEQ ID NO:171, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CP258_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:172. Amino acids 3 to 15 of SEQ ID NO:172 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal

sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CP258_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CP258_3 should be approximately 560 bp.

5 The nucleotide sequence disclosed herein for CP258_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

CP258_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 26 kDa was detected in conditioned medium and membrane fractions using
10 SDS polyacrylamide gel electrophoresis.

Clone "CW1155_3"

A polynucleotide of the present invention has been identified as clone "CW1155_3". CW1155_3 was isolated from a human fetal brain cDNA library using methods which are
15 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CW1155_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CW1155_3 protein").

The nucleotide sequence of CW1155_3 as presently determined is reported in SEQ ID
20 NO:173, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CW1155_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:174. Amino acids 220 to 232 of SEQ ID NO:174 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 233. Due to the hydrophobic nature of the predicted leader/signal
25 sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the CW1155_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CW1155_3 should be approximately 1170 bp.

The nucleotide sequence disclosed herein for CW1155_3 was searched against the
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CW1155_3 demonstrated at least some similarity with sequences identified as AA169043 (ms36h08.r1 Stratagene mouse heart (#937316) Mus musculus cDNA clone 613695 5'), D86145 (Rat mRNA), and H29261 (ym32b03.s1 Homo sapiens cDNA clone 49733 3'). Based upon sequence similarity, CW1155_3 proteins and each similar protein or peptide may share at
35 least some activity.

Clone "CZ247_2"

A polynucleotide of the present invention has been identified as clone "CZ247_2". CZ247_2 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
5 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CZ247_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CZ247_2 protein").

The nucleotide sequence of CZ247_2 as presently determined is reported in SEQ ID NO:175, and includes a poly(A) tail. What applicants presently believe to be the proper reading
10 frame and the predicted amino acid sequence of the CZ247_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:176. Amino acids 545 to 557 of SEQ ID NO:176 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 558. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal
15 sequence not be separated from the remainder of the CZ247_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CZ247_2 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for CZ247_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
20 protocols. CZ247_2 demonstrated at least some similarity with sequences identified as T09256 (Human ara Kb beta-galactosidase fusion protein coding sequence), W27222 (26h9 Human retina cDNA randomly primed sublibrary Homo sapiens cDNA), and W72736 (zd71e02.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 346106 3'). The predicted amino acid sequence disclosed herein for CZ247_2 was searched against the GenPept and GeneSeq amino acid
25 sequence databases using the BLASTX search protocol. The predicted CZ247_2 protein demonstrated at least some similarity to sequences identified as R88069 (Human ara Kb beta-galactosidase fusion protein). Based upon sequence similarity, CZ247_2 proteins and each similar protein or peptide may share at least some activity.

30 Clone "AM666_1"

A polynucleotide of the present invention has been identified as clone "AM666_1". AM666_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino

acid sequence of the encoded protein. AM666_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "AM666_1 protein").

The nucleotide sequence of AM666_1 as presently determined is reported in SEQ ID NO:177, and includes a poly(A) tail. What applicants presently believe to be the proper reading
5 frame and the predicted amino acid sequence of the AM666_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:178. Amino acids 15 to 27 of SEQ ID NO:178 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the predicted leader/signal
10 sequence not be separated from the remainder of the AM666_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM666_1 should be approximately 1300 bp.

The nucleotide sequence disclosed herein for AM666_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA
15 search protocols. AM666_1 demonstrated at least some similarity with sequences identified as AA493985 (nh07g08.s1 NCI_CGAP_Thy1 Homo sapiens cDNA clone). Based upon sequence similarity, AM666_1 proteins and each similar protein or peptide may share at least some activity.

AM666_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 17 kDa was detected in membrane fractions using SDS polyacrylamide gel
20 electrophoresis.

Clone "BN387_3"

A polynucleotide of the present invention has been identified as clone "BN387_3". BN387_3 was isolated from a human adult placenta cDNA library using methods which are
25 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BN387_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "BN387_3 protein").

The nucleotide sequence of BN387_3 as presently determined is reported in SEQ ID
30 NO:179, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BN387_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:180. Amino acids 14 to 26 of SEQ ID NO:180 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal

sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BN387_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BN387_3 should be approximately 2000 bp.

5 The nucleotide sequence disclosed herein for BN387_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. BN387_3 demonstrated at least some similarity with sequences identified as H16912 (ym39d01.r1 Homo sapiens cDNA clone 50771 5'). Based upon sequence similarity, BN387_3 proteins and each similar protein or peptide may share at least some activity.

10 BN387_3 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 30 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

Clone "BQ135_2"

15 A polynucleotide of the present invention has been identified as clone "BQ135_2". BQ135_2 was isolated from a human adult colon (adenocarcinoma Caco2) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. BQ135_2 is a full-length clone,
20 including the entire coding sequence of a secreted protein (also referred to herein as "BQ135_2 protein").

 The nucleotide sequence of BQ135_2 as presently determined is reported in SEQ ID NO:181, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the BQ135_2 protein corresponding to the
25 foregoing nucleotide sequence is reported in SEQ ID NO:182.

 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BQ135_2 should be approximately 1200 bp.

 The nucleotide sequence disclosed herein for BQ135_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search
30 protocols. BQ135_2 demonstrated at least some similarity with sequences identified as AA023751 (mh81f01.r1 Soares mouse placenta 4NbMP13.5 14.5 Mus musculus cDNA clone 457369 5'), AA105433 (ml83g01.r1 Stratagene mouse kidney (#937315) Mus musculus cDNA clone 518640 5'), D64061 (Rat brain mRNA for annexin V-binding protein (ABP-7), partial cds), and N67257 (yz49b08.s1 Homo sapiens cDNA clone 286359 3'). The predicted amino acid
35 sequence disclosed herein for BQ135_2 was searched against the GenPept and GeneSeq amino

acid sequence databases using the BLASTX search protocol. The predicted BQ135_2 protein demonstrated at least some similarity to sequences identified as D64061 (annexin V-binding protein (ABP-7) [Rattus norvegicus]). Annexins associate with membranes and act as ion channels, they can also act as an autocrine factor that enhances osteoclast formation and bone resorption. Annexins have been localized in nucleoli and mitochondria but also in the cytoplasm, plasma (i.e. blood) and in association with vesicles. They are probably involved in fusing vesicles to each other and to plasma membranes causing secretion of vesicular contents. Specifically they have a calcium-dependent ability to bind phospholipids. Thus they are membrane associated. It is possible that annexin-binding proteins are also membrane associated even though they are highly hydrophilic through the same mechanism (electrostatic interaction with phospholipids of membranes). Based upon sequence similarity, BQ135_2 proteins and each similar protein or peptide may share at least some activity.

Clone "CR678_1"

A polynucleotide of the present invention has been identified as clone "CR678_1". CR678_1 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CR678_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CR678_1 protein").

The nucleotide sequence of CR678_1 as presently determined is reported in SEQ ID NO:183, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CR678_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:184.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CR678_1 should be approximately 870 bp.

The nucleotide sequence disclosed herein for CR678_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CR678_1 demonstrated at least some similarity with sequences identified as X85232 (H.sapiens chromosome 3 sequences). Based upon sequence similarity, CR678_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of CR678_1 indicates that it may contain an Alu repetitive element.

Clone "CW420_2"

A polynucleotide of the present invention has been identified as clone "CW420_2". CW420_2 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CW420_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CW420_2 protein").

The nucleotide sequence of CW420_2 as presently determined is reported in SEQ ID NO:185, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CW420_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:186.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CW420_2 should be approximately 5100 bp.

The nucleotide sequence disclosed herein for CW420_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. CW420_2 demonstrated at least some similarity with sequences identified as T55440 (yb38e09.s1 Homo sapiens cDNA clone 73480 3'). Based upon sequence similarity, CW420_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the CW420_2 protein sequence centered around amino acids 500 and 1270 of SEQ ID NO:186, respectively.

Clone "CW795_2"

A polynucleotide of the present invention has been identified as clone "CW795_2". CW795_2 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CW795_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CW795_2 protein").

The nucleotide sequence of CW795_2 as presently determined is reported in SEQ ID NO:187, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CW795_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:188.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CW795_2 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for CW795_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA

search protocols. CW795_2 demonstrated at least some similarity with sequences identified as AA115676 (zl86a09.s1 Stratagene colon (#937204) Homo sapiens cDNA clone 511480 3'), N22955 (yw44h07.s1 Homo sapiens cDNA clone 255133 3'), and W56804 (zd16g06.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 340858 3'). The predicted amino acid sequence disclosed herein for CW795_2 was searched against the GenPept, GeneSeq, and SwissProt amino acid sequence databases using the BLASTX search protocol. The predicted CW795_2 protein demonstrated at least some similarity to sequences identified as X81068 (probable mitochondrial protein) and the yeast proteins rca1 and afg3 (tat-binding homologues). Based upon sequence similarity, CW795_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the CW795_2 protein sequence centered around amino acids 60 and 170 of SEQ ID NO:188, respectively.

CW795_2 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 10 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "CW823_3"

A polynucleotide of the present invention has been identified as clone "CW823_3". CW823_3 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. CW823_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "CW823_3 protein").

The nucleotide sequence of CW823_3 as presently determined is reported in SEQ ID NO:189, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the CW823_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:190.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone CW823_3 should be approximately 600 bp.

The nucleotide sequence disclosed herein for CW823_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

Clone "DF989_3"

A polynucleotide of the present invention has been identified as clone "DF989_3". DF989_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding
5 a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. DF989_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "DF989_3 protein").

The nucleotide sequence of the 5' portion of DF989_3 as presently determined is reported in SEQ ID NO:191. What applicants presently believe is the proper reading frame for the coding
10 region is indicated in SEQ ID NO:192. The predicted amino acid sequence of the DF989_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:192. Amino acids 2 to 14 of SEQ ID NO:192 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the
15 predicted leader/signal sequence not be separated from the remainder of the DF989_3 protein. Additional nucleotide sequence from the 3' portion of DF989_3, including a poly(A) tail, is reported in SEQ ID NO:193.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone DF989_3 should be approximately 1800 bp.

20 The nucleotide sequence disclosed herein for DF989_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. DF989_3 demonstrated at least some similarity with sequences identified as R24724 (yg43c05.r1 Homo sapiens cDNA clone 35337 5') and T33717 (EST58870 Homo sapiens cDNA 5' end similar to None). Based upon sequence similarity, DF989_3 proteins and each similar
25 protein or peptide may share at least some activity.

Clone "DL162_1"

A polynucleotide of the present invention has been identified as clone "DL162_1". DL162_1 was isolated from a human adult brain cDNA library using methods which are selective
30 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. DL162_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "DL162_1 protein").

The nucleotide sequence of DL162_1 as presently determined is reported in SEQ ID
35 NO:194, and includes a poly(A) tail. What applicants presently believe to be the proper reading

frame and the predicted amino acid sequence of the DL162_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:195. Amino acids 28 to 40 of SEQ ID NO:195 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 41. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the DL162_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone DL162_1 should be approximately 875 bp.

The nucleotide sequence disclosed herein for DL162_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. No hits were found in the database.

Clone "DL162_2"

A polynucleotide of the present invention has been identified as clone "DL162_2". DL162_2 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. DL162_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "DL162_2 protein").

The nucleotide sequence of DL162_2 as presently determined is reported in SEQ ID NO:196, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the DL162_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:197. Amino acids 1 to 13 of SEQ ID NO:197 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 14. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the DL162_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone DL162_2 should be approximately 4000 bp.

The predicted amino acid sequence disclosed herein for DL162_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted DL162_2 protein demonstrated at least some similarity to sequences identified as AB002309 (KIAA0311 protein [Homo sapiens]). The TopPredII computer program predicts a potential transmembrane domains within the DL162_2 protein sequence near the carboxyl terminus of SEQ ID NO:197.

DL162_2 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 160 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5 Clone "EC172_1"

A polynucleotide of the present invention has been identified as clone "EC172_1". EC172_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. EC172_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "EC172_1 protein").

10 The nucleotide sequence of EC172_1 as presently determined is reported in SEQ ID NO:198, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the EC172_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:199. Amino acids 659 to 671 of SEQ ID NO:199 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 672. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the EC172_1 protein.

15 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone EC172_1 should be approximately 4000 bp.

20 The nucleotide sequence disclosed herein for EC172_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. EC172_1 demonstrated at least some similarity with sequences identified as H31192 (EST104991 Rattus sp. cDNA 3' end similar to C.elegans hypothetical protein ZK1098.10) and U29585 (Streptococcus pyogenes emm18.1). The predicted amino acid sequence disclosed herein for EC172_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted EC172_1 protein demonstrated at least some similarity to sequences identified as Z22176 (ZK1098.10 [Caenorhabditis elegans]). Based upon sequence similarity, EC172_1 proteins and each similar protein or peptide may share at least some activity.

25 Deposit of Clones

30 Clones AX65_22, BD335_14, BG241_1, BL187_4, BL249_18, BO71_1, BO365_2, BV51_1, BV140_3, BV141_2, CC194_4, and DA136_11 were deposited on October 3, 1996 with

the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98196, from which each clone comprising a particular polynucleotide is obtainable.

5 Clones AR415_4, AS63_29, BG160_1, BO432_4, BO538_2, BR595_4, CI490_2, CI522_1, CN238_1, CO390_1, and AY304_1 (an additional isolate of clone AY304_14) were deposited on October 25, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98232, from which each clone comprising
10 a particular polynucleotide is obtainable. Clone AY304_14 was deposited on October 23, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 98561.

Clones AJ20_2, AR440_1, AS164_1, AX8_1, BD176_3, BD339_1, BD427_1, BL229_22,
15 BV123_16, and CH377_1 were deposited on November 15, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98261, from which each clone comprising a particular polynucleotide is obtainable.

Clones BD441_1, BD441_2, BG102_3, BK158_1, BP163_1, BZ16_3, CC182_1,
20 CG109_1 and CJ397_1 were deposited on November 20, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98264, from which each clone comprising a particular polynucleotide is obtainable.

Clones AM795_4, AT340_1, BG132_1, BG219_2, BG366_2, BV172_2, CC247_10,
25 CI480_9, CO722_1, and CT748_2 were deposited on December 5, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98271, from which each clone comprising a particular polynucleotide is obtainable.

Clones AJ1_1, AQ73_3, BG142_1, BV66_1, BV291_3, CK201_1, CQ331_2, CT550_1,
30 CT585_1 and CT797_3 were deposited on December 13, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98278, from which each clone comprising a particular polynucleotide is obtainable.

Clones CB107_1, CG300_3, CJ145_1, CJ160_11, CO20_1, CO223_1, CO310_2,
35 CP258_3, CW1155_3 and CZ247_2 were deposited on December 17, 1996 with the ATCC

(American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98279, from which each clone comprising a particular polynucleotide is obtainable. Clone CO223_3 was deposited on January 9, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 98291.

Clones AM666_1, BN387_3, BQ135_2, CR678_1, CW420_2, CW795_2, CW823_3, DF989_3, DL162_2, DL162_1, and EC172_1 were deposited on January 10, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98292, from which each clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R. § 1.806.

Each clone has been transfected into separate bacterial cells (*E. coli*) in this composite deposit. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or pNOTs vector depicted in Figures 1A and 1B, respectively. The pED6dpc2 vector ("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and insertion of the M13 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences provided herein, or from a combination of those sequences. The sequence of an oligonucleotide probe that was used to isolate or to sequence each full-length clone is identified below, and should be most reliable in isolating the clone of interest.

	<u>Clone</u>	<u>Probe Sequence</u>
	AX65_22	SEQ ID NO:200
	BD335_14	SEQ ID NO:201
	BG241_1	SEQ ID NO:202
5	BL187_4	SEQ ID NO:203
	BL249_18	SEQ ID NO:204
	BO71_1	SEQ ID NO:205
	BO365_2	SEQ ID NO:206
	BV51_1	SEQ ID NO:207
10	BV140_3	SEQ ID NO:208
	BV141_2	SEQ ID NO:209
	CC194_4	SEQ ID NO:210
	DA136_11	SEQ ID NO:211
	AR415_4	SEQ ID NO:212
15	AS63_29	SEQ ID NO:213
	AY304_14	SEQ ID NO:214
	BG160_1	SEQ ID NO:215
	BO432_4	SEQ ID NO:216
	BO538_2	SEQ ID NO:217
20	BR595_4	SEQ ID NO:218
	CI490_2	SEQ ID NO:219
	CI522_1	SEQ ID NO:220
	CN238_1	SEQ ID NO:221
	CO390_1	SEQ ID NO:222
25	AJ20_2	SEQ ID NO:223
	AR440_1	SEQ ID NO:224
	AS164_1	SEQ ID NO:225
	AX8_1	SEQ ID NO:226
	BD176_3	SEQ ID NO:227
30	BD339_1	SEQ ID NO:228
	BD427_1	SEQ ID NO:229
	BL229_22	SEQ ID NO:230
	BV123_16	SEQ ID NO:231
	CH377_1	SEQ ID NO:232
35	BD441_1	SEQ ID NO:233

	BD441_2	SEQ ID NO:234
	BG102_3	SEQ ID NO:235
	BK158_1	SEQ ID NO:236
	BP163_1	SEQ ID NO:237
5	BZ16_3	SEQ ID NO:238
	CC182_1	SEQ ID NO:239
	CG109_1	SEQ ID NO:240
	CJ397_1	SEQ ID NO:241
	AM795_4	SEQ ID NO:242
10	AT340_1	SEQ ID NO:243
	BG132_1	SEQ ID NO:244
	BG219_2	SEQ ID NO:245
	BG366_2	SEQ ID NO:246
	BV172_2	SEQ ID NO:247
15	CC247_10	SEQ ID NO:248
	CI480_9	SEQ ID NO:249
	CO722_1	SEQ ID NO:250
	CT748_2	SEQ ID NO:251
	AJ1_1	SEQ ID NO:252
20	AQ73_3	SEQ ID NO:253
	BG142_1	SEQ ID NO:254
	BV66_1	SEQ ID NO:255
	BV291_3	SEQ ID NO:256
	CK201_1	SEQ ID NO:257
25	CQ331_2	SEQ ID NO:258
	CT550_1	SEQ ID NO:259
	CT585_1	SEQ ID NO:260, SEQ ID NO:262
	CT797_3	SEQ ID NO:261
	CB107_1	SEQ ID NO:263
30	CG300_3	SEQ ID NO:264
	CJ145_1	SEQ ID NO:265
	CJ160_11	SEQ ID NO:266
	CO20_1	SEQ ID NO:267
	CO223_3	SEQ ID NO:268
35	CO310_2	SEQ ID NO:269

	CP258_3	SEQ ID NO:270
	CW1155_3	SEQ ID NO:271
	CZ247_2	SEQ ID NO:272
	AM666_1	SEQ ID NO:273
5	BN387_3	SEQ ID NO:274
	BQ135_2	SEQ ID NO:275
	CR678_1	SEQ ID NO:276
	CW420_2	SEQ ID NO:277
	CW795_2	SEQ ID NO:278
10	CW823_3	SEQ ID NO:279
	DF989_3	SEQ ID NO:280
	DL162_1, DL162_2	SEQ ID NO:281
	EC172_1	SEQ ID NO:282

15 In the sequences listed above which include an N at position 2, that position is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

20 The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;
- (b) It should be designed to have a T_m of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).

25 The oligonucleotide should preferably be labeled with γ -³²P ATP (specific activity 6000 Ci/mmole) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a
30 scintillation counter. Preferably, specific activity of the resulting probe should be approximately 4e+6 dpm/pmole.

The bacterial culture containing the pool of full-length clones should preferably be thawed and 100 μ l of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100 μ g/ml. The culture should preferably be grown to saturation at 37°C,
35 and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions

should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100 µg/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at 37°C. Other known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100 µg/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at a concentration greater than or equal to 1e+6 dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein

may be obtained by expression of the disclosed full-length polynucleotide (preferably those deposited with the ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

5 The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3'
10 untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene"
15 is a gene that has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

 The chromosomal location corresponding to the polynucleotide sequences disclosed herein may also be determined, for example by hybridizing appropriately labeled polynucleotides of the present invention to chromosomes *in situ*. It may also be possible to determine the
20 corresponding chromosomal location for a disclosed polynucleotide by identifying significantly similar nucleotide sequences in public databases, such as expressed sequence tags (ESTs), that have already been mapped to particular chromosomal locations. For at least some of the polynucleotide sequences disclosed herein, public database sequences having at least some similarity to the polynucleotide of the present invention have been listed by database accession
25 number. Searches using the GenBank accession numbers of these public database sequences can then be performed at an Internet site provided by the National Center for Biotechnology Information having the address <http://www.ncbi.nlm.nih.gov/UniGene/>, in order to identify "UniGene clusters" of overlapping sequences. Many of the "UniGene clusters" so identified will already have been mapped to particular chromosomal sites.

30 Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, *Trends Pharmacol. Sci.* 15(7): 250-254; Lavarosky *et al.*, 1997, *Biochem. Mol. Med.* 62(1): 11-22; and
35 Hampel, 1998, *Prog. Nucleic Acid Res. Mol. Biol.* 58: 1-39; all of which are incorporated by

reference herein). The desired change in gene expression can also be achieved through the use of double-stranded ribonucleotide molecules having some complementarity to the mRNA transcribed from the gene, and which interfere with the transcription, stability, or expression of the mRNA ("RNA interference" or "RNAi"; Fire *et al.*, 1998, *Nature* 391 (6669): 806-811; 5 Montgomery *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95 (26): 15502-15507; and Sharp, 1999, *Genes Dev.* 13 (2): 139-141; all of which are incorporated by reference herein). Transgenic animals that have multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic 10 animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the 15 corresponding gene(s) or through deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, *Bioessays* 14(9): 629-633; Zwaal *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* 90(16): 7431-7435; Clark *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91(2): 719-722; all of which are incorporated by reference herein), or through 20 homologous recombination, preferably detected by positive/negative genetic selection strategies (Mansour *et al.*, 1988, *Nature* 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models 25 for the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part or all of the 30 intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of transmembrane domains in an amino acid sequence, domains

which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* 266: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of Molecular Biology* 215: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* 3: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins

and BLASTP, and Q=10 for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

Species homologues of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuánez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity
5 (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic
10 acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides that hybridize under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent
15 conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) [‡]	Hybridization Temperature and Buffer [†]	Wash Temperature and Buffer [†]
5	A	≥ 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
	B	<50	T _B *; 1xSSC	T _B *; 1xSSC
	C	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
	D	<50	T _D *; 1xSSC	T _D *; 1xSSC
	E	≥ 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
	F	<50	T _F *; 1xSSC	T _F *; 1xSSC
10	G	≥ 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
	H	<50	T _H *; 4xSSC	T _H *; 4xSSC
	I	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
	J	<50	T _J *; 4xSSC	T _J *; 4xSSC
	K	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
	L	<50	T _L *; 2xSSC	T _L *; 2xSSC
15	M	≥ 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
	N	<50	T _N *; 6xSSC	T _N *; 6xSSC
	O	≥ 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
	P	<50	T _P *; 6xSSC	T _P *; 6xSSC
	Q	≥ 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
	R	<50	T _R *; 4xSSC	T _R *; 4xSSC

[‡]: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

[†]: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

*T_B - T_R: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(°C) = 81.5 + 16.6(log₁₀[Na⁺]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer ([Na⁺] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory*

Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

The isolated polynucleotide encoding the protein of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, California, U.S.A. (the
5 MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under
10 culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-
15 toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of
20 maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available
25 from the Eastman Kodak Company (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially
30 homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, *e.g.*, as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized
35 by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means, are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

USES AND BIOLOGICAL ACTIVITY

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding

protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential
5 genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as
10 an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, those described in Gyuris *et al.*, 1993, *Cell* 75: 791-803 and in Rossi *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* 94: 8405-8410, all of which are incorporated by reference
15 herein) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in
20 assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can
25 be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

30 Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate.

- 5 In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

10

Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date,

- 15 including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, 20 CTLL2, TF-1, Mo7e and CMK. The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In

25 Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

- 30 Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon γ , Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991;

5 deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark,

10 S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins

15 that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic

20 studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

Immune Stimulating or Suppressing Activity

25 A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells

30 and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as

candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, 5 rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitis, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression 10 is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The 15 functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists 20 after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), *e.g.*, preventing high level 25 lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule 30 which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the 35 corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter

prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, *Science* 257:789-792 (1992) and Turka *et al.*, *Proc. Natl. Acad. Sci USA*, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/*lpr/lpr* mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral

diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I α chain protein and β_2 microglobulin protein or an MHC class II α chain protein and an MHC class II β chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (*e.g.*, B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B

lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al., *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512, 1988; Herrmann et al., *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann et al., *J. Immunol.* 128:1968-1974, 1982; Handa et al., *J. Immunol.* 135:1564-1572, 1985; Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bowman et al., *J. Virology* 61:1992-1998; Takai et al., *J. Immunol.* 140:508-512, 1988; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Brown et al., *J. Immunol.* 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, *J. Immunol.* 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512, 1988; Bertagnolli et al., *J. Immunol.* 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., *J. Immunol.* 134:536-544, 1995; Inaba et al., *Journal of Experimental Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine* 182:255-260, 1995; Nair et al., *Journal of Virology*

67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

Hematopoiesis Regulating Activity

A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow

transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

5 Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and*
10 *Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al.,
15 *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21,
20 Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

25

Tissue Growth Activity

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

30 A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an

osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with

the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

5 Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including
10 vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions
15 resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the
20 following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

25 Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

30 Activin/Inhibin Activity

A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers
35 with a member of the inhibin α family, may be useful as a contraceptive based on the ability of

inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- β group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation,

- those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al. Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

Hemostatic and Thrombolytic Activity

- A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

- Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

Receptor/Ligand Activity

- A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

10

Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

25

Cadherin/Tumor Invasion Suppressor Activity

Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus vulgaris and pemphigus foliaceus (auto-immune blistering skin diseases), Crohn's disease, and some developmental abnormalities.

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The

35

cadherin domains bind calcium to form their tertiary structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this recognition site can change the specificity of a cadherin so that instead of recognizing only itself, the mutant molecule can now also bind to a different
5 cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal,
10 restoring cells' adhesiveness to each other and to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the present invention
15 encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different tissue in the body.
20 Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

Additionally, proteins of the present invention with cadherin activity, and polynucleotides
25 of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this
30 decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.

Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and poly-nucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable
35 effects. Additionally, fragments of proteins of the present invention with cadherin activity,

preferably truncated soluble cadherin fragments which have been found to be stable in the circulation of cancer patients, and polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al. Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an

immune response against such protein or another material or entity which is cross-reactive with such protein.

ADMINISTRATION AND DOSING

5 A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, 10 G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent. 15 20

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form. 25

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind 30 35

surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in
5 which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such
10 liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a
15 meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination,
20 serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments
25 employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering
30 protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous,

intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01 μ g

to about 100 mg (preferably about 0.1mg to about 10 mg, more preferably about 0.1 µg to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. As used herein, the term "antibody" includes without limitation a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a single-chain antibody, a CDR-grafted antibody, a humanized antibody, or fragments thereof which bind to the indicated protein. Such term also includes any other species derived from an antibody or antibody sequence which is capable of binding the indicated protein.

Antibodies to a particular protein can be produced by methods well known to those skilled in the art. For example, monoclonal antibodies can be produced by generation of antibody-producing hybridomas in accordance with known methods (see for example, Goding, 1983, *Monoclonal antibodies: principles and practice*, Academic Press Inc., New York; and Yokoyama, 1992, "Production of Monoclonal Antibodies" in *Current Protocols in Immunology*, Unit 2.5, Greene Publishing Assoc. and John Wiley & Sons). Polyclonal sera and antibodies can be produced by inoculation of a mammalian subject with the relevant protein or fragments thereof in accordance with known methods. Fragments of antibodies, receptors, or other reactive peptides can be produced from the corresponding antibodies by cleavage of and collection of the desired fragments in accordance with known methods (see for example, Goding, supra; and Andrew et al., 1992, "Fragmentation of Immunoglobulins" in *Current Protocols in Immunology*, Unit 2.8, Greene Publishing Assoc. and John Wiley & Sons). Chimeric antibodies and single chain antibodies can also be produced in accordance with known recombinant methods (see for example, 5,169,939, 5,194,594, and 5,576,184). Humanized antibodies can also be made from corresponding murine antibodies in accordance with well known methods (see for

example, U.S. Patent Nos. 5,530,101, 5,585,089, and 5,693,762). Additionally, human antibodies may be produced in non-human animals such as mice that have been genetically altered to express human antibody molecules (see for example Fishwild *et al.*, 1996, *Nature Biotechnology* 14: 845-851; Mendez *et al.*, 1997, *Nature Genetics* 15: 146-156 (erratum *Nature Genetics* 16: 410); and U.S. Patents 5,877,397 and 5,625,126). Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, FEBS Lett. 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite,

polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins of the present invention.

The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue

(e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final
5 composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a
10 mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can
15 then be introduced *in vivo* for therapeutic purposes.

Patent and literature references cited herein are incorporated by reference as if fully set forth.

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) the nucleotide sequence of SEQ ID NO:41;
 - (b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 102 to nucleotide 2027;
 - (c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1902 to nucleotide 2027;
 - (d) the nucleotide sequence of SEQ ID NO:41 from nucleotide 1 to nucleotide 431;
 - (e) the nucleotide sequence of the full-length protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232;
 - (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;
 - (g) the nucleotide sequence of a mature protein coding sequence of clone BG160_1 deposited with the ATCC under accession number 98232;
 - (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;
 - (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
 - (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;
 - (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
 - (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:41.
2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.

4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:
 - (a) growing a culture of a host cell in a suitable culture medium, wherein the host cell has been transformed with the polynucleotide of claim 2; and
 - (b) purifying said protein from the culture.
6. A protein produced according to the process of claim 5.
7. An isolated polynucleotide encoding the protein of claim 6.
8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232.
9. A protein comprising an amino acid sequence selected from the group consisting of:
 - (a) the amino acid sequence of SEQ ID NO:42;
 - (b) the amino acid sequence of SEQ ID NO:42 from amino acid 1 to amino acid 110;
 - (c) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
 - (d) the amino acid sequence encoded by the cDNA insert of clone BG160_1 deposited with the ATCC under accession number 98232;the protein being substantially free from other mammalian proteins.
10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:42.
11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.
12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) the nucleotide sequence of SEQ ID NO:129;

- (b) the nucleotide sequence of SEQ ID NO:129 from nucleotide 383 to nucleotide 3958;
- (c) the nucleotide sequence of SEQ ID NO:129 from nucleotide 470 to nucleotide 3958;
- (d) the nucleotide sequence of SEQ ID NO:129 from nucleotide 271 to nucleotide 488;
- (e) the nucleotide sequence of the full-length protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271;
- (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;
- (g) the nucleotide sequence of a mature protein coding sequence of clone CO722_1 deposited with the ATCC under accession number 98271;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271;
- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:130;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:129.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:130;
- (b) the amino acid sequence of SEQ ID NO:130 from amino acid 1 to amino acid 34;
- (c) a fragment of the amino acid sequence of SEQ ID NO:130, the fragment comprising eight contiguous amino acids of SEQ ID NO:130; and

(d) the amino acid sequence encoded by the cDNA insert of clone CO722_1 deposited with the ATCC under accession number 98271; the protein being substantially free from other mammalian proteins.




Fig. 1A




Fig. 1B

SEQUENCE LISTING

<110> Jacobs, Kenneth
McCoy, John M.
LaVallie, Edward R.
Collins-Racie, Lisa A.
Evans, Cheryl
Merberg, David
Treacy, Maurice
Bowman, Michael R.
Spaulding, Vikki
Agostino, Michael J.
Genetics Institute, Inc.

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6011-18X

<140>
<141>

<160> 283

<170> PatentIn Ver. 2.0

<210> 1
<211> 565
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> (20)

<220>
<221> unsure
<222> (49)

<220>
<221> unsure
<222> (58)

<220>
<221> unsure
<222> (144)

<220>
<221> unsure
<222> (178)

<220>
<221> unsure
<222> (215)

<220>
<221> unsure
<222> (218)

<220>
<221> unsure

<222> (228)

<220>

<221> unsure

<222> (239)

<220>

<221> unsure

<222> (541)

<400> 1

```

gattgtcagt ttccaaaaan gaggaggatt tgatattcac ttggcccng gtgatgcntt 60
tgagggtggc cgcgtccatc tggtcagaaa agacaatctt tttgtgtca agcttgaggt 120
gtggcaggct tgagatctgg ccanacactt gaggacaat gacatccact ttgctttntc 180
tccacagggtg tccactccca ggtccaactg caganttna attcggcctt catggcctnt 240
gaatagagac ttctggactc tatagaacct actgcctcct gatgaagtcc ctactgttca 300
cccttgtagt ttttatgctc ctggcccaat tggctcagg taattggtat gtgaaaaagt 360
gtctaaacga cgttggaatt tgcaagaaga agtgcaaac tgaagagatg catacaaaga 420
ctacaagaat ttcaacagta acagcaacaa cagtcaacaa caactttgat gatgactact 480
gcttcgatgt cttcgatggc tcctaccctt ttctccact ggtggaacat tcccagcctc 540
ngtctcctgc tctaggatcc ccgac
565

```

<210> 2

<211> 94

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (87)

<400> 2

```

Met Lys Ser Leu Leu Phe Thr Leu Ala Val Phe Met Leu Leu Ala Gln
 1             5             10             15
Leu Val Ser Gly Asn Trp Tyr Val Lys Lys Cys Leu Asn Asp Val Gly
          20             25             30
Ile Cys Lys Lys Lys Cys Lys Pro Glu Glu Met His Thr Lys Thr Thr
          35             40             45
Arg Ile Ser Thr Val Thr Ala Thr Thr Val Asn Asn Asn Phe Asp Asp
          50             55             60
Asp Tyr Cys Phe Asp Val Phe Asp Gly Ser Tyr Pro Phe Leu Pro Leu
          65             70             75             80
Val Glu His Ser Gln Pro Xaa Ser Pro Ala Leu Gly Ser Pro
          85             90

```

<210> 3

<211> 92

<212> DNA

<213> Homo sapiens

<400> 3

```

aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa
92

```

<210> 4
 <211> 2949
 <212> DNA
 <213> Homo sapiens

<400> 4
 acctgctgga acgctcggac gtccttgccg gccctgtcca tgtgaggtcc aggtcatctc 60
 ccgccaccgc agacaagcgc caggatgtgc ttttggggca agacagtggc tttggtcttg 120
 tgaaggatcc atgttatctg gctggctctg gatctaggtc actcagttgt tcagaaagag 180
 gccaagaaga gatgctgccg ctcttcacc atctcaccac tcgttggggg ggttcaggct 240
 gcaaagccat tgggtgattcc tccgttccca gtgaatgtcc tggaaacctg gaccatcaga 300
 ggcaagccag taggacaccc tgccccaggc caccactggc aggaacgcaa gggctgggtca 360
 cagacaccag ggctgcaccc ctgaccccaa ttggcaccac tctgccttca gccattccct 420
 ctggctactg ctacacaggc ggtcagacag ggcgacagcc tctcccgccc tacacccttg 480
 ccatgatgca cagaagcaat ggtcacaccc tgaccacagc tcccgggtcca agaggctgtg 540
 agggcgatgg ccagagcat ggggtagaag agggaacgag gaagaggggtc tcgctgcctc 600
 agtggccacc tccttctcga gcaaagtggg ccacgcagc cagagaggac agccttccctg 660
 aggaatcctc agccccctgat tttgcaaac tgaaagcact tcaaaaacag cagagtcttc 720
 caagtttatg cagcacttct gaccagaca cactctcttg ggcccgcagc actccaggga 780
 ggatctccct ccgaatatct gactctgtcc tgcgggactc cccgccacct catgaggatt 840
 atgaagacga agtggtttgtg agggatccgc accccaagga cagctccagc cccacatttg 900
 aacctcttcc cccaccccca cctcctccac cgagtcagga aaccccgggtg tatakcatgg 960
 atgacttccc tccacctcct cccacactg tatgtgaggc gcagctggac agtgaggatc 1020
 ccgagggggc acgccccagc ttcaacaaac tttctaaagt gacaattgca agggaaaggc 1080
 acatgcctgg tgcagcccat gtggtaggta gtcagacact ggcttccaga ctccaaactt 1140
 ctatcaaggg ttccagaggct gactccacac caccctctct catgagcgtt cagcccaac 1200
 ttgctgggtc tcttgggtgg cagccagcac ccattccagc tcaaagcctc agccatgatc 1260
 cagtgcagtg aactcagggt tttagaaaaga aagtcagctc tgatcctcag aagagttcag 1320
 aagacatcag aacagaggct ttggccaagg aaattgtcca ccaagacaaa tctctagcag 1380
 acatttttga tccagactcc aggtgaaga caacaatgga cctgatggaa ggtttgtttc 1440
 cccgagatgt gaacttgcctg aaggaaaaca gtgtaaagag gaaggccata cagagaactg 1500
 tcagctcttc aggatgtgaa ggcaagagga atgaagacaa ggaagcagtg agcatgttgg 1560
 ttaactgccc tgcctactac agtgtgtctg ctcccaaggc tgagctactg aaaaaatca 1620
 aagagatgcc agcagaagtg aatgaggaag aggaacaggc agatgtcaat gaaaagaagg 1680
 ctgagctcat tggaaagtcc acccacaagc tggagaccct ccaggaggcg aaggggagcc 1740
 tgctcacgga catcaagctc aacaacgccc tgggagaaga ggtggaggct ctgatcagcg 1800
 agctctgcaa gcccaatgag tttagacaagc ataggatgtc cataggggat ttggacaagg 1860
 tgggtcaacct gctgctctcc ctctcggggc gtctagcccg tgttgagaat gtccttagcg 1920
 gccttgggtg agatgccagt aatgaagaaa ggagctctct ttacgagaaa aggaagatcc 1980
 tggctgggtc catgaggat gcccgggagc tgaaggagaa cctggatcgc agggagcgag 2040
 tagtgctggg catcttggcc aattaccttt cagaggagca gctccaggac taccagcact 2100
 tcgtgaaaat gaagtccagc ctctcattg agcaacggaa gctggatgac aagatcaagc 2160
 tgggcccagga gcaggtcaag tgcctgctgg agagcctgcc ctccagatttc attcccaagg 2220
 ctggggccct ggctctgccc ccaaacctca cgagttagcc cattcctgct gggggctgta 2280
 ctcttcagtgg tattttccca acattaacct ctccacttta acctcttcta aaatacccaa 2340
 ccaaaagatc actgtttctc tcaacactat ttaattctgaa aaatgtttca gtacaaacca 2400
 ctgtttgaac tatctgggtt attggtgttt gttcctgatg aaaggaaaaa aattctctcc 2460
 aggaggaagc ctttttccct cttgcccttc ctgattgatc ttctgagagc tcgaatgtcg 2520
 ctggacacgt accccttcta ttattacttt gtagtagaaa gaaagttaat gaaactgaga 2580
 actgattgga ggggtgttga tcatttagtt ttttaacaggc tgaggcaaca tggatcagtg 2640
 tgtgtccccc tcaggaatgt atccacagtg gccttccttg ctggtgggca gtgtatcctg 2700
 atggcagggt acaagtacca ttaatgaagg gtctgcaaca taaagcctta aaaagacaca 2760
 cactaagaaa actgtaaaaa cttgaacatt gttatttata ttttttaaaa tggaaaagat 2820
 cactatgttt gttgtgctaa ccacttatct gattctgttt tgtgggtggc atagatgatt 2880
 acgtttgagc tttgtatctt gtgaaaacct taatgaaatg aattccaaag ataaaaaaaa 2940
 aaaaaaaaaa 2949

<210> 5
 <211> 709
 <212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (255)

<400> 5

```

Met Leu Pro Leu Phe His His Leu Thr Pro Arg Trp Gly Gly Ser Gly
 1              5              10              15

Cys Lys Ala Ile Gly Asp Ser Ser Val Pro Ser Glu Cys Pro Gly Thr
      20              25              30

Leu Asp His Gln Arg Gln Ala Ser Arg Thr Pro Cys Pro Arg Pro Pro
      35              40              45

Leu Ala Gly Thr Gln Gly Leu Val Thr Asp Thr Arg Ala Ala Pro Leu
      50              55              60

Thr Pro Ile Gly Thr Pro Leu Pro Ser Ala Ile Pro Ser Gly Tyr Cys
      65              70              75              80

Ser Gln Asp Gly Gln Thr Gly Arg Gln Pro Leu Pro Pro Tyr Thr Pro
      85              90              95

Ala Met Met His Arg Ser Asn Gly His Thr Leu Thr Gln Pro Pro Gly
      100              105              110

Pro Arg Gly Cys Glu Gly Asp Gly Pro Glu His Gly Val Glu Glu Gly
      115              120              125

Thr Arg Lys Arg Val Ser Leu Pro Gln Trp Pro Pro Pro Ser Arg Ala
      130              135              140

Lys Trp Ala His Ala Ala Arg Glu Asp Ser Leu Pro Glu Glu Ser Ser
      145              150              155              160

Ala Pro Asp Phe Ala Asn Leu Lys His Tyr Gln Lys Gln Gln Ser Leu
      165              170              175

Pro Ser Leu Cys Ser Thr Ser Asp Pro Asp Thr Pro Leu Gly Ala Pro
      180              185              190

Ser Thr Pro Gly Arg Ile Ser Leu Arg Ile Ser Glu Ser Val Leu Arg
      195              200              205

Asp Ser Pro Pro Pro His Glu Asp Tyr Glu Asp Glu Val Phe Val Arg
      210              215              220

Asp Pro His Pro Lys Ala Thr Ser Ser Pro Thr Phe Glu Pro Leu Pro
      225              230              235              240

Pro Pro Pro Pro Pro Pro Pro Ser Gln Glu Thr Pro Val Tyr Xaa Met
      245              250              255

Asp Asp Phe Pro Pro Pro Pro Pro His Thr Val Cys Glu Ala Gln Leu
      260              265              270

Asp Ser Glu Asp Pro Glu Gly Pro Arg Pro Ser Phe Asn Lys Leu Ser
      275              280              285

```

Lys Val Thr Ile Ala Arg Glu Arg His Met Pro Gly Ala Ala His Val
 290 295 300
 Val Gly Ser Gln Thr Leu Ala Ser Arg Leu Gln Thr Ser Ile Lys Gly
 305 310 315 320
 Ser Glu Ala Glu Ser Thr Pro Pro Ser Phe Met Ser Val His Ala Gln
 325 330 335
 Leu Ala Gly Ser Leu Gly Gly Gln Pro Ala Pro Ile Gln Thr Gln Ser
 340 345 350
 Leu Ser His Asp Pro Val Ser Gly Thr Gln Gly Leu Glu Lys Lys Val
 355 360 365
 Ser Pro Asp Pro Gln Lys Ser Ser Glu Asp Ile Arg Thr Glu Ala Leu
 370 375 380
 Ala Lys Glu Ile Val His Gln Asp Lys Ser Leu Ala Asp Ile Leu Asp
 385 390 395 400
 Pro Asp Ser Arg Leu Lys Thr Thr Met Asp Leu Met Glu Gly Leu Phe
 405 410 415
 Pro Arg Asp Val Asn Leu Leu Lys Glu Asn Ser Val Lys Arg Lys Ala
 420 425 430
 Ile Gln Arg Thr Val Ser Ser Ser Gly Cys Glu Gly Lys Arg Asn Glu
 435 440 445
 Asp Lys Glu Ala Val Ser Met Leu Val Asn Cys Pro Ala Tyr Tyr Ser
 450 455 460
 Val Ser Ala Pro Lys Ala Glu Leu Leu Asn Lys Ile Lys Glu Met Pro
 465 470 475 480
 Ala Glu Val Asn Glu Glu Glu Gln Ala Asp Val Asn Glu Lys Lys
 485 490 495
 Ala Glu Leu Ile Gly Ser Leu Thr His Lys Leu Glu Thr Leu Gln Glu
 500 505 510
 Ala Lys Gly Ser Leu Leu Thr Asp Ile Lys Leu Asn Asn Ala Leu Gly
 515 520 525
 Glu Glu Val Glu Ala Leu Ile Ser Glu Leu Cys Lys Pro Asn Glu Phe
 530 535 540
 Asp Lys Tyr Arg Met Phe Ile Gly Asp Leu Asp Lys Val Val Asn Leu
 545 550 555 560
 Leu Leu Ser Leu Ser Gly Arg Leu Ala Arg Val Glu Asn Val Leu Ser
 565 570 575
 Gly Leu Gly Glu Asp Ala Ser Asn Glu Glu Arg Ser Ser Leu Tyr Glu
 580 585 590
 Lys Arg Lys Ile Leu Ala Gly Gln His Glu Asp Ala Arg Glu Leu Lys
 595 600 605

Glu Asn Leu Asp Arg Arg Glu Arg Val Val Leu Gly Ile Leu Ala Asn
 610 615 620
 Tyr Leu Ser Glu Glu Gln Leu Gln Asp Tyr Gln His Phe Val Lys Met
 625 630 635 640
 Lys Ser Thr Leu Leu Ile Glu Gln Arg Lys Leu Asp Asp Lys Ile Lys
 645 650 655
 Leu Gly Gln Glu Gln Val Lys Cys Leu Leu Glu Ser Leu Pro Ser Asp
 660 665 670
 Phe Ile Pro Lys Ala Gly Ala Leu Ala Leu Pro Pro Asn Leu Thr Ser
 675 680 685
 Glu Pro Ile Pro Ala Gly Gly Cys Thr Phe Ser Gly Ile Phe Pro Thr
 690 695 700
 Leu Thr Ser Pro Leu
 705

<210> 6
 <211> 190
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (4)

<220>
 <221> unsure
 <222> (30)

<220>
 <221> unsure
 <222> (97)

<220>
 <221> unsure
 <222> (154)

<220>
 <221> unsure
 <222> (174)

<400> 6
 cgangcgtct gcacccacac gctcacgaan catcaggagc ctgtctatag cgtatctttc 60
 agccctgatg ggaagtactt ggccagtga tccttcnaca agtgcgtcca tatctggaat 120
 actcagagtg gaaatcttct ccacagctac cgangcactg gcggcatctt ccangtgtgc 180
 tggaaacgccc 190

<210> 7
 <211> 391
 <212> DNA
 <213> Homo sapiens

<400> 7

```

gagtggaaag gcagggggag gggcagggag caacagaagg aagagacaac aagcccaaga 60
cagcttccat ttcagacgga agggcccccag aagatagaat tccagccgac tgaaaaacca 120
cccaatgaac aaagaagatt ttagaaaata gaaagctgtt gggattacaa agttgcgcgt 180
ttcatcggta caaactggtc tttgaacctc ctttgtgaga gcaattgtag tgtccaaatt 240
gttagggaaa acaaaaaaaaa aaaatcccaa ggaggagggt ttttccccct tccctgtttg 300
gtttatcaca gcattttgct ttttttttgg cacagctttt tacgtttctt tccattcagc 360
catcacagag cctgttccgg gtggaaacca a

```

391

<210> 8

<211> 62

<212> PRT

<213> Homo sapiens

<400> 8

```

Thr Ser Phe Val Arg Ala Ile Val Val Ser Lys Leu Leu Gly Lys Thr
  1             5             10             15

```

```

Lys Lys Lys Asn Pro Lys Glu Glu Gly Phe Ser Pro Phe Pro Val Trp
          20             25             30

```

```

Phe Ile Thr Ala Phe Cys Phe Phe Gly Thr Ala Phe Tyr Val Ser
  35             40             45

```

```

Phe His Ser Ala Ile Thr Glu Pro Val Pro Gly Gly Asn Gln
  50             55             60

```

<210> 9

<211> 267

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (21)

<400> 9

```

tctctctctc tctccctctg nacctttctc atagttgctt cagatcttag gtctcaaggg 60
cactttggcg cgtagtaagt gctttatgta agaaggcagg gcaggggggc tttttacagg 120
agaaaaaaaa atgacttata agagaaagag cctggagtat ttttgaaaaa aaaaataata 180
tttttatgtt aaaacaattt taaaatctta aaatggccat cagacataga gagctttgtg 240
tgattcatgt tttaaaaaaaa aaaaaaa

```

267

<210> 10

<211> 2178

<212> DNA

<213> Homo sapiens

<400> 10

```

cttttaacgg tcaactcggcc ttaarecgtc ctgggtgtat gtgaggctga aagggaagga 60
rgaacagggt ctgggccgct gtgtgctctg ccgtcatctc taaraaacia ccataccact 120
gaatgctact gtgtacttac aaaccacact catattcgtc acgtcatttc atcttcaccc 180
ccacctccaa atgaggcttg aaatgaagat gaagatgttc ctcctcccc tttcaaccat 240
ggacctcaca ctgtggactt cctcttagag cctctgagtt aggtacccaa gccaggcac 300
aatgtccctg cccccataa gactgcccag cccctatggc tctgatcggc tggtagagct 360
agcagccagg ctccggccag cactctgtga tactctgata accgtaggga gccaggagtt 420
ccccgccac agcctggtgc tagcagggtg cagccagcag ctgggccgca ggggccagtg 480
ggctctggga gaaggcatca gcccttctac ctttgcccag ctctgaact ttgtgtatgg 540
ggagagtgtg gagctgcagc ctggagagct aaggccctt caggaggcgg ccagggcctt 600
gggagtgcag tccctggaag aggcattgctg gagggctcga ggggacaggg ctaaaaagcc 660

```

```

agatccaggc ctgaagaaac atcaggagga gccagagaaa ccctcaagga atcctgagag 720
agaactgggg gaccctggag agaagcagaa accagaacag gtttctagaa ctggtgggag 780
agaacaggag atgttgacac agcactcgcc accaagaggc agaccgcaga tggcaggagc 840
aacgcaggag gctcagcagg aacagaccag gtcaaaggag aaacgcctcc aagcccctgt 900
tggccaaagg ggagcagatg ggaagcatgg agtgctcacg tggttgaggg aaaatccagg 960
gggctctgag gaaagtctgc gcaagctccc tggccccctt cccccagcag gctccctgca 1020
aaccagcgtc acccctaggc cctcgtgggc tgaggccccct tggttggtgg gggggccagcc 1080
tgccctgtgg agcatcctgc tgatgccgcc cagatatggc attcccttct accatagcac 1140
cccaccact ggagcctggc aggaggtctg gcgggaacag aggatccac tgctccctaaa 1200
tgccccaaa gggtcttgga gccagaacca gttggcctcc tccagcccta cccagggttc 1260
cctccccag ggccccgcac agctcagccc tggggagatg gaagagtctg atcaggggca 1320
cacaggcgca cttgcaacct gtgcgggtca tgaggacaag gcaggctgcc cactcgccc 1380
gcacctccc cgggccctc ctgctcggtc tcggccctat gcgtgctctg tctgtggaaa 1440
gaggttttca ctcaagcatc agatggagac gcactaccga gtccacacag gagagaagcc 1500
cttctcctgt agcctttgtc ctacgcgctc cggggaactt tcggccatga ccaagcacct 1560
gcggacacac gggggccgctc cgtaccgctg ctccctgtgc gggggccggt gtcccagcct 1620
ggcctccatg caggcgacac tgcccggtca ctgcgccagc caactccgc ccggatggag 1680
catccgctcc accttctct actcctctc gaggcggtct cggccctcga cctctccctg 1740
ttgtccttct tctccacca cctgacgggg tgctcggtagc gtcttagcca agagtccaat 1800
taaagaacga aaagcgggcc ggctcggctt ctgacctggc accgctgcta cggcgcccta 1860
gcaaattccg ccccgagaagc gcccaggaa gggcgccgag tgccctctcc tggacgatcg 1920
cgggtcgtag aagcccaggc cagcgagccc tacagagtga ggacactgaa gtgtgcagga 1980
gcgaaggtta acagttaggg agattgtcga tctcatcacc ataataaaga gtttctgtt 2040
gccctccctt caaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2100
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2160
aaaaaaaaaa aaaaaaaaaa

```

2178

<210> 11

<211> 487

<212> PRT

<213> Homo sapiens

<400> 11

```

Met Ser Leu Pro Pro Ile Arg Leu Pro Ser Pro Tyr Gly Ser Asp Arg
 1              5              10              15
Leu Val Gln Leu Ala Ala Arg Leu Arg Pro Ala Leu Cys Asp Thr Leu
          20              25              30
Ile Thr Val Gly Ser Gln Glu Phe Pro Ala His Ser Leu Val Leu Ala
      35              40              45
Gly Val Ser Gln Gln Leu Gly Arg Arg Gly Gln Trp Ala Leu Gly Glu
      50              55              60
Gly Ile Ser Pro Ser Thr Phe Ala Gln Leu Leu Asn Phe Val Tyr Gly
      65              70              75              80
Glu Ser Val Glu Leu Gln Pro Gly Glu Leu Arg Pro Leu Gln Glu Ala
          85              90              95
Ala Arg Ala Leu Gly Val Gln Ser Leu Glu Glu Ala Cys Trp Arg Ala
      100              105              110
Arg Gly Asp Arg Ala Lys Lys Pro Asp Pro Gly Leu Lys Lys His Gln
      115              120              125
Glu Glu Pro Glu Lys Pro Ser Arg Asn Pro Glu Arg Glu Leu Gly Asp
      130              135              140

```


Pro Gly Glu Lys Gln Lys Pro Glu Gln Val Ser Arg Thr Gly Gly Arg
 145 150 155 160
 Glu Gln Glu Met Leu His Lys His Ser Pro Pro Arg Gly Arg Pro Glu
 165 170 175
 Met Ala Gly Ala Thr Gln Glu Ala Gln Gln Glu Gln Thr Arg Ser Lys
 180 185 190
 Glu Lys Arg Leu Gln Ala Pro Val Gly Gln Arg Gly Ala Asp Gly Lys
 195 200 205
 His Gly Val Leu Thr Trp Leu Arg Glu Asn Pro Gly Gly Ser Glu Glu
 210 215 220
 Ser Leu Arg Lys Leu Pro Gly Pro Leu Pro Pro Ala Gly Ser Leu Gln
 225 230 235 240
 Thr Ser Val Thr Pro Arg Pro Ser Trp Ala Glu Ala Pro Trp Leu Val
 245 250 255
 Gly Gly Gln Pro Ala Leu Trp Ser Ile Leu Leu Met Pro Pro Arg Tyr
 260 265 270
 Gly Ile Pro Phe Tyr His Ser Thr Pro Thr Thr Gly Ala Trp Gln Glu
 275 280 285
 Val Trp Arg Glu Gln Arg Ile Pro Leu Ser Leu Asn Ala Pro Lys Gly
 290 295 300
 Leu Trp Ser Gln Asn Gln Leu Ala Ser Ser Ser Pro Thr Pro Gly Ser
 305 310 315 320
 Leu Pro Gln Gly Pro Ala Gln Leu Ser Pro Gly Glu Met Glu Glu Ser
 325 330 335
 Asp Gln Gly His Thr Gly Ala Leu Ala Thr Cys Ala Gly His Glu Asp
 340 345 350
 Lys Ala Gly Cys Pro Pro Arg Pro His Pro Pro Pro Ala Pro Pro Ala
 355 360 365
 Arg Ser Arg Pro Tyr Ala Cys Ser Val Cys Gly Lys Arg Phe Ser Leu
 370 375 380
 Lys His Gln Met Glu Thr His Tyr Arg Val His Thr Gly Glu Lys Pro
 385 390 395 400
 Phe Ser Cys Ser Leu Cys Pro Gln Arg Ser Arg Asp Phe Ser Ala Met
 405 410 415
 Thr Lys His Leu Arg Thr His Gly Ala Ala Pro Tyr Arg Cys Ser Leu
 420 425 430
 Cys Gly Ala Gly Cys Pro Ser Leu Ala Ser Met Gln Ala His Met Arg
 435 440 445
 Gly His Ser Pro Ser Gln Leu Pro Pro Gly Trp Thr Ile Arg Ser Thr
 450 455 460

Phe Leu Tyr Ser Ser Ser Arg Pro Ser Arg Pro Ser Thr Ser Pro Cys
 465 470 475 480

Cys Pro Ser Ser Ser Thr Thr
 485

<210> 12
 <211> 2436
 <212> DNA
 <213> Homo sapiens

<400> 12
 caacaacccat ctacagctgt attccttgttc ccttgatggc acaattaaac tgtgggacta 60
 tatagatggc atcttaataa agactttcat agttggatgt aaacttcacg ccctctttac 120
 tcttgcccaa gctgaggatt ctgtctttgt tatagtgaat aaagaaaaac cagatatatt 180
 tcagctgggt tcagtgaaac tgccaaaatc ctcaagccag gaagtagaag ccaaggagct 240
 gtcttttgtt ttggattaca taaaccagtc acccaagtc attgcctttg gaaacgagg 300
 agtatatgtt gctgcagtac gggaaattta cttgtctgtt tttttttca aaaagaaaac 360
 aacatcaagg tttactttat catcatcaag aaataagaag catgctaaaa acaattttac 420
 gtgtgtagca tgtcacccaa cggaagactg catcgcatct ggtcacatgg atggcaaaat 480
 ccgtctttgg aggaattttt atgatgataa gaaatatacg tacacatgtt tacattggca 540
 coatgatatg gttatggatt tggctttttc agtgacaggg accagtctgc tgagtggcgg 600
 tcgtgaatct gtactttag agtggcgcca tgcaacagag aagaataagg agtttctccc 660
 gcgttttaga gctactattg aacatatctc agtctcgcc gcaggagatt tattctgcac 720
 ttctcactct gataataaga taataattat tcaccgaaac cttgaagcat ccgcagtaat 780
 tcaaggccta gtgaaagata ggagtatctt cactggtttg atgattgac caagaactaa 840
 agctttgggt ttgaatggaa aacctggcca cctgcagttt tttctctcc agagtataa 900
 acagttatag aatttagata ttatacagca agaataatatt aatgattatg gtctgatcca 960
 aattgaacta acaaaggctg catttggctg ctttggtaac tggcttgcaa cagtggaaac 1020
 ggggcaagaa aaggaaactg agcttgaatt gcaaatgaaa ctgtggatgt ataataagaa 1080
 aacacaaggg tttattctta acactaaaat taacatgcca cacgaagact gcattacagc 1140
 tctctgtttc tgtaatgcag aaaaacttga acagcccacc ttggttacag ctagcaaaga 1200
 tggttacttc aaagtatgga tattaacaga tgactctgac atatacaaaa aagctgttgg 1260
 ctggacctgt gactttgttg gtagttatca caagtatcaa gcaactaact gttgtttctc 1320
 cgaagatggt tctttactag cagttagttt tgaggaaata gtcacaatat gggattctgt 1380
 aacatgggaa cttaaatgta cattttgcca acgagctggg aaaataaggc accttgcct 1440
 tgggagattg acgtgttcaa agtatctact cgggtgctact gaaaatggca ttctttgctg 1500
 ttggaatctg ctgagctgtg cattggagtg gaatgcaaaa ttaaatgtta gatttatgga 1560
 acccgatcct aattcagaga atattgctgc aatctctcag tcttcagtgg gttcagactt 1620
 gtttgtatth aaacctagtg agccaaggcc attgtatatt caaaagggtta tctccagaga 1680
 gaaagtccag tggggagtggt ttgttccacg agatgtccct gaatccttca cctcagaagc 1740
 ttaccagtgg ctaaatagat cccagtttta cttcctaaca aaatcacaga gtttattgac 1800
 attcagtaca aagtctccag aagaaaaact cacaccaaca agcaaacagc tgctagcaga 1860
 agaaagtctt cccacaaccc cattttatth catattggga aaacacagc aacagcagga 1920
 tgaaaaacta aacgaaactt tagagaatga gctggtacaa ctaccttaa cagaaaacat 1980
 acccgcaatt agtgagcttc ttcacactcc agcccatgtc ctgccatctg ctgctttcct 2040
 gtgctccatg tttgtaaatt cattgctgct gtctaaagag actaagagtg ctaaggaaat 2100
 tcctgaagat gtagatatgg aagaagaaaa agaaagtga gattcagatg aagaaaatga 2160
 ttttaccgaa aaagtccagg atacaagtaa cacaggttta ggagaagaca ttatacatca 2220
 gttgtcaaaa tctgaagaaa aagaactgag aaaatttagg aaaatagact acagctggat 2280
 agctgccctt taagccttgg agatggggag gatccttgga ctttgtgttt ttgattgtat 2340
 gttgatattc taaaaacatc ttttttaatg ttatttctgt tctaaaaata agataataaa 2400
 tattaacaaa ctttaaaaaa aaaaaaaaaa aaaaaa 2436

<210> 13
 <211> 763
 <212> PRT
 <213> Homo sapiens

<400> 13
 Asn Asn His Leu Gln Leu Tyr Ser Cys Ser Leu Asp Gly Thr Ile Lys
 1 5 10 15
 Leu Trp Asp Tyr Ile Asp Gly Ile Leu Ile Lys Thr Phe Ile Val Gly
 20 25 30
 Cys Lys Leu His Ala Leu Phe Thr Leu Ala Gln Ala Glu Asp Ser Val
 35 40 45
 Phe Val Ile Val Asn Lys Glu Lys Pro Asp Ile Phe Gln Leu Val Ser
 50 55 60
 Val Lys Leu Pro Lys Ser Ser Ser Gln Glu Val Glu Ala Lys Glu Leu
 65 70 75 80
 Ser Phe Val Leu Asp Tyr Ile Asn Gln Ser Pro Lys Cys Ile Ala Phe
 85 90 95
 Gly Asn Glu Gly Val Tyr Val Ala Ala Val Arg Glu Phe Tyr Leu Ser
 100 105 110
 Val Tyr Phe Phe Lys Lys Lys Thr Thr Ser Arg Phe Thr Leu Ser Ser
 115 120 125
 Ser Arg Asn Lys Lys His Ala Lys Asn Asn Phe Thr Cys Val Ala Cys
 130 135 140
 His Pro Thr Glu Asp Cys Ile Ala Ser Gly His Met Asp Gly Lys Ile
 145 150 155 160
 Arg Leu Trp Arg Asn Phe Tyr Asp Asp Lys Lys Tyr Thr Tyr Thr Cys
 165 170 175
 Leu His Trp His His Asp Met Val Met Asp Leu Ala Phe Ser Val Thr
 180 185 190
 Gly Thr Ser Leu Leu Ser Gly Gly Arg Glu Ser Val Leu Val Glu Trp
 195 200 205
 Arg Asp Ala Thr Glu Lys Asn Lys Glu Phe Leu Pro Arg Leu Gly Ala
 210 215 220
 Thr Ile Glu His Ile Ser Val Ser Pro Ala Gly Asp Leu Phe Cys Thr
 225 230 235 240
 Ser His Ser Asp Asn Lys Ile Ile Ile Ile His Arg Asn Leu Glu Ala
 245 250 255
 Ser Ala Val Ile Gln Gly Leu Val Lys Asp Arg Ser Ile Phe Thr Gly
 260 265 270
 Leu Met Ile Asp Pro Arg Thr Lys Ala Leu Val Leu Asn Gly Lys Pro
 275 280 285
 Gly His Leu Gln Phe Tyr Ser Leu Gln Ser Asp Lys Gln Leu Tyr Asn
 290 295 300
 Leu Asp Ile Ile Gln Gln Glu Tyr Ile Asn Asp Tyr Gly Leu Ile Gln
 305 310 315 320

Ile Glu Leu Thr Lys Ala Ala Phe Gly Cys Phe Gly Asn Trp Leu Ala
 325 330 335
 Thr Val Glu Gln Arg Gln Glu Lys Glu Thr Glu Leu Glu Leu Gln Met
 340 345 350
 Lys Leu Trp Met Tyr Asn Lys Lys Thr Gln Gly Phe Ile Leu Asn Thr
 355 360 365
 Lys Ile Asn Met Pro His Glu Asp Cys Ile Thr Ala Leu Cys Phe Cys
 370 375 380
 Asn Ala Glu Lys Ser Glu Gln Pro Thr Leu Val Thr Ala Ser Lys Asp
 385 390 395 400
 Gly Tyr Phe Lys Val Trp Ile Leu Thr Asp Asp Ser Asp Ile Tyr Lys
 405 410 415
 Lys Ala Val Gly Trp Thr Cys Asp Phe Val Gly Ser Tyr His Lys Tyr
 420 425 430
 Gln Ala Thr Asn Cys Cys Phe Ser Glu Asp Gly Ser Leu Leu Ala Val
 435 440 445
 Ser Phe Glu Glu Ile Val Thr Ile Trp Asp Ser Val Thr Trp Glu Leu
 450 455 460
 Lys Cys Thr Phe Cys Gln Arg Ala Gly Lys Ile Arg His Leu Cys Phe
 465 470 475 480
 Gly Arg Leu Thr Cys Ser Lys Tyr Leu Leu Gly Ala Thr Glu Asn Gly
 485 490 495
 Ile Leu Cys Cys Trp Asn Leu Leu Ser Cys Ala Leu Glu Trp Asn Ala
 500 505 510
 Lys Leu Asn Val Arg Val Met Glu Pro Asp Pro Asn Ser Glu Asn Ile
 515 520 525
 Ala Ala Ile Ser Gln Ser Ser Val Gly Ser Asp Leu Phe Val Phe Lys
 530 535 540
 Pro Ser Glu Pro Arg Pro Leu Tyr Ile Gln Lys Gly Ile Ser Arg Glu
 545 550 555 560
 Lys Val Gln Trp Gly Val Phe Val Pro Arg Asp Val Pro Glu Ser Phe
 565 570 575
 Thr Ser Glu Ala Tyr Gln Trp Leu Asn Arg Ser Gln Phe Tyr Phe Leu
 580 585 590
 Thr Lys Ser Gln Ser Leu Leu Thr Phe Ser Thr Lys Ser Pro Glu Glu
 595 600 605
 Lys Leu Thr Pro Thr Ser Lys Gln Leu Leu Ala Glu Glu Ser Leu Pro
 610 615 620
 Thr Thr Pro Phe Tyr Phe Ile Leu Gly Lys His Arg Gln Gln Gln Asp
 625 630 635 640

Glu Lys Leu Asn Glu Thr Leu Glu Asn Glu Leu Val Gln Leu Pro Leu
 645 650 655
 Thr Glu Asn Ile Pro Ala Ile Ser Glu Leu Leu His Thr Pro Ala His
 660 665 670
 Val Leu Pro Ser Ala Ala Phe Leu Cys Ser Met Phe Val Asn Ser Leu
 675 680 685
 Leu Leu Ser Lys Glu Thr Lys Ser Ala Lys Glu Ile Pro Glu Asp Val
 690 695 700
 Asp Met Glu Glu Glu Lys Glu Ser Glu Asp Ser Asp Glu Glu Asn Asp
 705 710 715 720
 Phe Thr Glu Lys Val Gln Asp Thr Ser Asn Thr Gly Leu Gly Glu Asp
 725 730 735
 Ile Ile His Gln Leu Ser Lys Ser Glu Glu Lys Glu Leu Arg Lys Phe
 740 745 750
 Arg Lys Ile Asp Tyr Ser Trp Ile Ala Ala Leu
 755 760

<210> 14
 <211> 137
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (2)

<220>
 <221> unsure
 <222> (18)

<220>
 <221> unsure
 <222> (29) .. (30)

<220>
 <221> unsure
 <222> (36)

<220>
 <221> unsure
 <222> (44)

<220>
 <221> unsure
 <222> (49)

<220>
 <221> unsure
 <222> (62)

<220>

<221> unsure
<222> (67)

<220>
<221> unsure
<222> (80)

<220>
<221> unsure
<222> (93)

<220>
<221> unsure
<222> (99)

<220>
<221> unsure
<222> (108)

<220>
<221> unsure
<222> (112)

<220>
<221> unsure
<222> (121)

<400> 14
gnggggggagg atgaatantc ggaagctggn gcggcngcgg gganccaang aagaggttcc 60
gnacccngct tacgaaccgn ggactcttga ctncaccaac cctgaacnct cncctccaaa 120
ngcgtcatgg atggaat 137

<210> 15
<211> 539
<212> DNA
<213> Homo sapiens

<400> 15
gtggagggag agtgaattct gggaatctct cagcagcttt ttgccaaaca gatgggcccag 60
gagccgcgga accaggctga ggaatgttgc ctcacgatct ctcatatcca ttcttggcac 120
ccaccagccc aggggaatgcc tctaccagtt gtcagcgaga ggcttacaca gcattctaaa 180
taaaagggat tattgaacca agaggccagg gactgatgga aatgcccacc ttgctggctc 240
attgaaaaag tttggcaagg ttgtcaggag acatgaatta gatgggcttg ggtcttgtgc 300
cctttgtctaa accaagtgtc gtattgggaa agagacgggg agagaagtgt tggagatgct 360
cttttagtcag gcctgagtca cttgcccac cctggagtgt gagttgggga tggagccagg 420
atctccaaac cacatgcccc tagagtttca gggaaaatat ggattgtgaa ttgaagatgg 480
ggggtgatgt aaggcagaca aggacagaaa atccctcttc cagctgtgat ttggctgtg 539

<210> 16
<211> 27
<212> PRT
<213> Homo sapiens

<400> 16
Met Asp Cys Glu Leu Lys Met Gly Gly Asp Val Arg Gln Thr Arg Thr
1 5 10 15
Glu Asn Pro Ser Ser Ser Cys Asp Leu Ala Val
20 25

<210> 17
 <211> 99
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (50)

<220>
 <221> unsure
 <222> (55)..(56)

<220>
 <221> unsure
 <222> (62)

<220>
 <221> unsure
 <222> (64)

<220>
 <221> unsure
 <222> (84)

<400> 17
 agagaggcag atccccaatg aggggtcagg acgggtcttg gctgcacatn ctggmntcat 60
 antnatccca tggggagcag cacnttatga aaaaaaaaaa 99

<210> 18
 <211> 2608
 <212> DNA
 <213> Homo sapiens

<400> 18
 attagccact gcacctggct ggtgtctcca ttttacagat gaggtctgtca cccccaata 60
 tcacacagcc aagtggcaga gttgtgttca aatgctggca gtcaggccat gctctgtctg 120
 artccaract ctgccacttt ctarctgtgt gttctcgggc aggttatatta ccaactttga 180
 agtccatttt cttatcccta aaagagggtc gagaagctca ccctacett tgaaggttgt 240
 ttgaggatta agcaagacca gcatctgaag ggcctgaagc atctagggca tcatcagtgc 300
 actttatctc tggcctctaa gaagtgcac tagaaggac cactgtcac agcttccttg 360
 tacccttgcc ttgtcctgtc aggtagcact tcggtttggc cctcactttt tcaggatttg 420
 ttggaacaa gatcccttat cttctctagg tcagatttgg ggtgaacttg gctgacagta 480
 ttccagggca ggcagaggtc aggttggtg gtgacctgct gcctccttcc catcggcagc 540
 tagttctagg gccatccaca gggatcccca gccctccata gtaccagtga gcctcactgg 600
 catgtgccag ggccaagagg gccttcagcg tcgcctccag tccagccctg tcctcgtatt 660
 cttggggaca ctgtggctca gccagcacag tgcttagccc acagtacccc agcaggcgtc 720
 cctcgagtgg gccagtggca ggcacagagc tgctctcccc ttgcctggat gcgtgaggaa 780
 aggaactctt ctccctgcag ccagccctg caggctctct gctgccagat gacgagttaa 840
 gatgaaagtc ctgatgtcta ttgtctgtgt gcacagagct ttggagtggc ctggccagga 900
 agagggggaa cgtgtgtggg gtcagcaciaa ggggttgccc ggatcttggg tactggttct 960
 gctctgctgc tggttccctc tgaaaaatgtg ggtttctccc ctccgtgctg agcctcagaa 1020
 aggggcaggg agcccatgc cagcacacat ggcccctcgg ctctctggcc tctgcagggt 1080
 cagcccagct ccttcacctc tctcctgctt tcccctcagg cctcagtgc cccacacccc 1140
 cttcaggccc aagcctggat ggtaccggat ggaacagctt ccagtcgtcg gatgccactg 1200
 agccccccag cccctgctct ggcccaggcc ccagtatgg aaagccgacc ccagcgagc 1260
 acatccctgc cagcaagcag cggctctggac gacctagacc tcctggggaa gacctcctg 1320
 cagcagtcgc tgccccgga atcccagcaa gtgcggtggg agaagcagca gccaaacccc 1380
 cggctcacac tccgggacct gcagaataag agcagcagct gcagctcccc cagctccagc 1440

gccaccagcc ttctccacac cgtgtcccca gagcccccca ggctccgca gcagcccgta 1500
 ccaaccgagc tctcactggc cagcatcact gtgcccctgg agtccatcaa acccagcaac 1560
 atcctgcccc tgactgtgta tgaccagcac ggcttccgca tcctcttcca ttttgccegg 1620
 gaccactgc cagggcgctc cgacgtgctg gtgggtggtg tttccatgct gagcaccgcc 1680
 ccccagccca tccgcaacat cgtgttccag tcagctgtcc ccaaggttat gaaggtgaag 1740
 ctgcagccac cctcgggcac ggagctgcca gcttttaacc ccatcgctcca cccctcagca 1800
 atcacccagg tcttctgtct tgccaacccc cagaaggaga aggttcgcct ccgctacaag 1860
 ctcaccttca ccatgggtga ccagacctac aacgagatgg gggatgtgga ccagtcccc 1920
 ccacctgaaa cctggggtag cctctagaac agaggggctg gggagaggaa ggggcagagg 1980
 gaccggtcac tgtccagcct ggagggagggc attggtggcc aaggacaccc tttgttgccc 2040
 atggccattc acccccaggc ctggtgcttc tccccacacc cctgtaggcc tcaagtact 2100
 cttccccctc ctgctccggc cccgcccctg ctgagccaaa ccagtagga ggctgggcct 2160
 gggtttgtgc cgctggggtc tccatcacgg ggacctggag agggaggggc tgtgtagcct 2220
 tggaagaact tgggtcatgg ggaggaagca cagctgttgg ggaagggcca ggacctcagg 2280
 cccagcccca accccagctg ggggtgggtc tccccacct gtctcttatg ccttatggga 2340
 agggccagcc ataactcggg ggccatgctc gagctgggga ccagcttagg cctcctatgg 2400
 gtgtcttacg tctgtccatc catctgtccg tggtcagaag tggggtcagt gtgtgagtga 2460
 gagcaggagt atttatgaaa ataaaacgtc gtttttctcg gaaaaaaaaa aaaaaaaaaa 2520
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2580
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2608

<210> 19

<211> 236

<212> PRT

<213> Homo sapiens

<400> 19

Met Glu Ser Arg Pro Pro Ala Gln Thr Ser Leu Pro Ala Ser Ser Gly
 1 5 10 15
 Leu Asp Asp Leu Asp Leu Leu Gly Lys Thr Leu Leu Gln Gln Ser Leu
 20 25 30
 Pro Pro Glu Ser Gln Gln Val Arg Trp Glu Lys Gln Gln Pro Thr Pro
 35 40 45
 Arg Leu Thr Leu Arg Asp Leu Gln Asn Lys Ser Ser Ser Cys Ser Ser
 50 55 60
 Pro Ser Ser Ser Ala Thr Ser Leu Leu His Thr Val Ser Pro Glu Pro
 65 70 75 80
 Pro Arg Pro Pro Gln Gln Pro Val Pro Thr Glu Leu Ser Leu Ala Ser
 85 90 95
 Ile Thr Val Pro Leu Glu Ser Ile Lys Pro Ser Asn Ile Leu Pro Val
 100 105 110
 Thr Val Tyr Asp Gln His Gly Phe Arg Ile Leu Phe His Phe Ala Arg
 115 120 125
 Asp Pro Leu Pro Gly Arg Ser Asp Val Leu Val Val Val Val Ser Met
 130 135 140
 Leu Ser Thr Ala Pro Gln Pro Ile Arg Asn Ile Val Phe Gln Ser Ala
 145 150 155 160
 Val Pro Lys Val Met Lys Val Lys Leu Gln Pro Pro Ser Gly Thr Glu
 165 170 175

Leu Pro Ala Phe Asn Pro Ile Val His Pro Ser Ala Ile Thr Gln Val
 180 185 190

Leu Leu Leu Ala Asn Pro Gln Lys Glu Lys Val Arg Leu Arg Tyr Lys
 195 200 205

Leu Thr Phe Thr Met Gly Asp Gln Thr Tyr Asn Glu Met Gly Asp Val
 210 215 220

Asp Gln Phe Pro Pro Pro Glu Thr Trp Gly Ser Leu
 225 230 235

<210> 20

<211> 328

<212> DNA

<213> Homo sapiens

<400> 20

agacttcgaa ttcggccttc atggcctara aaaactgatt cacctggcag agacctaccc 60
 catccacatg cacagccagc tggaccacct tagcctctat tactgcaggt gtactctgcc 120
 agagaatcca aacaatcaca cctccagta ctggaaggac cacaacatcg tgacagcaga 180
 agtccactgg gctaacctga ctgtcagtga atgccaggag atgcatggag agttcatggg 240
 atctgcgtgc ggccatcatg gacctacac tcctgatgtc ctcttttggt cctgtattct 300
 ctttttcacc accttcatcc tctcaagc 328

<210> 21

<211> 87

<212> PRT

<213> Homo sapiens

<400> 21

Met His Ser Gln Leu Asp His Leu Ser Leu Tyr Tyr Cys Arg Cys Thr
 1 5 10 15

Leu Pro Glu Asn Pro Asn Asn His Thr Leu Gln Tyr Trp Lys Asp His
 20 25 30

Asn Ile Val Thr Ala Glu Val His Trp Ala Asn Leu Thr Val Ser Glu
 35 40 45

Cys Gln Glu Met His Gly Glu Phe Met Gly Ser Ala Cys Gly His His
 50 55 60

Gly Pro Tyr Thr Pro Asp Val Leu Phe Trp Ser Cys Ile Leu Phe Phe
 65 70 75 80

Thr Thr Phe Ile Leu Ser Ser
 85

<210> 22

<211> 326

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (1)

<220>
 <221> unsure
 <222> (31)

<220>
 <221> unsure
 <222> (82)

<220>
 <221> unsure
 <222> (84)

<220>
 <221> unsure
 <222> (129)

<400> 22
 naaatgtatt tatatgtatg tcttattata nacaaggcag atttcccctg gaataaaaagt 60
 ctagaatgta ctgcttaatt tnanacatgt gtgcaggcaa tattatctgt gaggtaaaag 120
 tggataana cgtggattgg gtcaactgat ttcagcttg ttaggagtc tctgtgtgag 180
 acatgggtgtg ataattgtga agttctcact gtatgtggat gttcatgtga aagatagtag 240
 tttcttcccg taaatatctt ttgatttcca tttgtatgga atcccaatga atgtatcttt 300
 ggaaaacaaa aaaaaaaaaa aaaaaa 326

<210> 23
 <211> 194
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (55)

<220>
 <221> unsure
 <222> (73)

<220>
 <221> unsure
 <222> (119)

<220>
 <221> unsure
 <222> (126)

<220>
 <221> unsure
 <222> (137)

<220>
 <221> unsure
 <222> (174)

<400> 23
 aagtctcatt ctcaagagtg atgctgcaaa accccttttg ggccatgatg ctgtnatcca 60
 ggcttttagca canaaaggtc tttatgtcac tgaccaggaa aaattggtaa ctgaacgana 120
 tctccncaag aaaccntac agatgagtgc acatttgggc catgatcgat accncatgat 180
 ggcttatact gtat 194

<210> 24

<211> 396
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (139)

<400> 24
 ttcattttta ctgtcctgat gttgtcagat tagaggatat ttgtttgaaa gaaactatgt 60
 ctttggtctga tagcctgtat aatctgcagc tgattcaaga attttgccaa gaatacttga 120
 accagtgttg ccatttcant ctggaagata tgctctatgc tgcttcatcc ataaagagta 180
 attatttgggt gttcatggcg gaactgttct ggtgggttga agtgggtgaag ccgtcttttg 240
 tacagcctcg tgttgttcgt ccacaaggag ctgaacctgt aaaagatatg ccttcaattc 300
 ctgtcttgaa tgctgccaaa agaaatgtct tagatagtag ttctgacttc ccttcaagt 360
 gggaaggagc tacatttaca cagtctcatc tcgagg 396

<210> 25
 <211> 113
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (28)

<400> 25
 Met Ser Leu Ala Asp Ser Leu Tyr Asn Leu Gln Leu Ile Gln Glu Phe
 1 5 10 15
 Cys Gln Glu Tyr Leu Asn Gln Cys Cys His Phe Xaa Leu Glu Asp Met
 20 25 30
 Leu Tyr Ala Ala Ser Ser Ile Lys Ser Asn Tyr Leu Val Phe Met Ala
 35 40 45
 Glu Leu Phe Trp Trp Phe Glu Val Val Lys Pro Ser Phe Val Gln Pro
 50 55 60
 Arg Val Val Arg Pro Gln Gly Ala Glu Pro Val Lys Asp Met Pro Ser
 65 70 75 80
 Ile Pro Val Leu Asn Ala Ala Lys Arg Asn Val Leu Asp Ser Ser Ser
 85 90 95
 Asp Phe Pro Ser Ser Gly Glu Gly Ala Thr Phe Thr Gln Ser His Leu
 100 105 110
 Glu

<210> 26
 <211> 336
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (87)

<220>

<221> unsure

<222> (123)

<400> 26

aatttgatga tgaccagaaa gtatgctgtg gttctttttt aaggatgac aaaaagcaga 60
 aaatgatatg gcaatgaaac gggcagnttt gttggagaaa agattaagaa gggaaaagga 120
 aantcagctc cggaacaac agttggaagc agaaatggag cataagaagg aggaacaag 180
 gcgtaadact gaggaagaac gtcagaagaa agaagatgag agagcacgca gagaatttat 240
 taggcaagaa tatatgaggc ggaacaact gaaactaatg gaagatatgg atacagtaat 300
 taaaccccgct cctcaagtag taaaaaaaa aaaaaa 336

<210> 27

<211> 917

<212> DNA

<213> Homo sapiens

<400> 27

cacctggatt atctcagtag tttcccaact ggtttccttg tttccattct tgccctcttc 60
 tgtctactct caatataaca gctagaacaa tccttttaca atggaattca gatcatgttt 120
 accctctgtg tcaaattctc cagtgaactt ccagttttta catgatctgg ctctactac 180
 ctgtctcact gtgtttccta ctactctctt gccctttctc ctcttaataa aactgggct 240
 catgggtgtt cctttaacat gccaggcatg cttgacctg tcctgtctca gggccctgct 300
 gttccctctg cctggaacat tcttccata gtgtctgcat ggctcgctct ctactgctt 360
 tggattgctg ctcaaaagtc acctatcaaa aggcctttcc caaaggttta aaaatcattc 420
 tactataaag acacatgcat acatatgttt attgcagcac tattcacaat acaaaagact 480
 tggaaccaac ccaaatgccc atcaatgata gactggataa agaaaatatg gcacgtaagc 540
 accatggaat actatgcagc cataaaaaag aatgagtcca tgtcctttgc agggacatgg 600
 atgaagctgg aaaccattat tctcagcaaa ctaacacagg aacagaaaac caaacaccgc 660
 atgttctcac tcataagtgg gagttgaaca atgagaacat acgggcacag tggggggaac 720
 atcacacacc agggcctgtc ggggggtgag aggcaaggga agtgatagca ttaagagaaa 780
 tacctaattg agattatggg ttgatggggg cagcaaacca ccatggcaca tgtgtacct 840
 tgtaacaaac ctgcacattc tgcacatata tcccagaact taaagtataa ttaagaaaa 900
 agaaaaaaaa aaaaaa 917

<210> 28

<211> 76

<212> PRT

<213> Homo sapiens

<400> 28

Met Glu Phe Arg Ser Cys Leu Pro Leu Cys Ser Asn Ser Pro Val Thr
 1 5 10 15
 Phe Gln Phe Leu His Asp Leu Ala Pro Thr Thr Cys Leu Thr Val Phe
 20 25 30
 Pro Thr Thr Leu Leu Pro Phe Leu Leu Leu Ile Asn Thr Gly Leu Met
 35 40 45
 Val Phe Pro Leu Thr Cys Gln Ala Cys Leu Thr Leu Ser Cys Leu Arg
 50 55 60
 Ala Leu Leu Phe Pro Leu Pro Gly Thr Phe Phe Pro
 65 70 75

<210> 29

<211> 351

<212> DNA

<213> Homo sapiens

<400> 29

```

ggcttttgacc gctatcgcca ggagtggatg gactatggct gtgcacagga ggcagagggc 60
aggatgtgcg aggacttcca ggatgaggac cacgactcag cctcccctga cacttccttc 120
agcccctatg atggagacct caccamtacc tcctcctccc tcttcacga cagcctcacc 180
acagaagatg acaccaagtt gaatccctat gcaggaggag acggccttca gaacaacctg 240
tcccccaaga caaagggcac tcctgtgcac ctggggacca tcgtgggcat cgtgctggca 300
gtcctcctcg tggcggccat catcctggct ggaatttaca tcaatggcca c 351

```

<210> 30

<211> 108

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (40)

<400> 30

```

Met Asp Tyr Gly Cys Ala Gln Glu Ala Glu Gly Arg Met Cys Glu Asp
 1              5              10              15
Phe Gln Asp Glu Asp His Asp Ser Ala Ser Pro Asp Thr Ser Phe Ser
      20              25              30
Pro Tyr Asp Gly Asp Leu Thr Xaa Thr Ser Ser Ser Leu Phe Ile Asp
      35              40              45
Ser Leu Thr Thr Glu Asp Asp Thr Lys Leu Asn Pro Tyr Ala Gly Gly
      50              55              60
Asp Gly Leu Gln Asn Asn Leu Ser Pro Lys Thr Lys Gly Thr Pro Val
      65              70              75              80
His Leu Gly Thr Ile Val Gly Ile Val Leu Ala Val Leu Leu Val Ala
      85              90              95
Ala Ile Ile Leu Ala Gly Ile Tyr Ile Asn Gly His
      100              105

```

<210> 31

<211> 179

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (24)

<220>

<221> unsure

<222> (33)

<220>

<221> unsure

<222> (56)

<220>
 <221> unsure
 <222> (82)

<220>
 <221> unsure
 <222> (93)

<220>
 <221> unsure
 <222> (99)

<220>
 <221> unsure
 <222> (117)

<220>
 <221> unsure
 <222> (137)

<400> 31
 gttgtcagcc ccttcctttc ccngaccta aanataaaag acaaggcaaa gcccgntaa 60
 ttttaagacg gtttttttagg anattagtc acnattttnt tggtttgatg gttttcngaa 120
 ataaagtccc tttcctngct ccaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 179

<210> 32
 <211> 3906
 <212> DNA
 <213> Homo sapiens

<400> 32
 ggaaactcag ccacctgtga caaatttgag tgtctctggt gaaaacctct gcacagtaat 60
 atggacatgg aatccacccg agggagccag ctcaaattgt agtctatggt attttagtca 120
 ttttggcgac aaacaagata agaaaatagc tccggaaact cgtcgttcaa tagaagtacc 180
 cctgaatgag aggatttgtc tgcaagtggg gtcccagtggt agcaccaatg agagtggaga 240
 gcctagcatt ttggttgaaa aatgcatctc acccccagaa ggtgatcctg agtctgctgt 300
 gactgagctt caatgcattt ggcaacaact gagctacatg aagtgttctt ggctccctgg 360
 aaggaatacc agtcccagaca ctaactatac tctctactat tggcacagaa gcctgggaaa 420
 aattcatcaa tgtgaaaaca tcttttagaga agggcaatac ttgggttggt cctttgatct 480
 gaccaaagtg aaggattcca gttttgaaca acacagtgtc caaataatgg tcaaggataa 540
 tgcagggaaa attaaaccat ccttcaatat agtgccttta acttcccgtg tgaaacctga 600
 tcttccacat attaaaaacc tctccttcca caatgatgac ctatatgtgc aatgggagaa 660
 tccacagaat tttatttagc gatgcctatt ttatgaagta gaagtcaata acagccaaac 720
 tgagacacat aatgttttct acgtccaaga ggctaaatgt gagaatccag aatttgagag 780
 aaatgtggag aatacatctt gtttcatggt ccctggtggt ctctctgata ctttgaacac 840
 agtcagaata agagtcaaaa caaataagtt atgctatgag gatgacaaac tctggagtaa 900
 ttggagccaa gaaatgagta taggtaagaa gcgcaattcc acactctaca taaccatggt 960
 actcattggt ccagtcacgt tgcgaggtgc aatcatagta ctctgctttt acctaaaaag 1020
 gctcaagatt attatattcc ctccaattcc tgatcctggc aagattttta aagaaatggt 1080
 tggagaccag aatgatgata ctctgcactg gaagaagtac gacatctatg agaagcaaac 1140
 caaggaggaa accgactctg tagtgctgat agaaaacctg aagaaagcct ctcaagtgatg 1200
 gagataattt atttttacct tcaactgtgac cttgagaaga ttcttcccat tctccatttg 1260
 ttatctggga acttattaaa tggaaactga aactactgca ccatttataa acaggcagct 1320
 cataagagcc acaggtcttt atgttgagtc gcgcaccgaa aaactaaaaa taatgggagc 1380
 tttggagaag agtgtggagt catttccatt gaattataaa agccagcagg cttcaacta 1440
 ggggacaaag caaaaagtga tgatagtggt ggagttaatc ttatcaagag ttgtgacaac 1500
 tctctgaggg atctatactt gctttgtggt ctttgtgtca acatgaacaa attttatttg 1560
 taggggaact catttggggt gcaaatgcta atgtcaaaact tgagtcacaa agaacatgta 1620
 gaaaacaaaa tggataaaat ctgatatgta ttgtttggga tcctattgaa ccatgtttgt 1680
 ggctattaaa actcttttaa cagtcctggg tgggtccggg ggctcacgcc tgtaatccca 1740

gcaatttggg agtccgaggc gggcggatca ctcgaggtca ggagttccag accagcctga 1800
 ccaaattggt gaaacctcct ctctactaaa actacaaaaa ttaactgggt gtggtggcgc 1860
 gtgcctgtaa tcccagctac tcgggaagct gaggcagggtg aattgtttga acctgggagg 1920
 tggaggttgc agtgagcaga gatcacacca ctgcactcta gcctgggtga cagagcaaga 1980
 ctctgtctaa aaaacaaaac aaaacaaaac aaaacaaaaa aacctcttaa tattctggag 2040
 tcacattcc ctcgacagc atttctctct gctttgaaag cccagaaaat cagtgttggc 2100
 catgatgaca actacagaaa aaccagaggc agcttctttg ccaagacctt tcaaagccat 2160
 tttaggctgt taggggcagt ggaggtagaa tgactccttg ggtattagag tttcaacct 2220
 gaagtctcta acaatgtatt ttcttcacct ctgctactca agtagcattt actgtgtctt 2280
 tggtttgtgc taggcccccg ggtgtgaagc acagaccctt tccaggggtt tacagtctat 2340
 ttgagactcc tcagttcttg ccactttttt tttaatctcc accagtcatt tttcagacct 2400
 tttactcct caattccaac actgatttcc ccttttgcct tctcctcctt tcccttcctt 2460
 gtacgctttt gactttcatt ggaaattagg atgtaaatct gctcaggaga cctggaggag 2520
 cagaggataa ttagcatctc aggttaagtg tgagtaatct gagaacaat gactaattct 2580
 tgcattttt gtaacttcca tgtgagggtt ttcagcattg atatttgtgc attttctaaa 2640
 cagagatgag gtggtatctt cacgtagaac attggtattc gcttgagaaa aaaagaatag 2700
 ttgaacctat ttctctttct ttacaagatg ggtccaggat tcctctttt tctgccataa 2760
 atgattaatt aaatagcttt tgtgtcttac attggtagcc agccagccaa ggctctgttt 2820
 atgcttttgg ggggcatata ttgggttcca ttctcaccta tccacacaac atatccgtat 2880
 atatccctc tactcttact tcccccaat ttaaagaagt atgggaaatg agaggcattt 2940
 cccccacccc atttctctcc tcacacacag actcatatta ctggtaggaa cttgagaact 3000
 ttatttccaa gttgttccaa catttaccaa tcatattaat acaatgatgc tatttgcaat 3060
 tcctgtctct aggggagggg agataagaaa cctcactct ctacagggtt ggggtacaagt 3120
 ggcaacctgc ttccatggcc gtgtagaagc atgggtgccct ggcttctctg aggaagctgg 3180
 ggttcatgac aatggcagat gtaaagtat tcttgaagtc agattgaggc tgggagacag 3240
 ccgtagtaga tgttctactt tgttctgctg ttctctagaa agaataattt gtttctctgt 3300
 ataggaatga gattaattcc ttccaggta tttataatt ctgggaagca aaacccatgc 3360
 ctccccctag ccatttttac tgttatccta tttagatggc catgaagagg atgctgtgaa 3420
 attccaaca aacattgatg ctgacagtc tgcagtctgg gagggggaa gtgatctttt 3480
 gttcccatcc tcttctttta gcagtaaaat agctgaggga aaagggaggg aaaaggaagt 3540
 tatgggaata cctgtggtgg ttgtgatccc taggtcttgg gagctcttgg aggtgtctgt 3600
 atcagtggat tccccatccc ctgtgggaaa ttagtaggct catttactgt tttagggtcta 3660
 gcctatgtgg attttttcct aacatacyta agcaaaccca gtgtcaggat ggtaattctt 3720
 attctttcgt tcagttaagt tttcccttc atctgggcac tgaagggaata tgtgaaacaa 3780
 tgttaacatt tttggtagtc ttcaaccagg gattgtttct gtttaacttc ttataggaaa 3840
 gcttgagtaa aataaatatt gtctttttgt atgtcaccca aaaaaaaaaa aaaaaaaaaa 3900
 aaaaaa 3906

<210> 33

<211> 286

<212> PRT

<213> Homo sapiens

<400> 33

Met Lys Cys Ser Trp Leu Pro Gly Arg Asn Thr Ser Pro Asp Thr Asn
 1 5 10 15

Tyr Thr Leu Tyr Trp His Arg Ser Leu Glu Lys Ile His Gln Cys
 20 25 30

Glu Asn Ile Phe Arg Glu Gly Gln Tyr Leu Gly Cys Ser Phe Asp Leu
 35 40 45

Thr Lys Val Lys Asp Ser Ser Phe Glu Gln His Ser Val Gln Ile Met
 50 55 60

Val Lys Asp Asn Ala Gly Lys Ile Lys Pro Ser Phe Asn Ile Val Pro
 65 70 75 80

Leu Thr Ser Arg Val Lys Pro Asp Pro Pro His Ile Lys Asn Leu Ser

85 90 95
 Phe His Asn Asp Asp Leu Tyr Val Gln Trp Glu Asn Pro Gln Asn Phe
 100 105 110
 Ile Ser Arg Cys Leu Phe Tyr Glu Val Glu Val Asn Asn Ser Gln Thr
 115 120 125
 Glu Thr His Asn Val Phe Tyr Val Gln Glu Ala Lys Cys Glu Asn Pro
 130 135 140
 Glu Phe Glu Arg Asn Val Glu Asn Thr Ser Cys Phe Met Val Pro Gly
 145 150 155 160
 Val Leu Pro Asp Thr Leu Asn Thr Val Arg Ile Arg Val Lys Thr Asn
 165 170 175
 Lys Leu Cys Tyr Glu Asp Asp Lys Leu Trp Ser Asn Trp Ser Gln Glu
 180 185 190
 Met Ser Ile Gly Lys Lys Arg Asn Ser Thr Leu Tyr Ile Thr Met Leu
 195 200 205
 Leu Ile Val Pro Val Ile Val Ala Gly Ala Ile Ile Val Leu Leu Leu
 210 215 220
 Tyr Leu Lys Arg Leu Lys Ile Ile Ile Phe Pro Pro Ile Pro Asp Pro
 225 230 235 240
 Gly Lys Ile Phe Lys Glu Met Phe Gly Asp Gln Asn Asp Asp Thr Leu
 245 250 255
 His Trp Lys Lys Tyr Asp Ile Tyr Glu Lys Gln Thr Lys Glu Glu Thr
 260 265 270
 Asp Ser Val Val Leu Ile Glu Asn Leu Lys Lys Ala Ser Gln
 275 280 285

 <210> 34
 <211> 1605
 <212> DNA
 <213> Homo sapiens

 <400> 34
 acgctgacca tcacaggcac acagaggcac atccacctca catccacctc atacttggtgt 60
 actctcaggg ttcagtcttt catcctatcc ctctctgata tgtgctctcc aataccttcc 120
 aagatgttta cagagaccct tctccctgtg cagttaggag tgtaaggcaa gagagcccct 180
 acttcatggg gcagatcaag agctgagacc aaagatgggc tatgttgctg accttgctct 240
 gtccctctgc tgccttaaac tatgatccct gctgcgggtc ctgaagcctt tccctgtgag 300
 cagtgggtgt tgagagccag gcgtccctct gcctgcccac tcagtggcaa caccggggag 360
 ctgttttgtc ctttgtggag cctcagcagt tccctctttc agaactcact gccaagagcc 420
 ctgaacagga gccaccatgc agtgcttcag cttcattaag accatgatga tctcttcaa 480
 tttgctcatc tttctgtgtg gtgcagccct gttggcagtg ggcattctggg tgtcaatcga 540
 tggggcatcc tttctgaaga tcttcggggc actgtcgtcc agtgccatgc agtttgtcaa 600
 cgtgggctac ttcctcatcg cagccggcgt tgtggtcttt gctcttggtt tccctgggctg 660
 ctatgggtgt aagactgaga gcaagtgtgc cctcgtgacg tctcttctca tccctctcct 720
 catcttcatt gctgaggttg cagctgctgt ggtcgccctg gtgtacacca caatggctga 780
 gcacttctct acgttgctgg tagtgctgc catcaagaaa gattatgggt cccaggaaga 840
 cttcactcaa gtgtggaaca ccaccatgaa agggctcaag tgctgtgggt tcaccaacta 900


```

tacggatttt gaggactcac cctacttcaa agagaacagt gcctttcccc cattctggtg 960
caatgacaac gtcaccaaca cagccaatga aacctgcacc aagcaaaaagg ctcacgacca 1020
aaaagtagag ggttgcttca atcagctttt gtatgacatc cgaactaatg cagtcaccgt 1080
gggtggtgtg gcagctggaa ttgggggcct cgagctggtt gccatgattg tgtccatgta 1140
tctgtactgc aatctacaat aagtccactt ctgcctctgc cactactgct gccacatggg 1200
aactgtgaag aggcaccctg gcaagcagca gtgattgggg gaggggacag gatctaacaa 1260
tgtcacttgg gccagaatgg acctgccctt tctgctccag acttgggggt agatagggac 1320
cactcctttt aggcgatgcc tgactttcct tccattggtg ggtggatggg tggggggcat 1380
tccagagcct ctaaggtagc cagttctggt gccattccc ccagtctatt aaacccttga 1440
tatgccccct aggcctagtg gtgatcccag tgctctactg ggggatgaga gaaaggcatt 1500
ttatagcctg ggcataagtg aaatcagcag agcctctggg tggatgtgta gaaggcatt 1560
caaaatgcat aaacctgtta caatgttgaa aaaaaaaaaa aaaaaa 1605

```

<210> 35

<211> 241

<212> PRT

<213> Homo sapiens

<400> 35

```

Met Gln Cys Phe Ser Phe Ile Lys Thr Met Met Ile Leu Phe Asn Leu
  1             5             10             15

Leu Ile Phe Leu Cys Gly Ala Ala Leu Leu Ala Val Gly Ile Trp Val
          20             25             30

Ser Ile Asp Gly Ala Ser Phe Leu Lys Ile Phe Gly Pro Leu Ser Ser
          35             40             45

Ser Ala Met Gln Phe Val Asn Val Gly Tyr Phe Leu Ile Ala Ala Gly
          50             55             60

Val Val Val Phe Ala Leu Gly Phe Leu Gly Cys Tyr Gly Ala Lys Thr
          65             70             75             80

Glu Ser Lys Cys Ala Leu Val Thr Phe Phe Phe Ile Leu Leu Leu Ile
          85             90             95

Phe Ile Ala Glu Val Ala Ala Ala Val Ala Leu Val Tyr Thr Thr
          100            105            110

Met Ala Glu His Phe Leu Thr Leu Leu Val Val Pro Ala Ile Lys Lys
          115            120            125

Asp Tyr Gly Ser Gln Glu Asp Phe Thr Gln Val Trp Asn Thr Thr Met
          130            135            140

Lys Gly Leu Lys Cys Cys Gly Phe Thr Asn Tyr Thr Asp Phe Glu Asp
          145            150            155            160

Ser Pro Tyr Phe Lys Glu Asn Ser Ala Phe Pro Pro Phe Cys Cys Asn
          165            170            175

Asp Asn Val Thr Asn Thr Ala Asn Glu Thr Cys Thr Lys Gln Lys Ala
          180            185            190

His Asp Gln Lys Val Glu Gly Cys Phe Asn Gln Leu Leu Tyr Asp Ile
          195            200            205

Arg Thr Asn Ala Val Thr Val Gly Gly Val Ala Ala Gly Ile Gly Gly
          210            215            220

```

Leu Glu Leu Ala Ala Met Ile Val Ser Met Tyr Leu Tyr Cys Asn Leu
 225 230 235 240

Gln

<210> 36
 <211> 377
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (346)

<400> 36
 caacctcgtg gtcaccgcac cggggctgat caaggggtgac gcctgcttca catctctaat 60
 gaacaccctc atgacgtcgc taccagcact agtgcagcaa caggggaaggc tgcttctggc 120
 tgctaatgtg gccaccctgg ggctcctcat ggcccggctc cttagcacct ctccagctct 180
 tcagggaaca ccagcatccc gagggttctt cgcagctgcc atcctcttcc taccacagtc 240
 ccacgtggcg cgggccaccc cgggctcaga ccaggcagtg ctagccctgt cccctgagta 300
 tgagggcatc tgggccgacc tgcaggagct ctgggttcctg ggcatncaag ccttcaccgg 360
 ctgtgtgcct ctgctgc 377

<210> 37
 <211> 106
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (96)

<400> 37
 Met Asn Thr Leu Met Thr Ser Leu Pro Ala Leu Val Gln Gln Gln Gly
 1 5 10 15
 Arg Leu Leu Leu Ala Ala Asn Val Ala Thr Leu Gly Leu Leu Met Ala
 20 25 30
 Arg Leu Leu Ser Thr Ser Pro Ala Leu Gln Gly Thr Pro Ala Ser Arg
 35 40 45
 Gly Phe Phe Ala Ala Ala Ile Leu Phe Leu Ser Gln Ser His Val Ala
 50 55 60
 Arg Ala Thr Pro Gly Ser Asp Gln Ala Val Leu Ala Leu Ser Pro Glu
 65 70 75 80
 Tyr Glu Gly Ile Trp Ala Asp Leu Gln Glu Leu Trp Phe Leu Gly Xaa
 85 90 95
 Gln Ala Phe Thr Gly Cys Val Pro Leu Leu
 100 105

<210> 38
 <211> 245

<212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (3)

<220>
 <221> unsure
 <222> (17)

<220>
 <221> unsure
 <222> (34)

<220>
 <221> unsure
 <222> (46)

<220>
 <221> unsure
 <222> (65)

<220>
 <221> unsure
 <222> (71)

<220>
 <221> unsure
 <222> (142)

<220>
 <221> unsure
 <222> (214)

<400> 38
 tangactcct ctctgcngag acgcgactgg cggntccagc agggantacc ttttttataa 60
 acccnggggg nccacacaca cacacacaca cacacacaca cacacacaca 120
 catttttgat cctttgcttc cntccccagc tgcgttctgt gatcgccaag ttcaaagctg 180
 tgcacatgtg gacactcaat aaatgttcat tggngacaaa aaaaaaaaaa aaaaaaaaaa 240
 aaaaa 245

<210> 39
 <211> 2384
 <212> DNA
 <213> Homo sapiens

<400> 39
 ccaaaacatg ctcaaaagta gaacattttg tttcaatatt aggaaagtgc tttgaatccc 60
 cttggacgac aaaagcgttg tctgagacag catgcgaaga ctcaagaggaa aacaagcaga 120
 gaataacagg tgcccagact ctaccaaagc atgtttctac cagcagtgat gaaggagacc 180
 ccagtgccag tacaccaatg atcaataaaa ctggctttta attttcagct gagaagcctg 240
 tgattgaagt tcccagcatg acaatcctgg ataaaaagga tggagagcag gccaaagccc 300
 tgtttgagaa agtgaggaag ttccgtgccc atgtggaaga tagtgacttg atctataaac 360
 tctatgtggt ccaaacagtt atcaaaacag ccaagtccat ttttattctc tgctatacag 420
 cgaactttgt caacgcaatc agctttgaac acgtctgcaa gcccaaagtt gagcatctga 480
 ttggttatga ggtatttgag tgcacccaca atatggctta catgttgaaa aagcttctca 540
 tcagttacat atccattatt tgtgtttatg gctttatctg cctctacact ctcttctggt 600
 tattcaggat acctttgaag gaatattctt tcgaaaaagt cagagaagag agcagtttta 660
 gtgacattcc agatgtcaaa aacgattttg cgttccttct tcacatggta gaccagtatg 720

```

accagctata ttccaagcgt tttggtgtgt tcttgtcaga agttagttaa aataaactta 780
gggaaattag tttgaaccat gagtggacat ttgaaaaact caggcagcac atttcacgca 840
acgcccagga caagcaggag ttgcatctgt tcatgctgtc gggggtgccc gatgctgtct 900
ttgacctcac agacctggat gtgctaaagc ttgaactaat tccagaagct aaaattcctg 960
ctaagatttc tcaaatgact aacctccaag agctccacct ctgccactgc cctgcaaaag 1020
ttgaacagac tgcttttagc tttcttcgcg atcacttgag atgccttcac gtgaagttca 1080
ctgatgtggc tgaatttcct gcctgggtgt atttgctcaa aaaccttcga gagttgtact 1140
taataggcaa tttgaactct gaaaacaata agatgatagg acttgaatct ctccgagagt 1200
tgcgccacct taagattctc cacgtgaaga gcaatttgac caaagttccc tccaacatta 1260
cagatgtggc tccacatctt acaaagttag tcattcataa tgacggcact aaactcttgg 1320
tactgaacag ccttaagaaa atgatgaatg tcgctgagct ggaactccag aactgtgagc 1380
tagagagaat cccacatgct attttcagcc tctctaattt acaggaactg gatttaaagt 1440
ccaataacat tcgcacaatt gaggaatca tcagtttcca gcatttaaaa cgactgactt 1500
gtttaaaatt atggcataac aaaattgtta ctattcctcc ctctattacc catgtcaaaa 1560
acttggagtc actttatttc tctaacaaca agctcgaatc cttaccagtg gcagtattta 1620
gtttacagaa actcagatgc ttagatgtga gctacaacaa catttcaatg attccaatag 1680
aataggatt gcttcagaac ctgcagcatt tgcatatcac tgggaacaaa gtggacattc 1740
tgccaaaaca attgtttaaa tgcataaagt tgaggacttt gaatctggga cagaactgca 1800
tcacctcact ccagagaaaa gttggtcagc tctcccagct cactcagctg gagctgaagg 1860
ggaactgctt ggaccgcctg ccagcccagc tgggccagtg tcggatgctc aagaaaagcg 1920
ggcttgttgt ggaagatcac ctttttgata ccctgccact cgaagtcaaa gaggcattga 1980
atcaagacat aatatattccc tttgcaaatg ggattttamac taagataata tatgcacagt 2040
gatgtgcagg aacaacttcc tagattgcaa gtgctcacgt acaagttatt acaagataat 2100
gcattttagg agtagataca tcttttataa taaaacagag aggatgcata gaaggctgat 2160
agaagacata actgaatggt caatgtttgt aggggtttta gtcattcatt tccaaatcat 2220
tttttttttt cttttgggga aagggaagga aaaattataa tcactaatct tgggttcttt 2280
taaattgttt gtaacttggg tgctgccgct actgaatggt tacaaattgc ttgcctgcta 2340
aagtaaatga ttaaattgac attttcttac tataaaaaaa aaaa 2384

```

<210> 40

<211> 614

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (607)

<400> 40

```

Met Ile Asn Lys Thr Gly Phe Lys Phe Ser Ala Glu Lys Pro Val Ile
  1                      5                      10                      15
Glu Val Pro Ser Met Thr Ile Leu Asp Lys Lys Asp Gly Glu Gln Ala
                20                      25                      30
Lys Ala Leu Phe Glu Lys Val Arg Lys Phe Arg Ala His Val Glu Asp
                35                      40                      45
Ser Asp Leu Ile Tyr Lys Leu Tyr Val Val Gln Thr Val Ile Lys Thr
                50                      55                      60
Ala Lys Phe Ile Phe Ile Leu Cys Tyr Thr Ala Asn Phe Val Asn Ala
                65                      70                      75                      80
Ile Ser Phe Glu His Val Cys Lys Pro Lys Val Glu His Leu Ile Gly
                85                      90                      95
Tyr Glu Val Phe Glu Cys Thr His Asn Met Ala Tyr Met Leu Lys Lys
                100                     105                     110

```

Leu Leu Ile Ser Tyr Ile Ser Ile Ile Cys Val Tyr Gly Phe Ile Cys
 115 120 125
 Leu Tyr Thr Leu Phe Trp Leu Phe Arg Ile Pro Leu Lys Glu Tyr Ser
 130 135 140
 Phe Glu Lys Val Arg Glu Glu Ser Ser Phe Ser Asp Ile Pro Asp Val
 145 150 155 160
 Lys Asn Asp Phe Ala Phe Leu Leu His Met Val Asp Gln Tyr Asp Gln
 165 170 175
 Leu Tyr Ser Lys Arg Phe Gly Val Phe Leu Ser Glu Val Ser Glu Asn
 180 185 190
 Lys Leu Arg Glu Ile Ser Leu Asn His Glu Trp Thr Phe Glu Lys Leu
 195 200 205
 Arg Gln His Ile Ser Arg Asn Ala Gln Asp Lys Gln Glu Leu His Leu
 210 215 220
 Phe Met Leu Ser Gly Val Pro Asp Ala Val Phe Asp Leu Thr Asp Leu
 225 230 235 240
 Asp Val Leu Lys Leu Glu Leu Ile Pro Glu Ala Lys Ile Pro Ala Lys
 245 250 255
 Ile Ser Gln Met Thr Asn Leu Gln Glu Leu His Leu Cys His Cys Pro
 260 265 270
 Ala Lys Val Glu Gln Thr Ala Phe Ser Phe Leu Arg Asp His Leu Arg
 275 280 285
 Cys Leu His Val Lys Phe Thr Asp Val Ala Glu Ile Pro Ala Trp Val
 290 295 300
 Tyr Leu Leu Lys Asn Leu Arg Glu Leu Tyr Leu Ile Gly Asn Leu Asn
 305 310 315 320
 Ser Glu Asn Asn Lys Met Ile Gly Leu Glu Ser Leu Arg Glu Leu Arg
 325 330 335
 His Leu Lys Ile Leu His Val Lys Ser Asn Leu Thr Lys Val Pro Ser
 340 345 350
 Asn Ile Thr Asp Val Ala Pro His Leu Thr Lys Leu Val Ile His Asn
 355 360 365
 Asp Gly Thr Lys Leu Leu Val Leu Asn Ser Leu Lys Lys Met Met Asn
 370 375 380
 Val Ala Glu Leu Glu Leu Gln Asn Cys Glu Leu Glu Arg Ile Pro His
 385 390 395 400
 Ala Ile Phe Ser Leu Ser Asn Leu Gln Glu Leu Asp Leu Lys Ser Asn
 405 410 415
 Asn Ile Arg Thr Ile Glu Glu Ile Ile Ser Phe Gln His Leu Lys Arg
 420 425 430

Leu Thr Cys Leu Lys Leu Trp His Asn Lys Ile Val Thr Ile Pro Pro
 435 440 445
 Ser Ile Thr His Val Lys Asn Leu Glu Ser Leu Tyr Phe Ser Asn Asn
 450 455 460
 Lys Leu Glu Ser Leu Pro Val Ala Val Phe Ser Leu Gln Lys Leu Arg
 465 470 475 480
 Cys Leu Asp Val Ser Tyr Asn Asn Ile Ser Met Ile Pro Ile Glu Ile
 485 490 495
 Gly Leu Leu Gln Asn Leu Gln His Leu His Ile Thr Gly Asn Lys Val
 500 505 510
 Asp Ile Leu Pro Lys Gln Leu Phe Lys Cys Ile Lys Leu Arg Thr Leu
 515 520 525
 Asn Leu Gly Gln Asn Cys Ile Thr Ser Leu Pro Glu Lys Val Gly Gln
 530 535 540
 Leu Ser Gln Leu Thr Gln Leu Glu Leu Lys Gly Asn Cys Leu Asp Arg
 545 550 555 560
 Leu Pro Ala Gln Leu Gly Gln Cys Arg Met Leu Lys Lys Ser Gly Leu
 565 570 575
 Val Val Glu Asp His Leu Phe Asp Thr Leu Pro Leu Glu Val Lys Glu
 580 585 590
 Ala Leu Asn Gln Asp Ile Asn Ile Pro Phe Ala Asn Gly Ile Xaa Thr
 595 600 605
 Lys Ile Ile Tyr Ala Gln
 610

<210> 41
 <211> 2386
 <212> DNA
 <213> Homo sapiens

<400> 41
 ccaaaacatc aaaccctttt cttgtagcag cacaggattc tgagacagat tatgtcacia 60
 cagataattt aacaaagggtg actgaggaag tcgtggcaaa catgcctgaa ggcctgactc 120
 cagatttagt acaggaagca tgtgaaagt aattgaatga agttactggt acaaagattg 180
 cttatgaaac aaaaatggac ttggttcaaa catcagaagt tatgcaagag tcaactctatc 240
 ctgcagcaca gctttgcccc tcatttgaag agtcagaagc tactccttca ccagttttgc 300
 ctgacattgt tatggaagca ccattgaatt ctgcagttcc tagtgctggt gcttccgtga 360
 tacagcccag ctcatcacca ttagaagcct cttcagttaa ttatgaaagc ataaaacatg 420
 agcctgaaaa cccccacca tatgaagagg ccatgagtgt atcactaaaa aaagtatcag 480
 gaataaagga agaaattaaa gagcctgaaa atattaatgc agctcttcaa gaaacagaag 540
 ctccttatat atctattgca tgtgatttaa ttaaagaaac aaagctttct gctgaaccag 600
 ctccggattt ctctgattat tcagaaatgg caaaagttga acagccagtg cctgatcatt 660
 ctgagctagt tgaagattcc tcacctgatt ctgaaccagt tgacttattt agtgatgatt 720
 caatacctga cgttccacaa aaacaagatg aaactgtgat gcttggtgaa gaaagtctca 780
 ctgagacttc atttgagtca atgatagaat atgaaaataa gaaaaaactc agtgctttgc 840
 cacctgaggg aggaaagcca tatttggaat cttttaagct cagtttagat aacacaaaag 900
 ataccctgtt acctgatgaa gtttcaacat tgagcaaaaa ggagaaaatt cctttgcaga 960
 tggaggagct cagtactgca gtttattcaa atgatgactt atttatttct aaggaagcac 1020

```

agataagaga aactgaaacg ttttcagatt catctccaat tgaaattata gatgagttcc 1080
ctacattgat cagttctaaa actgattcat tttctaaaatt agccagggaa tatactgacc 1140
tagaagtatc ccacaaaagt gaaattgcta atgccccgga tggagctggg tcattgcctt 1200
gcacagaatt gccccatgac ctttctttga agaacataca acccaaagtt gaagagaaaa 1260
tcagtttctc agatgacttt tctaaaaatg ggtctgctac atcaaagggtg ctcttattgc 1320
ctccagatgt ttctgctttg gccactcaag cagagataga gagcatagtt aaacccaaag 1380
ttcttgtaaa agaagctgag aaaaaacttc cttccgatac agaaaaagag gacagatcac 1440
catctgctat attttcagca gagctgagta aaacttcagt tgttgacctc ctgtactgga 1500
gagacattaa gaagactgga gtggtgtttg gtgccagcct attcctgctg ctttcattga 1560
cagtattcag cattgtgagc gtaacagcct acattgcctt ggccctgctc tctgtgacca 1620
tcagcttttag gatatacaag ggtgtgatcc aagctatcca gaaatcagat gaaggccacc 1680
cattcaggga agttgctata tctgaggagt tgggttcagaa gtacagtaat tctgctcttg 1740
gtcatgtgaa ctgcacgata aaggaactca ggccgctctt cttagttgat gatttagttg 1800
attctctgaa gtttgacgtg ttgatgtggg tatttaccta tgttggtgcc ttgtttaatg 1860
gtctgacact actgattttg gctctcattt cactcttcag tgttcctggt atttatgaac 1920
ggcatcaggc acagatagat cattatctag gacttgcaaa taagaatggt aaagatgcta 1980
tggtctaaaat ccaagcaaaa atccctggat tgaagcgcaa agctgaatga aaacgcccaa 2040
aataattagt aggagtcat ctttaaaggg gatattcatt tgattatacg ggggagggtc 2100
agggaagaac gaaccttgac gttgcagtgc agtttcacag atcgttggtta gatctttatt 2160
tttagccatg cactgttggt aggaataatt acctgtcttg actgccatgt gttcatcatc 2220
ttaagtattg taagctgcta tgtagggatt taaaccgtaa tcatatcttt ttcctatctg 2280
aggcactggt ggaataaaaa acctgtatat ttactttgt tgcagatagt cttgccgcat 2340
cttggaagt tgcagagatg gtggagctag aaaaaaaaaa aaaaaa 2386

```

<210> 42

<211> 642

<212> PRT

<213> Homo sapiens

<400> 42

```

Met Pro Glu Gly Leu Thr Pro Asp Leu Val Gln Glu Ala Cys Glu Ser
  1              5              10              15

Glu Leu Asn Glu Val Thr Gly Thr Lys Ile Ala Tyr Glu Thr Lys Met
      20              25              30

Asp Leu Val Gln Thr Ser Glu Val Met Gln Glu Ser Leu Tyr Pro Ala
      35              40              45

Ala Gln Leu Cys Pro Ser Phe Glu Glu Ser Glu Ala Thr Pro Ser Pro
      50              55              60

Val Leu Pro Asp Ile Val Met Glu Ala Pro Leu Asn Ser Ala Val Pro
      65              70              75              80

Ser Ala Gly Ala Ser Val Ile Gln Pro Ser Ser Ser Pro Leu Glu Ala
      85              90              95

Ser Ser Val Asn Tyr Glu Ser Ile Lys His Glu Pro Glu Asn Pro Pro
      100             105             110

Pro Tyr Glu Glu Ala Met Ser Val Ser Leu Lys Lys Val Ser Gly Ile
      115             120             125

Lys Glu Glu Ile Lys Glu Pro Glu Asn Ile Asn Ala Ala Leu Gln Glu
      130             135             140

Thr Glu Ala Pro Tyr Ile Ser Ile Ala Cys Asp Leu Ile Lys Glu Thr
      145             150             155             160

```

Lys Leu Ser Ala Glu Pro Ala Pro Asp Phe Ser Asp Tyr Ser Glu Met
 165 170 175
 Ala Lys Val Glu Gln Pro Val Pro Asp His Ser Glu Leu Val Glu Asp
 180 185 190
 Ser Ser Pro Asp Ser Glu Pro Val Asp Leu Phe Ser Asp Asp Ser Ile
 195 200 205
 Pro Asp Val Pro Gln Lys Gln Asp Glu Thr Val Met Leu Val Lys Glu
 210 215 220
 Ser Leu Thr Glu Thr Ser Phe Glu Ser Met Ile Glu Tyr Glu Asn Lys
 225 230 235 240
 Glu Lys Leu Ser Ala Leu Pro Pro Glu Gly Gly Lys Pro Tyr Leu Glu
 245 250 255
 Ser Phe Lys Leu Ser Leu Asp Asn Thr Lys Asp Thr Leu Leu Pro Asp
 260 265 270
 Glu Val Ser Thr Leu Ser Lys Lys Glu Lys Ile Pro Leu Gln Met Glu
 275 280 285
 Glu Leu Ser Thr Ala Val Tyr Ser Asn Asp Asp Leu Phe Ile Ser Lys
 290 295 300
 Glu Ala Gln Ile Arg Glu Thr Glu Thr Phe Ser Asp Ser Ser Pro Ile
 305 310 315 320
 Glu Ile Ile Asp Glu Phe Pro Thr Leu Ile Ser Ser Lys Thr Asp Ser
 325 330 335
 Phe Ser Lys Leu Ala Arg Glu Tyr Thr Asp Leu Glu Val Ser His Lys
 340 345 350
 Ser Glu Ile Ala Asn Ala Pro Asp Gly Ala Gly Ser Leu Pro Cys Thr
 355 360 365
 Glu Leu Pro His Asp Leu Ser Leu Lys Asn Ile Gln Pro Lys Val Glu
 370 375 380
 Glu Lys Ile Ser Phe Ser Asp Asp Phe Ser Lys Asn Gly Ser Ala Thr
 385 390 395 400
 Ser Lys Val Leu Leu Leu Pro Pro Asp Val Ser Ala Leu Ala Thr Gln
 405 410 415
 Ala Glu Ile Glu Ser Ile Val Lys Pro Lys Val Leu Val Lys Glu Ala
 420 425 430
 Glu Lys Lys Leu Pro Ser Asp Thr Glu Lys Glu Asp Arg Ser Pro Ser
 435 440 445
 Ala Ile Phe Ser Ala Glu Leu Ser Lys Thr Ser Val Val Asp Leu Leu
 450 455 460
 Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser Leu
 465 470 475 480

Phe Leu Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr Ala
 485 490 495
 Tyr Ile Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr
 500 505 510
 Lys Gly Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro Phe
 515 520 525
 Arg Glu Val Ala Ile Ser Glu Leu Val Gln Lys Tyr Ser Asn Ser
 530 535 540
 Ala Leu Gly His Val Asn Cys Thr Ile Lys Glu Leu Arg Arg Leu Phe
 545 550 555 560
 Leu Val Asp Asp Leu Val Asp Ser Leu Lys Phe Ala Val Leu Met Trp
 565 570 575
 Val Phe Thr Tyr Val Gly Ala Leu Phe Asn Gly Leu Thr Leu Leu Ile
 580 585 590
 Leu Ala Leu Ile Ser Leu Phe Ser Val Pro Val Ile Tyr Glu Arg His
 595 600 605
 Gln Ala Gln Ile Asp His Tyr Leu Gly Leu Ala Asn Lys Asn Val Lys
 610 615 620
 Asp Ala Met Ala Lys Ile Gln Ala Lys Ile Pro Gly Leu Lys Arg Lys
 625 630 635 640
 Ala Glu

<210> 43
 <211> 344
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (13)

<220>
 <221> unsure
 <222> (39)

<220>
 <221> unsure
 <222> (185)

<220>
 <221> unsure
 <222> (260)

<400> 43
 ggcggctgcg ganccggcgg tccttgcgct ccccaacanc ggcgccgjgg gcgcgggggc 60
 gccgtcgggc acagtcccgg tgctcttctg tttctcagtc ttcgcgcgac cctcgctcggt 120
 gccacacggg gcgggctaca agctgctcat ccagaagttc ctcagcctgt acggcgacca 180
 gatcnacatg caccgcaaatt tcgtggtgca gctgttcgcc gaggagtggg gccagtacgt 240

ggacttgccc aagggcttcn cggtagcgga gcgctgcaag gtgcgcctcg tgccgctgca 300
tatccagctc actaccctgg gaaatcttac accttcaagc actg 344

<210> 44
<211> 631
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> (73)

<220>
<221> unsure
<222> (369)

<400> 44
acatgattca gaggctctgt ggtaaattca tatgcaataa tcttattcca atcaatctgt 60
aaagtaaaaag cantacatcc acattaacat tataacatct tacagtaata taaaagccaa 120
atcattgttg gtacgtcatt ttctttaaaag tgaacaatct aagaaaactt cacaagagtc 180
tgcaacttttg aaagatacga tcagagtaca cagtagagac aaaacaggca tcttcattgt 240
aatttttttt aataaataaa agcacattaa caaaaaagga aggtaagcag caccggaagc 300
ctttgacgtt tgtaactaaa tgctggtact caattgaatc gagctgggta agtttacta 360
ggaggcgcn aaaaaggagcc gtttttgact taacatttta attctagtag agataagaag 420
agcttggttg ggcttacagt ccttcacctg actgtccttc accagttagt agcataccag 480
ttcttcaaat gtctataact ttggaaaagca gacccgactc tggagcactc gccttaatta 540
gattctgaat ttccttgaat tttggatggt ccttatcage taccagctga agcagaacag 600
cctcactcgt ggtcactatg atcccgggtc g 631

<210> 45
<211> 22
<212> PRT
<213> Homo sapiens

<400> 45
Met Val Leu Ile Ser Tyr Gln Leu Lys Gln Asn Ser Leu Thr Arg Gly
1 5 10 15

His Tyr Asp Pro Gly Ser
20

<210> 46
<211> 70
<212> DNA
<213> Homo sapiens

<400> 46
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaaa 70

<210> 47
<211> 428
<212> DNA
<213> Homo sapiens

<400> 47
agcacgcggt cctgcccgtg gacggggcaa cgctggcaga tgtgatgcgc cagcggggca 60
tcaacatgcg ctacctgggc aagggtgctg agctggtgct gcggarcccg gcccgccacc 120
agctggacca cgtctttaa atcggcattg gagaactcat caccgctcg sccaagcaca 180

tcttcaagac gtacttacag ggagtcgagc tctccggcct ctcagccgcc atcagccact 240
 tcctgaactg cttcctgagc tctacccaa acccctgtggc ccacctgccc gccgacgagc 300
 tgggtctccaa gaagcggat aagaggagga aaaaccggcc cccgggggct gcagataaca 360
 cagcctgggc tgtcatgacc ccccaggagc tctggaagaa catctgccag gaggccaaga 420
 actacttt 428

<210> 48

<211> 128

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (21)

<220>

<221> UNSURE

<222> (43)

<400> 48

Met Arg Gln Arg Gly Ile Asn Met Arg Tyr Leu Gly Lys Val Leu Glu
 1 5 10 15

Leu Val Leu Arg Xaa Pro Ala Arg His Gln Leu Asp His Val Phe Lys
 20 25 30

Ile Gly Ile Gly Glu Leu Ile Thr Arg Ser Xaa Lys His Ile Phe Lys
 35 40 45

Thr Tyr Leu Gln Gly Val Glu Leu Ser Gly Leu Ser Ala Ala Ile Ser
 50 55 60

His Phe Leu Asn Cys Phe Leu Ser Ser Tyr Pro Asn Pro Val Ala His
 65 70 75 80

Leu Pro Ala Asp Glu Leu Val Ser Lys Lys Arg Asn Lys Arg Arg Lys
 85 90 95

Asn Arg Pro Pro Gly Ala Ala Asp Asn Thr Ala Trp Ala Val Met Thr
 100 105 110

Pro Gln Glu Leu Trp Lys Asn Ile Cys Gln Glu Ala Lys Asn Tyr Phe
 115 120 125

<210> 49

<211> 245

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (46)

<220>

<221> unsure

<222> (138)

<220>

<221> unsure

<222> (147)

<400> 49

tggtggggga ggatgtgccc accctgagac cgggaggaga cgggcntttg cctgggtttg 60
 cggagagccg cttatgggtg tgggccgtcc agacaccttg tttcaagggg gatgggcgtg 120
 agcgggcaag cagagcanc caccgntga gcaagaactt tttttgttt ttaaaccatc 180
 acgtcctcat ttcacattgg aataaagtga gtttttgaaa aaaaaaaaaa aaaaaaaaaa 240
 aaaaaa 245

<210> 50

<211> 566

<212> DNA

<213> Homo sapiens

<400> 50

cagtgaagccc tttgaaaaat aaacatccag atgaagatgc tgtggaagct gaggggcatg 60
 aggtaaaaag actcaggttt gacaaagaag gtgaagtcag agaaacagcc agtcaaacga 120
 cttccagcga aattttcttca gttatggtag gagaaacaga agcatcatct tcatctcagg 180
 ataaagacaa agatagccgt tgtwcccgcc agcactgtwc agaagaggat gaagaagagg 240
 atgaagagga agaagaagag tcttttatga catcaagaga aatgatccca gaaagaaaaa 300
 atcaagaaaa agaattctgat gatgccttaa ctgtgaatga agagacttct gaggaaaata 360
 atcaaatgga ggaattctgat gtgtctcaag ctgagaaaga tttgctacat tctgaaggta 420
 gtgaaaacga aggcctctgta agtagtagtt cttctgactg ccgtgaaaca gaagaattag 480
 taggatccaa ttccagtaaa actggagaga ttctttcaga atcatccatg gaaaatgatg 540
 acgaagccac agaagtcacc gatgaa 566

<210> 51

<211> 141

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (21)

<220>

<221> UNSURE

<222> (26)

<400> 51

Met Val Gly Glu Thr Glu Ala Ser Ser Ser Ser Gln Asp Lys Asp Lys
 1 5 10 15
 Asp Ser Arg Cys Xaa Arg Gln His Cys Xaa Glu Glu Asp Glu Glu Glu
 20 25 30
 Asp Glu Glu Glu Glu Glu Ser Phe Met Thr Ser Arg Glu Met Ile
 35 40 45
 Pro Glu Arg Lys Asn Gln Glu Lys Glu Ser Asp Asp Ala Leu Thr Val
 50 55 60
 Asn Glu Glu Thr Ser Glu Glu Asn Asn Gln Met Glu Glu Ser Asp Val
 65 70 75 80
 Ser Gln Ala Glu Lys Asp Leu Leu His Ser Glu Gly Ser Glu Asn Glu
 85 90 95
 Gly Pro Val Ser Ser Ser Ser Asp Cys Arg Glu Thr Glu Glu Leu
 100 105 110

Val Gly Ser Asn Ser Ser Lys Thr Gly Glu Ile Leu Ser Glu Ser Ser
 115 120 125

Met Glu Asn Asp Asp Glu Ala Thr Glu Val Thr Asp Glu
 130 135 140

<210> 52
 <211> 531
 <212> DNA
 <213> Homo sapiens

<400> 52
 tcatcatggc tataaataacc aaaacgattt ggatccattt atgtttgtag gataatatac 60
 tactgactga cttgactgtc aggttcacaa cagctagatg atatatattat gactatgtct 120
 aatagttgaa ataaaaatctg aatattgatt tactataccc aagaggggag aaaaattaac 180
 cattgtaaat ttttaaaaat tttttcaaaa atgttaaaat gaggcaaatt taagtttaca 240
 aattttgaaa ttttcttttg aatatttatg aaattgtcag taaacttacc taagatcctg 300
 tgaccttttg atatttttta ttttaattgt agtgccatgg accatttgta aacaaattga 360
 tttacttttg ttggttgtaa gttgaagatt tagcattatg actttgaggt ctgtgggttt 420
 atttgtaaac ttgcaattgc tatatttgca agggcaaattg tatttcttta ttaaataaag 480
 tacaataatg gtgaatgtac caaaatgaca tcacttaaaa aaaaaaaaaa a 531

<210> 53
 <211> 1163
 <212> DNA
 <213> Homo sapiens

<400> 53
 kctggaacca cgcggargaa ggaagagacg caggcaggct gcggttacc aagcggscac 60
 ccgggacctca gggacccttc ccgagagac gccaccatga ccaggggaaa gctctccgtg 120
 gctaacaagc ccctgggacc gaggggcagc agcakgtgca tggcgagaag aaggagctcc 180
 agcagtgcctc tcagccccac cctcctatga ggaaccacct ctggggaggg gatgaaggca 240
 ggggaccttc cccagcccc cacagcggty cctctccacc ctactgggc ctatgtggac 300
 cccagcagca gtcacagcta tgacaacggt ttccccaccg gagaccatga gctcttcacc 360
 actttcagct gggatgacca gaaagtctgt cgagtctttg tcagaaaggc ctacaccatc 420
 ctgctgattc agctgctggg gacctgggt gtcgtggctc tctttacttt ctgtgacctc 480
 gtcaaggact atgtccaggc caaccaggc tggtagctggg catcctatgc tgtgttctct 540
 gcaacctacc tgacctggc ttgctgttct ggaccagga ggcatttccc ctggaacctg 600
 attctcctga ccgtctttac cctgtccatg gcctacctca ctgggatgct gtccagctac 660
 tacaacacca cctcctgct gctgtgcctg ggcacacagg cccttgctg cctctcagtc 720
 accgtcttca gcttccagac caagtctgac ttcacctcct gccagggcgt gctcttcgtg 780
 cttctcatga ctcttttctt cagcggactc atcctggcca tcctcctacc cttccaatat 840
 gtgccctggc tccatgcagt ttatgcagca ctgggagcgg gtgtatttac attgttctctg 900
 gcacttgaca cccagtgtgt gatgggtaac cgacgccact cgctgagccc tgaggagtat 960
 atttttggag ccctcaacat ttacctagac atcatctata tcttcacctt ctctctgcag 1020
 ctttttggca ctaaccgaga atgaggagcc ctccctgccc caccgtcctc cagagaatgc 1080
 gcccctcctg gttccctgtc cctccctgct gctcctgcga gaccagatat aaaactagct 1140
 gccaacccaa aaaaaaaaaa aaa 1163

<210> 54
 <211> 270
 <212> PRT
 <213> Homo sapiens

<400> 54
 Met Lys Ala Gly Ala Phe Pro Pro Ala Pro Thr Ala Val Pro Leu His
 1 5 10 15

Pro Ser Trp Ala Tyr Val Asp Pro Ser Ser Ser Ser Ser Tyr Asp Asn
 20 25 30
 Gly Phe Pro Thr Gly Asp His Glu Leu Phe Thr Thr Phe Ser Trp Asp
 35 40 45
 Asp Gln Lys Val Arg Arg Val Phe Val Arg Lys Val Tyr Thr Ile Leu
 50 55 60
 Leu Ile Gln Leu Leu Val Thr Leu Ala Val Val Ala Leu Phe Thr Phe
 65 70 75 80
 Cys Asp Pro Val Lys Asp Tyr Val Gln Ala Asn Pro Gly Trp Tyr Trp
 85 90 95
 Ala Ser Tyr Ala Val Phe Phe Ala Thr Tyr Leu Thr Leu Ala Cys Cys
 100 105 110
 Ser Gly Pro Arg Arg His Phe Pro Trp Asn Leu Ile Leu Leu Thr Val
 115 120 125
 Phe Thr Leu Ser Met Ala Tyr Leu Thr Gly Met Leu Ser Ser Tyr Tyr
 130 135 140
 Asn Thr Thr Ser Val Leu Leu Cys Leu Gly Ile Thr Ala Leu Val Cys
 145 150 155 160
 Leu Ser Val Thr Val Phe Ser Phe Gln Thr Lys Phe Asp Phe Thr Ser
 165 170 175
 Cys Gln Gly Val Leu Phe Val Leu Leu Met Thr Leu Phe Phe Ser Gly
 180 185 190
 Leu Ile Leu Ala Ile Leu Leu Pro Phe Gln Tyr Val Pro Trp Leu His
 195 200 205
 Ala Val Tyr Ala Ala Leu Gly Ala Gly Val Phe Thr Leu Phe Leu Ala
 210 215 220
 Leu Asp Thr Gln Leu Leu Met Gly Asn Arg Arg His Ser Leu Ser Pro
 225 230 235 240
 Glu Glu Tyr Ile Phe Gly Ala Leu Asn Ile Tyr Leu Asp Ile Ile Tyr
 245 250 255
 Ile Phe Thr Phe Phe Leu Gln Leu Phe Gly Thr Asn Arg Glu
 260 265 270

<210> 55

<211> 624

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (123)

<400> 55

cgcaccccct ccggccgcgg gcgcakcggg ggcgcctgggtg gacctgmgaaggccasgtc 60

```

cgggcgggcg ggcgggcggt ggcactggct cgggactctg cccggccagg gcggcggmte 120
cancggggag ggcgacgtgg agcgggccack tggakcggcc cggggggargc tggcgggcggg 180
akgcgagggcg cggggcgggcg akcakccakg agcgcccacg gagstggacc cccagakccg 240
cgcgcgcgccg cagcagttcc aggaaggatg ttacctttga cgatgacagt gttaatcctg 300
ctgctgctcc ccacgggtca ggctgcccc aaggatggag tcacaaggcc agaattctgaa 360
gtgcagcatc agctcctgcc caacccttc cagccaggcc aggagcagct cggacttctg 420
cagagctacc taaaggggact aggaaggaca gaagtgcac tggagcatct gagccgggag 480
caggttctcc tctacctctt tgccctccat gactatgacc agagtggaca gctggatggc 540
ctggagctgc tgtccatggt gacagctgct ctggcccctg gagctgccaa ctctcctacc 600
accaaccggg tgatcttgat agtg                                     624

```

<210> 56
 <211> 119
 <212> PRT
 <213> Homo sapiens

<400> 56
 Met Leu Pro Leu Thr Met Thr Val Leu Ile Leu Leu Leu Pro Thr
 1 5 10 15
 Gly Gln Ala Ala Pro Lys Asp Gly Val Thr Arg Pro Glu Ser Glu Val
 20 25 30
 Gln His Gln Leu Leu Pro Asn Pro Phe Gln Pro Gly Gln Glu Gln Leu
 35 40 45
 Gly Leu Leu Gln Ser Tyr Leu Lys Gly Leu Gly Arg Thr Glu Val Gln
 50 55 60
 Leu Glu His Leu Ser Arg Glu Gln Val Leu Leu Tyr Leu Phe Ala Leu
 65 70 75 80
 His Asp Tyr Asp Gln Ser Gly Gln Leu Asp Gly Leu Glu Leu Leu Ser
 85 90 95
 Met Leu Thr Ala Ala Leu Ala Pro Gly Ala Ala Asn Ser Pro Thr Thr
 100 105 110
 Asn Pro Val Ile Leu Ile Val
 115

<210> 57
 <211> 80
 <212> DNA
 <213> Homo sapiens

<400> 57
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa 80

<210> 58
 <211> 2160
 <212> DNA
 <213> Homo sapiens

<400> 58
 agacagggaa tactttatcc aaaacccatc acagaaatgg acagcttggg tctgtaacaa 60
 agcattcatg ttttagagca taggtcagta attgtatatg agagcataca ctgctacata 120
 caaattaact gatcagacca caacttttca atgttttaaaa cagaataagc ttccctgtaa 180

```

aagcagcacc tttgtgacgt ttttaacttta gtatttctct ccttcttctt caccctctcc 240
ttcaacagaa tccacaccaa cctcctcata atccttctct gcagcacatg aatcacaggt 300
attcctactg caagcgggag gcgaggagc ggggaagcggc ggagcgcgag gcgcgcgaga 360
aagggcactt ggaacccacc gagctgctga tgaaccgggc ttacttgagc agcattaccc 420
ctcaggggta ccttgactcg gaggagagg agagtatgcc gagggatggc gagagcgaga 480
aggagcacga gaaagaaggc gaggatggct acgggaagct gggcagacag gatggcgacg 540
aggagtctga ggaggaagag gaagaaagtg aaaataaaag tatggatacg gatcccgaaa 600
cgatacgaga tgaagaagag actggagatc actccatgga cgatagtctg gaggatggga 660
aatggaaac caaatcagac cagcaggaag acaatatgga agatggcatg taataaacta 720
ctgcatttta agcttctctat ttttttttcc agtagtattg ttacctgctt gaaaacactg 780
ctgtgttaag ctgttcatgc acgtgcctga cgcttcagg aagctgtaga gagggacaga 840
agggcggtt cagccaagac agatgtagac ggagtggag ctgggtattg ttaaaaactg 900
cattatgcaa aaattttgta cagtgttaag gcctaaaaac tgtgtggttc agagactaat 960
tcctgtgttt aatagcattt atactttaag cacaactaga aaattgtgag aattgcactc 1020
tacttatgta tcaactacaa ctttaaaaaa ctatgtctaa tttatattaa tacattttta 1080
aaaggtgccc gcactaccat acatcagtat ttttattatt attattgtta ttctttttta 1140
atttaatgtg ctgcactac aatgcacag tattatgatt cctctgtact ttcttttctg 1200
tattcatcaa ttctccattt tttttttcag cttaagtaac cacacaattt taggcctcaa 1260
tttttttttt tctgtgaagg aacttgaagt gatgcatgtg tgaatttaag ataccgaagt 1320
cttaagtgga cctggacgtg aaggaaaaag taagatgaga aataaagaaa gcctttgtta 1380
ggtgtgttta aaagccttat atgcaaacct tttaatctgt gtttctgcaa gtgccatcct 1440
tgtacagtgt taagagggtg acatgggtta cctttgcacc agcttcagtg ttaagctcac 1500
cctgttcttt gaagcaccca tgtcagtatt agaagaatag gcagcagttc cttagtttac 1560
atatgtttgt gcaattattt tctgtacttt tttgttcatt aattttgtca gtattacacc 1620
aaactgtttt tgcaacaaaa aaattttttt tgcattcatt taattttagg tcaaataaca 1680
ttttatttat gtggctcatt ttatatattc taattttatt tatttcatac tgtagtgtac 1740
agtattatag ttcttcaata tatagatata ttttagtaaa aaaggaacat gacgttgatc 1800
atttgggcaa attttacgta aagagaagag catttattgt gttttggaac attaatgtg 1860
agatgggatt ttcaattttt attattttat tttgtttttt ttccaattac tggaaattcc 1920
aaatttggga acttttgata cgatcttggt aaaacactgt attttcgact gaaaattcca 1980
ctttcttcat cttgtttttt agctaaaaag agggactgtt aaatacaatg tatgatacca 2040
tgacaaaaat ctttcttgaa ttgtctttgt aaaagtatta ttgaattttc aatttgaat 2100
ttcttttgaa aatgaccatg ctggaataaa aatgtagcca aactaaaaaa aaaaaaaaaa 2160

```

<210> 59

<211> 141

<212> PRT

<213> Homo sapiens

<400> 59

```

Met Asn His Arg Tyr Ser Tyr Cys Lys Arg Glu Ala Glu Glu Arg Glu
 1              5              10              15
Ala Ala Glu Arg Glu Ala Arg Glu Lys Gly His Leu Glu Pro Thr Glu
 20              25              30
Leu Leu Met Asn Arg Ala Tyr Leu Gln Ser Ile Thr Pro Gln Gly Tyr
 35              40              45
Ser Asp Ser Glu Glu Arg Glu Ser Met Pro Arg Asp Gly Glu Ser Glu
 50              55              60
Lys Glu His Glu Lys Glu Gly Glu Asp Gly Tyr Gly Lys Leu Gly Arg
 65              70              75              80
Gln Asp Gly Asp Glu Glu Phe Glu Glu Glu Glu Glu Ser Glu Asn
 85              90              95
Lys Ser Met Asp Thr Asp Pro Glu Thr Ile Arg Asp Glu Glu Glu Thr
100              105              110

```


Gly Asp His Ser Met Asp Asp Ser Ser Glu Asp Gly Lys Met Glu Thr
 115 120 125

Lys Ser Asp His Glu Glu Asp Asn Met Glu Asp Gly Met
 130 135 140

<210> 60
 <211> 2168
 <212> DNA
 <213> Homo sapiens

<400> 60
 gcagttactg ggagggggct tgctgtggcc ctgtcaggaa gagtagagct ctggtccagc 60
 tccgcgcagg gagggaggct gtcaccatgc cggcctgctg cagctgcagt gatgttttcc 120
 agtatgagac gaacaaagtc actcggatcc agagcatgaa ttatggcacc attaagtggt 180
 tcttccacgt gatcatcttt tcttacgttt gctttgctct ggtgagtgac aagctgtacc 240
 agcggaaaga gcctgtcatc agttctgtgc acaccaaggt gaaggggata gcagagggtga 300
 aagaggagat cgtggagaat ggagtgaaga agttgggtgca cagtgtcttt gacaccgcag 360
 actacacctt ccctttgcag gggaactctt tcttcgtgat gacaaacttt ctcaaaacag 420
 aaggccaaga gcagcggttg tgtcccgagt atcccaccgc caggacgctc tgttcctctg 480
 accgaggttg taaaaaggga tggatggacc cgcagagcaa aggaattcag accggaaggt 540
 gtgtagtgc tgaagggaac cagaagacct gtgaagtctc tgcctgggtgc cccatcgagg 600
 cagtggaaaga ggcccccccg cctgctctct tgaacagtgc cgaaaacttc actgtgctca 660
 tcaagaacaa tatcgacttc cccggccaca actacaccac gagaacatc ctgccagggt 720
 taaacatcac ttgtaccttc cacaagactc agaattccaca gtgtcccatc ttccgactag 780
 gagacatctt ccgagaaaca ggcgataatt tttcagatgt ggcaattcag ggcggaataa 840
 tgggcattga gatctactgg gactgcaacc tagaccgttg gttccatcac tgccatccca 900
 aatacagttt ccgtcgcttc gacgacaaga ccaccaacgt gtccttgtag cctggctaca 960
 acttcagata cgccaagtac tacaaggaaa acaatgttga gaaacggact ctgataaaag 1020
 tcttcgggat ccgttttgac atcctggttt ttggcaccgc aggaaaattt gacattatcc 1080
 agctgggtgt gtacatcggc tcaacctctt cctacttcgc tctggccgct gtgttcacg 1140
 acttctctcat cgacacttac tccagtaact gctgtcgcct ccatatttat ccctgggtgca 1200
 agtgctgtca gccctgtgtg gtcaacgaat actactacag gaagaagtgc gagtccattg 1260
 tggagccaaa gccgacatta aagtatgtgt cctttgtgga tgaatccac attaggatgg 1320
 tgaaccagca gctactaggg agaagtctgc aagatgtcaa gggccaagaa gtcccaagac 1380
 ctgcgatgga cttcacagat ttgtccaggc tgccccctggc cctccatgac acacccccga 1440
 ttcttgagca accagaggag atacagctgc ttagaaaagga ggcgactcct agatccaggg 1500
 atagccccgt ctggtgccag tgtggaagct gcctcccatc tcaactccct gagagccaca 1560
 ggtgcctgga ggagctgtgc tgccggaaaa agccgggggc ctgcatcacc acctcagagc 1620
 tgttcaggaa gctggtcctg tccagacacg tccctgcagt cctcctgctc taccaggagc 1680
 ccttgctggc gctggatgtg gattccacca acagccggct gcggcactgt gcctacaggt 1740
 gctacgccac ctggcgcttc ggctcccagg acatggctga ctttgccatc ctgccagct 1800
 gctgcccgtg gaggatccgg aaagagtttc cgaagagtga agggcagtac agtggcttca 1860
 agagtccctta ctgaagccag gcgccgtggc tcacgtctgt aatcccagcg ctttgggagg 1920
 ccgaggcagg cagatcacct gaggtcggga gttggagacc cgcttggtca acaaggcgaa 1980
 atcctgtctg tactaaaaat acaaaaatca gccagacatg gtggcatgca cctgcaatcc 2040
 cagctactcg ggaggctgag gcacaagaat cacttgaacc cgggaggcag aggttgtagt 2100
 gagcccagat tgtgccactg ctctccagcc tgggaggcac agcaaactgt ccccaaaaaa 2160
 aaaaaaaa 2168

<210> 61
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> 61
 Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30
 Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45
 Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60
 Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80
 Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95
 Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110
 Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125
 Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140
 Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160
 Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175
 Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
 180 185 190
 Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205
 Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
 210 215 220
 Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240
 Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255
 Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys His Pro Lys
 260 265 270
 Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285
 Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300
 Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320
 Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350
 Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365
 Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380
 Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400
 Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415
 Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430
 Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445
 Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460
 Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480
 Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495
 Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510
 Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525
 Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540
 Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560
 Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575
 Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590
 Ser Pro Tyr
 595

<210> 62

<211> 430

<212> DNA

<213> Homo sapiens

<400> 62

taaagatctg tggttcagagt cataactgaay agagacttct ggactctata gaaccactg 60

cctcctgatg aagtccttac tgttcaccct tgcagttttt atgctcctgg cccaattggg 120
 ctcaggtaat tggatatgta aaaagtgtct aaacgacgtt ggaatttgca agaagaagtg 180
 caaacctgaa gagatgcatg taaagaatgg ttgggcaatg tgcggcaaac aaagggactg 240
 ctgtgttcca gctgacagac gtgctaatta tcctgttttc tgtgtccaga caaagactac 300
 aagaatttca acagtaacag caacaacagc aacaacaact ttgatgatga ctactgcttc 360
 gatgtcttcg atggctccta cccgtttctc cactgggtg aacattccag cctctgtctc 420
 ctgctctagg 430

<210> 63

<211> 121

<212> PRT

<213> Homo sapiens

<400> 63

Met Lys Ser Leu Leu Phe Thr Leu Ala Val Phe Met Leu Leu Ala Gln
 1 5 10 15

Leu Val Ser Gly Asn Trp Tyr Val Lys Lys Cys Leu Asn Asp Val Gly
 20 25 30

Ile Cys Lys Lys Lys Cys Lys Pro Glu Glu Met His Val Lys Asn Gly
 35 40 45

Trp Ala Met Cys Gly Lys Gln Arg Asp Cys Cys Val Pro Ala Asp Arg
 50 55 60

Arg Ala Asn Tyr Pro Val Phe Cys Val Gln Thr Lys Thr Thr Arg Ile
 65 70 75 80

Ser Thr Val Thr Ala Thr Thr Ala Thr Thr Leu Met Met Thr Thr
 85 90 95

Ala Ser Met Ser Ser Met Ala Pro Thr Arg Phe Ser His Trp Leu Asn
 100 105 110

Ile Pro Ala Ser Val Ser Cys Ser Arg
 115 120

<210> 64

<211> 112

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (8)

<220>

<221> unsure

<222> (12)

<220>

<221> unsure

<222> (36)

<220>

<221> unsure

<222> (41)

<220>
 <221> unsure
 <222> (44)

<400> 64
 tttcctgntt tnggatecccc gattcattaa agcaangggg nttnaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 112

<210> 65
 <211> 324
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (1)

<220>
 <221> unsure
 <222> (69)

<220>
 <221> unsure
 <222> (74)

<220>
 <221> unsure
 <222> (125)

<220>
 <221> unsure
 <222> (159)

<400> 65
 nctaccccaa cctgtgtggc tgggccgagg tctccctca agggcctggg gccgtgcctc 60
 ggggtgtacnc gtanggggtct gtgtgctggg ggtggctcac cgggcagcgt gggtgagcgg 120
 cgcancggcg gcagcggaga acgagagagg ggagcagana cagaatcgcc taagctgaag 180
 tgtattggcg ccatcatggc tcaactgcgc ctccggctcc ttggctcggg tgattctcct 240
 gcctgagcct ccctagtagc taggactaca gtgctgtaga agaaaatcac atgattgggtg 300
 ccctcaaaaa attggtgcca cttg 324

<210> 66
 <211> 794
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (61)

<220>
 <221> unsure
 <222> (82)

<220>
 <221> unsure
 <222> (108)..(120)

<220>
 <221> unsure

<222> (184)

<220>

<221> unsure

<222> (754)

<400> 66

cattatttca tcaccagaga atacacatgc agcaaatagc attgtgagtc aaactattcc 60
naaagcacag attcagcaat cnacacacac tcattctggat atctcacnnn nnnnnnnnnn 120
ttaactgatg aaaaaagtaa tggaacaatt gcccttgtgg atgattctga ggatcctgga 180
gccnatgtat ctaacataca gcttcagcaa aaaatttcaa gtctggagat taaactcaaa 240
gtatctgaag aagaaaaaca gagaattaaa caggatgtgg aakcattgat ggaaaagcat 300
aatgtcttag aaaaaggctt tctaaaagaa aaagagcaag aggccatttc ttttcaagat 360
agatacaaaag aacttcagga aaaacataaa caagaattgg aagacatgag gaaagctggt 420
cacgaagccc tcagcattat tgtggatgaa tataaggcac tactgcagtc ttcagttaag 480
caacaagtag aagctattga aaaacagtac atttctgcaa ttgagaaaca ggcacacaag 540
tgtgaggagt tgctaaatgc tcagcatcag aggctccttg aagtgcataa tacagagaag 600
gaactgttaa aagaaaaaat aaaggaagct ttgattcagc aatctcaaga acagaaggaa 660
atattggaaa agtgttttga ggaagaaagg caaagaaata aagaggcatt agtatccgct 720
gcaaagcttg aaaaagaacc agtgaaggat gcanttttaa aattcgtaga agaagaaaga 780
aaaaaaaaaa aaaa 794

<210> 67

<211> 164

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (156)

<400> 67

Met Glu Lys His Asn Val Leu Glu Lys Gly Phe Leu Lys Glu Lys Glu
1 5 10 15
Gln Glu Ala Ile Ser Phe Gln Asp Arg Tyr Lys Glu Leu Gln Glu Lys
20 25 30
His Lys Gln Glu Leu Glu Asp Met Arg Lys Ala Gly His Glu Ala Leu
35 40 45
Ser Ile Ile Val Asp Glu Tyr Lys Ala Leu Leu Gln Ser Ser Val Lys
50 55 60
Gln Gln Val Glu Ala Ile Glu Lys Gln Tyr Ile Ser Ala Ile Glu Lys
65 70 75 80
Gln Ala His Lys Cys Glu Glu Leu Leu Asn Ala Gln His Gln Arg Leu
85 90 95
Leu Glu Val Leu Asp Thr Glu Lys Glu Leu Leu Lys Glu Lys Ile Lys
100 105 110
Glu Ala Leu Ile Gln Gln Ser Gln Glu Gln Lys Glu Ile Leu Glu Lys
115 120 125
Cys Leu Glu Glu Glu Arg Gln Arg Asn Lys Glu Ala Leu Val Ser Ala
130 135 140
Ala Lys Leu Glu Lys Glu Pro Val Lys Asp Ala Xaa Leu Lys Phe Val

145

150

155

160

Glu Glu Glu Arg

<210> 68

<211> 1494

<212> DNA

<213> Homo sapiens

<400> 68

```

cgtttgtcga cgccgctgcc accgcctgcc tgagagaagt cgtcgcggcc gaccccgctcg 60
cctccgcccgg ctaccatgtc cgcccaggcg cagatgcggg cctgctgga ccagctcatg 120
ggcacgggctc gggacgggaga cgaaaccaga cagaggggtca agtttacaga tgaccgtgtc 180
tgcaagagtc accttctgga ctgctgcccc catgacatcc tggctgggac gcgcatggat 240
ttaggagaat gtacaaaaat ccacgacttg gccctccgag cagattatga gattgcaagt 300
aaagaaagag acctgttttt tgaattagat gcaatggatc acttgagtc ctttattgct 360
gaatgtgatk ggagaactga gctcgccaag aagcggcttg cagaaacaca ggaggaaatc 420
agtgcggaag tttctgcaaa ggcaggaaaa gtacatgagt taaatgaaga aataggaaaa 480
ctccttgcta aagccgaaca gctaggggct gaaggtaatg tggatgaatc ccagaagatt 540
cttatggaag tggaaaaagt tcgtgcgaag aaaaaagaag ctgaggaaga atacagaaat 600
tccatgcctg catccagttt tcagcagcaa aagctgcgtg tctgcgaggt ctgttcagcc 660
taccttggtc tccatgacaa tgaccgtcgc ctggcagacc acttcggtgg caagttacac 720
ttgggggttca ttcagatccg agagaagctt gatcagttga ggaaaactgt cgttgaaaag 780
caggagaaga gaaatcagga tcgcttgagg aggagagagg agagggaacg ggaggagcgt 840
ctgagcagga ggtcgggac aagaaccaga gatcgagga ggtcacgctc ccgggacggt 900
cgtcggaggc ggtcaagatc tacctcccga gagcgacgga aattgtcccg gtcccgggtc 960
cgagatagac atcggcgcca ccgcagccgt tcccggagcc acagccgggg acatcgtcgg 1020
gcttcccggg accgaagtgc gaaatacaag taactactct gactccttcg gtagctgcaa 1080
ccaggagttc tccagagagc gggcatccag agaggagtcc tgggagagcg ggcggagcga 1140
gcgagggccc ccgactgga ggcttgagag ctccaacggg aagatggctt cacggaggtc 1200
agaagagaag gaggccggcg agatctgaac ccgtctcccg ggtgctgtaa atagtctgat 1260
aaacgttcac acagtctaaa attacccttt atatttgctg aatacaactc atcttttgta 1320
gtttaaaatt tctattgttt tggagctagc tgtgagtttc tagaagtgta cagagttgct 1380
cctgtgttcc cgggtcatgt tgagtaggaa taaataaatc tgatgctgcc tcctggaaaa 1440
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 1494

```

<210> 69

<211> 325

<212> PRT

<213> Homo sapiens

<400> 69

```

Met Ser Ala Gln Ala Gln Met Arg Ala Leu Leu Asp Gln Leu Met Gly
 1             5             10            15

Thr Ala Arg Asp Gly Asp Glu Thr Arg Gln Arg Val Lys Phe Thr Asp
 20            25            30

Asp Arg Val Cys Lys Ser His Leu Leu Asp Cys Cys Pro His Asp Ile
 35            40            45

Leu Ala Gly Thr Arg Met Asp Leu Gly Glu Cys Thr Lys Ile His Asp
 50            55            60

Leu Ala Leu Arg Ala Asp Tyr Glu Ile Ala Ser Lys Glu Arg Asp Leu
 65            70            75            80

Phe Phe Glu Leu Asp Ala Met Asp His Leu Glu Ser Phe Ile Ala Glu

```

```
<210> 70
<211> 1761
<212> DNA
<213> Homo sapiens
```

<400> 70									
caaggggaagt	tctgagggct	gagaggttgc	tcattctgtca	gagcgtgctg	cccaccctcc	60			
acccctgcat	ggcagaaact	gtgcagggga	cgaggccaag	gaatcaggag	accagaggcc	120			
aggggtggcc	cggagacggt	gaagaaacca	agacgcagag	aggccaagcc	ccttgccctg	180			
ggtcacacag	ccaaaggagg	cagagccaga	actcacacc	agatccaagc	gcaacaggga	240			
catggccacc	tgggacgaaa	aggcagtcac	ccgcagggcc	aaggtggctc	ccgctgagag	300			
gatgagcaag	ttcttaaggc	acttcacggt	ctggggagac	gactaccatg	cctggaacat	360			


```

caactacaag aaatgggaga atgaagagga ggaggaggag gaggagcagc caccacccac 420
accagtctca ggcgaggaag gcagagctgc agcccttgac gttgcccctg cccctggccc 480
cgcacccagg gcccccttg acttcagggg catgttgagg aaactgttca gctcccacag 540
gtttcaggtc atcatcatct gcttggtggt tctggatgcc ctcttggtgc ttgctgagct 600
catcctggac ctgaagatca tccagcccga caagaataac tatgctgcca tggattacca 660
ctacatgagc atcaccatct tggctctttt tatgatggag atcatcttta aattatttgt 720
cttcgcctg gagttctttc accacaagtt tgagatcctg gatgccgtcg tgggtggtgt 780
ctcattcatc ctgcacattg tcctcctggt ccaggagcac cagtttgagg ctctgggcct 840
gctgattctg ctccggctgt ggcgggtggc ccggatcatc aatgggatta tcattcagc 900
taagacacgt tcagaacggc aactcttaag gttaaacag atgaatgtac aattggccgc 960
caagattcaa cactttgagt tcagctgctc tgagaaggaa caagaaattg aaagacttaa 1020
caactattg cgacagcatg gacttcttgg tgaagtgaac tagaccgga ccagctcccc 1080
tcaaaaagaa gacactgtct catgggcctg tgctgtcacg agaggaacag ctgcccctcc 1140
tgggcgcgtt ggtgagaggt ttggtttgat acctctgcct ccctcctgcc agcatggatt 1200
ctgggtggac acagccttgt ggaagggtcca gtaccaccaa gagctgcccc tccactcccc 1260
ccccacactg tatcaaatgt atcacatttt ctcattgtga acacttttagc cttaattgaa 1320
aatgagcaac aaagctggac aattgctagt tgtatataaa atttaatctc accgaatgta 1380
cagttttcaa atttcacgtg tatattaagg aactgatgca tctgagcatt ctgaaagaaa 1440
gaaaaagaag ctacttttagc tgccacccca ttctagaaaa gtctcttatt ttcaagctgt 1500
tctaaatagc ttctgtctcag tttcccaaaa aggggtaccc agggccctcc tctgtgtgcc 1560
ccagctgcat cagccagctt ctagggtggc ccattgtttt ctgccacctg acaacatttt 1620
tcctcaatta ctgtacaact actgtataaa ataaaacaac tactgtataa aataaactct 1680
ctcttttccc tggaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1740
aaaaaaaaaa aaaaaaaaaa a 1761

```

<210> 71
 <211> 273
 <212> PRT
 <213> Homo sapiens

```

<400> 71
Met Ala Thr Trp Asp Glu Lys Ala Val Thr Arg Arg Ala Lys Val Ala
  1              5              10              15

Pro Ala Glu Arg Met Ser Lys Phe Leu Arg His Phe Thr Val Val Gly
      20              25              30

Asp Asp Tyr His Ala Trp Asn Ile Asn Tyr Lys Lys Trp Glu Asn Glu
      35              40              45

Glu Glu Glu Glu Glu Glu Glu Gln Pro Pro Pro Thr Pro Val Ser Gly
      50              55              60

Glu Glu Gly Arg Ala Ala Ala Pro Asp Val Ala Pro Ala Pro Gly Pro
      65              70              75              80

Ala Pro Arg Ala Pro Leu Asp Phe Arg Gly Met Leu Arg Lys Leu Phe
      85              90              95

Ser Ser His Arg Phe Gln Val Ile Ile Ile Cys Leu Val Val Leu Asp
      100             105             110

Ala Leu Leu Val Leu Ala Glu Leu Ile Leu Asp Leu Lys Ile Ile Gln
      115             120             125

Pro Asp Lys Asn Asn Tyr Ala Ala Met Val Phe His Tyr Met Ser Ile
      130             135             140

Thr Ile Leu Val Phe Phe Met Met Glu Ile Ile Phe Lys Leu Phe Val
      145             150             155             160

```

Asn

```

<400> 72
aatcgggagtg cctggaaagt taattccaat gtcatgacgt caaacttaaa atggctcgtct 60
gccaccatga agattgataa ccctcacata acatatactg ccagggtccc agtcgatgtc 120
catgaataca acctaacgca tctgcagcct tccacagatt atgaagtgtg tctcacagtg 180
tccaatatcc atcagcgacg tcaaaagtca tgcgtaaatg tcacaaccaa aaatgccgcc 240
ttcgcagtgcc acatctctga tcaagaaaacc agtacagccc ttgctgcagt aatggggggt 300
atgtttgccc tcattagcct tgcgtccatt gctgtggtac tttgccaaaa gattttaagag 360
aaaaaantac caccactcat taaaaaagta tatgcaaaaa acctcttcaa tcccactaaa 420
tgagctgtac ccaccactca ttaacctntg ggaaggtgac agcgagnaag acaaaagtgg 480

```

ttttgcagac accaagccaa cccaggtnga cacatccaga aggtattaca tgtggtaant 540
 cagaggatat tttgcttctg gtagtaagga gcacaaagac gtttttgctt tattctgcaa 600
 aagtgaacaa gttgaagact tttgtatttt tgactttgct agttttgtggc agagtggaga 660
 ggacgggtgg atatttcaaa tttttttagt atagcgtatc gcaagggttt gacacggctg 720
 ccagcgactc taggcttcca gtctgtgttt gggtttttatt cttatcatta ttatgattgt 780
 tattatatta ttatttttatt ttagttgttg tgctaaactc aataatgctg ttctaactac 840
 agtgctcaat aaaatgatta atgacaggat ggggttcccc tgtgctttta ccagtagcat 900
 gacccttcct gaagccatcc gtagaaag 928

<210> 73

<211> 52

<212> PRT

<213> Homo sapiens

<400> 73

Met Ile Val Ile Ile Leu Leu Phe Tyr Phe Ser Cys Cys Ala Lys Leu
 1 5 10 15

Asn Asn Ala Val Leu Thr Thr Val Leu Asn Lys Met Ile Asn Asp Arg
 20 25 30

Met Gly Phe Pro Cys Ala Phe Thr Ser Ser Met Thr Leu Pro Glu Ala
 35 40 45

Ile Arg Arg Lys
 50

<210> 74

<211> 49

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (9)

<400> 74

aaattaana aaaaaaaaaa aaaaaaaaaa aaataaagaa aaaaaaaaaa 49

<210> 75

<211> 597

<212> DNA

<213> Homo sapiens

<400> 75

attctacaag ataacttccc agtactttaaa aaaagtctca aagtcataaa caagaaagaa 60
 ctgaggggact attgcatatt ggagcgatct aaagaagtat tacaatttgt ggaattcttg 120
 attaaatcct ggaccagcaa aaggacatta gtgggaaaat tgatgaaatt caaatgagat 180
 cttatatatga agttaattgt gtcagtgtac atttcctggg ttccataatt gcaagtgatt 240
 atgtaagggtt tgtaaatatt aggagcagct gggtaaagggt tatacaaaaa ctctatacta 300
 tttttgcatt tttttctgta agttttaaac attttccaac taaaaagttg aaaacacatg 360
 tattagagac acatgcgtat gtgtctctaa taatcttaaa tatatttaag atgatagaag 420
 gaattcttga gatagtaaaa tgaagtcacc aaaaaacaaa caaagaaaca aaacgaaatc 480
 accaaaatct atcaataaat ttcaggtaat acttttggca gattcattcc tttgagatgg 540
 agtctcactc ccagctctggg caacgagcga aactccgtct aaaaaaaaaa aaaaaaa 597

<210> 76

<211> 89

<212> PRT

<213> Homo sapiens

<400> 76

Met Arg Ser Tyr Ile Glu Val Asn Cys Val Ser Val His Phe Leu Val
 1 5 10 15

Phe Ile Ile Ala Ser Asp Tyr Val Arg Phe Val Asn Ile Arg Ser Ser
 20 25 30

Trp Val Lys Val Ile Gln Lys Leu Tyr Thr Ile Phe Ala Phe Phe Ser
 35 40 45

Val Ser Leu Lys His Phe Pro Thr Lys Lys Leu Lys Thr His Val Leu
 50 55 60

Glu Thr His Ala Tyr Val Ser Leu Ile Ile Leu Asn Ile Phe Lys Met
 65 70 75 80

Ile Glu Gly Ile Leu Glu Ile Val Lys
 85

<210> 77

<211> 1804

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (1794)

<400> 77

ggagttcatt aggccagtgg cactgatttt tccttcccat catagctatt tattctagta 60
 agaggataat aaaagaaaatt tcctatacaa gaactgaaat tttccatttg tatggatcta 120
 gtcactttca gattttcaatt tgagggttaag tatataaagc acatcccaat tttatatgct 180
 gccttgagaa aattacagga tgcacggcaa tttgtaggaa tttcaaattg gatcatttaa 240
 acatttgaaa aattatttta aaaaccatct agtttgcttt tggatttttag acattaaagc 300
 ctatgttgtc ttgttaacag ggggtggaatg tataaccatc agattcagca tgtgatttca 360
 cctttgaatc tgagtatttc ttccctatct tctttgagtc atttttggag cagactgtca 420
 ccagtattga taactaagca ttaaagggaa aagttgcatt gcaactatgc attgggttcc 480
 tggaagaact tttcttttgt tttagtgaat gaagaggctt gatgggatca cttactgtaa 540
 ctccctctac ataaggaccc ctcttgcaag cagaacacaa aagaacatgc tcaaggagta 600
 tcccattttc tggataaatt gaagaagttt gctagtaatg tctttatact agcgtcttcc 660
 ttgtatccct ttgctggcaa gggaatacaa ggcgtcaaga ccacagatca aaacacccca 720
 catttgagtg gagtcttatt tttactccaa gaggagttat tcccttctag tctaaaattg 780
 gcagtttttt ctttttttta ataaaatttt taaaatattc ccaaaccagt ggaacacaga 840
 cactggctgc acttagtact gccaaaagcc aaggtcattt gcacatattc catcaacctg 900
 tcgagaatta ggcctcactt tataacccaa ggcagtggaag tgcattgcatt ctcttagctg 960
 ggcaacaat tatactgtag ttgtgatata acacatgttg cttttatttg tactgcacat 1020
 atccactgta cagccacttg ggagtatcgt ggttagcttg cagcaactgc tgtctgcatt 1080
 tatactgttt attgcataatt cttttccctg gaagtgaag agaaatgttt ttcttggtgc 1140
 attgattaca ttttataaat ttgcttagct ggaaagtgtg ggaaaagagg cctgtttgtc 1200
 aattgtacaa ccgattgtga agctctagtg tgaatatttt tacgtctgta ttagacattc 1260
 tctttgcaaa tctattgttc gattgaaatg taaatgaaat taaagatggg gtacacccat 1320
 catgtaaaaa cgaggcacca tctctaagat ggatttaatg ctcattttta aggcataatc 1380
 tcagcttcta tttaaaacta taatttaaaa taattctgta caatgaaatg gggaatatat 1440
 atgggaataa attctattcc atttatttca atttgaattt ccaaattgta atgtttccct 1500
 ttgtgtcata ggaataggat taaatggggg aagactagga tttataaggc ctgtatatgg 1560
 ggggagggca gagatggaac aatgaggggt gtgatgatag tgaatagcaa agagtgaatt 1620
 ctgtgtgttt ttgctgtagc actgaagtga agagatatta gctttggctg ttcacaaaat 1680

agagcatcat gattttcagt gtttgagaga aaattgatgg aaaaagtttg cagtacttga 1740
 catgtatttg catgcacaaa ataaaattat ttgtccacct taaaaaaaaa aaanaaaaaa 1800
 aaaa 1804

<210> 78
 <211> 37
 <212> PRT
 <213> Homo sapiens

<400> 78
 Met Lys Arg Leu Asp Gly Ile Thr Tyr Cys Asn Ser Phe Tyr Ile Arg
 1 5 10 15
 Thr Pro Ser Ala Ser Arg Thr Gln Lys Asn Met Leu Lys Glu Tyr Pro
 20 25 30
 Ile Phe Trp Ile Asn
 35

<210> 79
 <211> 360
 <212> DNA
 <213> Homo sapiens

<400> 79
 agaataccaa gactgtgtgt acacgcagat gtcagtggca gagaatgaag atcagcttcg 60
 tgcaagggt tatgacaaaa caccagactt cattttacaa gtaccagttg ctgtagaagg 120
 gcacataatt cactggattg aaagcaaagc ctcatttggg gatgaatgta gccaccacgc 180
 ctacctgcat gaccagttct ggagctactg gaataggggc ccaatataac agacaaatgg 240
 tgaaacagag ggatactcac taggaaacag atttgggccca ggcttagtca tctattggta 300
 tggatttatc caggagctgg actgcaaccg ggaaaggggc atcctgctca aagcctgttt 360

<210> 80
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 80
 Met Asp Leu Ser Arg Ser Trp Thr Ala Thr Gly Lys Gly Ala Ser Cys
 1 5 10 15
 Ser Lys Pro Val
 20

<210> 81
 <211> 202
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (136)

<220>
 <221> unsure
 <222> (138)

<400> 81

aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 120
 aaaaaaaaaa aaaaantnaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 180
 aaaaaaaaaa aaaaaaaaaa aa 202

<210> 82
 <211> 1189
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (1155)

<400> 82
 ggcccttcacg gcctaatacag atttttagtgg caagacagggt gatgataggg agaaagaaag 60
 agtcctgcct gtcaaaagtg gctgtgctat ataratgaac cctcacagggt agcacctctc 120
 agagagaata ratggtaaat gtttcttttc aggtctttaa aagtgtcagg ctatcagtta 180
 acctctccta gatctcagaa atgcctagaa agagaagtc tggctacatc aatggaaatt 240
 ctccacagat gcaaattttc tctacaaaag atggccttgc agagccacct cagtctgttg 300
 tccctgtagc agccatttca aattatgtca aagagatata ttttggggta aaatattttg 360
 attatcttca ttagtatatc tcaattttgt caatacaaac ctgagagtta tagtcagagg 420
 ttgaattttc atttcaaaat gtttctcctag ttttttttct cttttttgtt ttattgtaag 480
 ttgacaattt ataattgtat aaaagtatga ggtacaaagt gatgttatag cttagaata 540
 cagtatggta tgattaaatc aagttattaa cctatccttc acgttaaagt cttaaatttt 600
 ttgatagaaa catttgaaat ttactcttgg aaggtaaaaa aaaatctcag gaccccccaa 660
 attaaagcca tgaagctgaa ttgtgcaaca tcctcttcca aatggaagct tgtcttccag 720
 gtacagaaca aaaacaagac tcatttcttc acctgcctaa agatgtgcac ataattggct 780
 cctcctttac tccttttttc tcttctaaca ttcattatat cttgtgtaaa atgtagattt 840
 actggacact aactaaaatt tcacaggggt gtaccattt gccttactgc ctacctacct 900
 gtyttcttac gtaccttytc cccactttaa ggaatgcata catattaaac ctcccaaaaa 960
 cytytttaga aaaatagcca cagggttatc tgtggctgggt gtgtttccct agatgcacty 1020
 taaagctggc ttaataaacc tcagtattg aaacttatgc ctcaatcact ctttttggtt 1080
 gtcactgtct taccaatttg aaatgtaaaa tattytacta ctaattatat ttaccacatt 1140
 gtgcaacaga actcnaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1189

<210> 83
 <211> 56
 <212> PRT
 <213> Homo sapiens

<400> 83
 Met Arg Thr Phe Glu Ile Tyr Ser Trp Lys Val Lys Lys Asn Leu Arg
 1 5 10 15
 Thr Pro Gln Ile Lys Ala Met Lys Leu Asn Cys Ala Thr Ser Ser Ser
 20 25 30
 Lys Trp Lys Leu Val Phe Gln Val Gln Asn Lys Asn Lys Thr His Phe
 35 40 45
 Phe Thr Cys Leu Lys Met Cys Thr
 50 55

<210> 84
 <211> 525
 <212> DNA
 <213> Homo sapiens

<400> 84
cataacagcg tcagagagaa agaactgact gaaacgtttg agatgaagaa agttctcttc 60
ctgatcacag ccatcttggc agtggctgtt ggtttcccag tctctcaaga ccaggaacga 120
gaaaaaagaa gtatcagtga cagcgatgaa ttagcttcag ggttttttgt gttcccttac 180
ccatatccat ttcgcccact tccaccaatt ccatttccaa gatttccatg gtttagacgt 240
aattttccta ttccaatacc tgaatctgcc cctacaactc cccttcttag cgaaaagtaa 300
acaagaagga aaagtcacga taaacctggt cacctgaaat tgaaattgag ccacttcctt 360
gaagaatcaa aattcctgtt aataaaaagaa aaacaaatgt aattgaaata gcacacagca 420
ttctctagtc aatatcttta gtgatytyt ttaataaaca tgraagcaaa graaaaaaaa 480
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaa 525

<210> 85
<211> 85
<212> PRT
<213> Homo sapiens

<400> 85
Met Lys Lys Val Leu Leu Leu Ile Thr Ala Ile Leu Ala Val Ala Val
1 5 10 15
Gly Phe Pro Val Ser Gln Asp Gln Glu Arg Glu Lys Arg Ser Ile Ser
20 25 30
Asp Ser Asp Glu Leu Ala Ser Gly Phe Phe Val Phe Pro Tyr Pro Tyr
35 40 45
Pro Phe Arg Pro Leu Pro Pro Ile Pro Phe Pro Arg Phe Pro Trp Phe
50 55 60
Arg Arg Asn Phe Pro Ile Pro Ile Pro Glu Ser Ala Pro Thr Thr Pro
65 70 75 80
Leu Pro Ser Glu Lys
85

<210> 86
<211> 349
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> (9)

<220>
<221> unsure
<222> (159)

<220>
<221> unsure
<222> (188)

<220>
<221> unsure
<222> (230)

<220>
<221> unsure
<222> (232)

<220>

<221> unsure

<222> (270)

<400> 86

```

acttcgcanc ccatcccggc tggacgcgac cggggagtg agcagcccg tccccctcctc 60
ggtgccgcct ctgcccagcg tttgcttggc tgggctacca cctgcgctcg gacggcgctc 120
ggagggtcct cgccccggc ctgcctacct gaaaaccana actgatggct ctatttgag 180
tctttcanac aacattcttc ttaacattgc tgccttgag gacttaccan antgaagtct 240
tggctgaacg tttaccattg actcctgttn tcacttaaag tttccaccaa ttctacgcgt 300
cagagtttgc acttacaatg gactgtccac aaccttcctt atcatcagg 349

```

<210> 87

<211> 563

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (63)

<220>

<221> unsure

<222> (83)

<220>

<221> unsure

<222> (116)

<220>

<221> unsure

<222> (177)

<220>

<221> unsure

<222> (183)

<220>

<221> unsure

<222> (228)

<220>

<221> unsure

<222> (240)

<400> 87

```

tccattccag caccagccaa cagcacaaaa ttaatccttg acagggtgtc ctaccaaatt 60
tgngtcatag ccaacaacag tgnnggtgct tctcctgctt ctgtaatagt catttntgca 120
gaccccgaaa acaagaggt tgaggaagaa agaattgcag gcacagaggg tggattnttt 180
ttnttttggg aaccccaacc tggagatggt ataggttatg ttgtggantg gtgtgaccan 240
accaggatg tgcctcggcg atttccagtg gaagaatgta ggtcccaata ccacaagcac 300
agtcattagc acagatgctt ttaggccagg agttcgatat gacttcagaa tttatgggtt 360
atctacaaaa aggattgctt gtttattaga gaaaaaaaaac aggatactct caggaaacttg 420
ctccttcaga caaccctcac gtgctgggtg atacattgac atcccactcc ttcactctga 480
gttggaaga ttactctact gaatctcaac ctgggttttat acaagggtac catgtctatc 540
tgaaatccaa ggcgaggcag tgc 563

```

<210> 88

<211> 58

<212> PRT

<213> Homo sapiens

<400> 88

Arg Lys Lys Thr Gly Tyr Ser Gln Glu Leu Ala Pro Ser Asp Asn Pro
 1 5 10 15

His Val Leu Val Asp Thr Leu Thr Ser His Ser Phe Thr Leu Ser Trp
 20 25 30

Lys Asp Tyr Ser Thr Glu Ser Gln Pro Gly Phe Ile Gln Gly Tyr His
 35 40 45

Val Tyr Leu Lys Ser Lys Ala Arg Gln Cys
 50 55

<210> 89

<211> 361

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (102)

<220>

<221> unsure

<222> (105)

<220>

<221> unsure

<222> (153)

<220>

<221> unsure

<222> (186)

<220>

<221> unsure

<222> (191)

<220>

<221> unsure

<222> (252)..(253)

<400> 89

gaaaaggcag ttctttcaga tggttcagaa tgttgcaaatacaaaattga caaccggaa 60
 gaaaaggcat tgattgtgga caacctaaag ccagaatcct tntangagtt tttcatcact 120
 ccattcacta gtgctggtga aggccccagt gcnacgttca cgaagggtcac gactccggat 180
 gaacantcct ngatgetgat tcatatccta ctgcccattg ttttctgcgt cttgctcact 240
 atggtcatgt gnnacttgaa aagtcagtgg atcaaggaga cctgttatcc tgacatccct 300
 gacccttaca agagcagcat cctgtcatta ataaaattca aggtaaaaaa aaaaaaaaaa 360
 a 361

<210> 90

<211> 756

<212> DNA

<213> Homo sapiens

<220>
 <221> unsure
 <222> (37)

<220>
 <221> unsure
 <222> (54)

<220>
 <221> unsure
 <222> (376)

<220>
 <221> unsure
 <222> (433)

<400> 90
 attgaattct agacctgcgg ccgctgctcc ctcagangcc cctgatgtct gganaattgt 60
 gagcttggag ccaggaaaac atactgtgac cttattcttg aagccattat caaaactgca 120
 tgccaatgga aagatcctgt tctataatgt agttgtagaa aacctagaca aaccatccag 180
 ttcagagctc cattccattc cagcaccagc caacagcaca aaactaatcc ttgacagggtg 240
 ttccctacaa atctgcgtca tagccaacaa cagtgtgggt gcttctctctg cttctgtaat 300
 agtcattyct gcagaccccc aaaacaaaga gggtgaggaa gaaagaattg caggcacaga 360
 ggggtggattt ttttntttt ggaaacccca acctggagat gttataggtt atgttggtga 420
 ctggtgtgac canaccagg atgtgcctcg gtgatttcca gtggaagaat gtaggtccca 480
 ataccacaag cacagtcatt agcacagatg cttttaggcc aggagttcga tatgacttca 540
 gaatttatgg gttatctaca aaaaggattg cttgtttatt agagaaaaaa aacaggatac 600
 tctcaggaac ttgctccttc agacaacct cactgtctgg tggatacatt gacatccac 660
 tccttcactc tgagttggaa agattactct actgaatctc aacctgggtt tatacaaggg 720
 taccatgtct atctgaaatc caaggcgagg cagtgc 756

<210> 91
 <211> 58
 <212> PRT
 <213> Homo sapiens

<400> 91
 Arg Lys Lys Thr Gly Tyr Ser Gln Glu Leu Ala Pro Ser Asp Asn Pro
 1 5 10 15
 His Val Leu Val Asp Thr Leu Thr Ser His Ser Phe Thr Leu Ser Trp
 20 25 30
 Lys Asp Tyr Ser Thr Glu Ser Gln Pro Gly Phe Ile Gln Gly Tyr His
 35 40 45
 Val Tyr Leu Lys Ser Lys Ala Arg Gln Cys
 50 55

<210> 92
 <211> 79
 <212> DNA
 <213> Homo sapiens

<400> 92
 tcattaataa aattcaaggt aaatgttaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa 79

<210> 93

<211> 1939
 <212> DNA
 <213> Homo sapiens

<400> 93
 attggttttc aaatattaaa ccagcatttt attttaaaaa aaaccccact tagttatgat 60
 attttatctt ttttatataa tgtgggattt gatttgccga tattttatta aagatttttg 120
 catcatgttc tttttctgaa attttatttt gttctaattg cataaaataa gttggaaatt 180
 atcctctcta tctctacttt gtgaaagaat ttatattaca ttggtattat ttcttccttg 240
 aatgtttgat agactatact agtgaaagca tctgggccta ggggttttctc tgtggaagat 300
 tttcagttac aaattcaatt ttactgagtc aggtttgata agttacattt ctcaaggaaat 360
 ttatctattt aatcattgaa tgtattgaac attcatttgt ttataatttt gttttgttat 420
 tgaaaaatgtc tgtaagattt atagtgatgt tcccttttct attcctgaca ttgttaattt 480
 gtgttctctc tccctccatc cctctctcac tctctcttgt cagctgggtc gtgggattat 540
 caccctttatt aatcttttca gagaaccact ttttatttct ttgatttcct ttattgtttg 600
 ttaacttttt agttttattga ttccctctct tagctttatt acttccttcc ttcracttag 660
 taagagttaa atttgccctc atttttttag cttcttaagg taggaaattt gattatattt 720
 taaacttaaa aaaacttaaa gctataaaat tccctttaag cactgtttta gctgaattct 780
 acataatttg attgggtatg ttttcatcat tcaattaaaa acatagtttc taattttttc 840
 ttgcaatttc ttctttaacc cataggttat ttggcagttt gctgctgaat ttcaaagtgc 900
 ttgggagggt ttatagctat cttgttaaaa attgatttct aatttaattt tgtcagagaa 960
 catactttgt atgacttaaa tcttttaaaa tacgttcaga ctgatttagg gtccagcata 1020
 tgggtacgtc tagtgaatgt aatatatgcc attggaaatg atgcacattc tgcaattgtt 1080
 ggttgtaatg tgctataaat gtcaaccagg ttaacttggt tgatagtatt gttgaattcc 1140
 tgtgttccaa atgatatttt ttgtctgttt attctatcaa ttaatgagaa aggtttatta 1200
 aaatctccaa ccattgttgt gcatttattt ctatttcttt agttctgttg atttttgttt 1260
 tatatatttt gaagctctga gtacatctcc ctgatgagtt gaccctttta tcattatgaa 1320
 atgtcattct ttctctgtaa tgcttatctt aaaatctact ttgtttgata tgaatgtagc 1380
 cacttaagct ttcttatgca tactgtttgc aaagtgtgtg tgtgtgtgtg tgtgtgtgtg 1440
 tgatctatc tcaatccctt agtttccgcc tatctctgtc tttttattta aagtgtgttt 1500
 ttgcaggtag tgtataactt ggatcttgtt tttatattca gtcctgacaat ttatgcttct 1560
 tgattattgt gtttagttta tttgcatagt gtgtagtcca ttgacatttg gcataatatt 1620
 gataatttgt gttggagcgt attgtcttca tttctgattg tcccatctgg ttttgttctt 1680
 gttttgttgt tctctgttcc cacacttttt gtcttctttt gggttaatca aatattcttt 1740
 ggctttgcat cttatgtctt ctgttggatt tttatctata ccttgtctga tttttcttta 1800
 gtggtttctc taggggttac actatacatt attaatcaat catagtccat ttagaattaa 1860
 tattgtacac tttacataaa atgtaaaaca tttgtcacag tataaagtct attccttcca 1920
 aaaaaaaaaa aaaaaaaaaa 1939

<210> 94
 <211> 52
 <212> PRT
 <213> Homo sapiens

<400> 94
 Met Ser Val Arg Phe Ile Val Met Phe Pro Phe Leu Phe Leu Thr Leu
 1 5 10 15
 Leu Ile Cys Val Leu Ser Pro Ser Ile Pro Leu Ser Leu Ser Leu Val
 20 25 30
 Ser Trp Ser Val Gly Leu Ser Pro Leu Leu Ile Phe Ser Glu Asn His
 35 40 45
 Phe Leu Phe Leu
 50

<210> 95
 <211> 1252

<212> DNA

<213> Homo sapiens

<400> 95

```

gtagaaatcc caaaaagatt tggacagagg cagagcctat gtcactacac cttaaaggat 60
aacaagggtt' atatcaagac tcatgggtgaa catgtaggtt ttagaatttt catggatgcc 120
atactacttt ctttgactag aaagggtgaag atgccagatg tggagctctt tgtaattttg 180
ggagactggc ctttggaata aaagaaatcc aattcaaaca tccatccgat cttttcctgg 240
tgtggctcca cagattccaa ggatatcgtg atgcctacgt acgatttgac tgattctggt 300
ctggaaacca tgggcccggg aagtctggat atgatgtccg tgcaagctaa cacgggtcct 360
ccctgggaaa gcaaaaattc cactgccgtc tggagagggc gagacagccg caaagagaga 420
ctcgagctgg ttaaaactcag tagaaaacac ccagaactca tagacgctgb tttcaccaac 480
tttttcttct ttaaacacga tgaaaacctg tatggtccca ttgtgaaaca tatttcattt 540
tttgatttct tcaagcataa gtatcaaata aatattcgatg gcaactgtac agcttatcgc 600
ctgccatatt tgctagtgtg tgacagtgtt gtgctgaagc aggattccat ctactatgaa 660
catttttaca atgagctgca gccctggaaa cactacattc cagttaagag caacctgagc 720
gatctgctag aaaaacttaa atggggcgaaa gatcacgatg aagaggccaa aaagatagca 780
aaagcaggac aagaatttgc aagaaataat ctcatgggag atgacatatt ctgttattat 840
ttcaaaacttt tccaggaata tgccaattta caagtgaagt agccccaat ccgagagggc 900
atgaaaaggg tagaaccaca gactgaggac gacctcttcc cttgtacttg ccataggaaa 960
aagaccaaag atgaactctg atatgcaaaa taacttctat tagaataatg gtgctctgaa 1020
gactcttctt aactaaaaag aagaattttt ttaagtatta attccatgga caatataaaa 1080
tctgtgtgat tgtttgcagt atgaagacac atttctactt atgcagtatt ctcatgactg 1140
tactttaaag tacattttta gaattttata ataaaaccac ctttatttta aaggaaaaaa 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 1252

```

<210> 96

<211> 289

<212> PRT

<213> Homo sapiens

<400> 96

```

Met Asp Ala Ile Leu Leu Ser Leu Thr Arg Lys Val Lys Met Pro Asp
 1             5             10             15

Val Glu Leu Phe Val Asn Leu Gly Asp Trp Pro Leu Glu Lys Lys Lys
      20             25             30

Ser Asn Ser Asn Ile His Pro Ile Phe Ser Trp Cys Gly Ser Thr Asp
      35             40             45

Ser Lys Asp Ile Val Met Pro Thr Tyr Asp Leu Thr Asp Ser Val Leu
      50             55             60

Glu Thr Met Gly Arg Val Ser Leu Asp Met Met Ser Val Gln Ala Asn
      65             70             75             80

Thr Gly Pro Pro Trp Glu Ser Lys Asn Ser Thr Ala Val Trp Arg Gly
      85             90             95

Arg Asp Ser Arg Lys Glu Arg Leu Glu Leu Val Lys Leu Ser Arg Lys
      100            105            110

His Pro Glu Leu Ile Asp Ala Ala Phe Thr Asn Phe Phe Phe Lys
      115            120            125

His Asp Glu Asn Leu Tyr Gly Pro Ile Val Lys His Ile Ser Phe Phe
      130            135            140

Asp Phe Phe Lys His Lys Tyr Gln Ile Asn Ile Asp Gly Thr Val Ala

```

145 150 155 160
 Ala Tyr Arg Leu Pro Tyr Leu Leu Val Gly Asp Ser Val Val Leu Lys
 165 170 175
 Gln Asp Ser Ile Tyr Tyr Glu His Phe Tyr Asn Glu Leu Gln Pro Trp
 180 185 190
 Lys His Tyr Ile Pro Val Lys Ser Asn Leu Ser Asp Leu Leu Glu Lys
 195 200 205
 Leu Lys Trp Ala Lys Asp His Asp Glu Glu Ala Lys Lys Ile Ala Lys
 210 215 220
 Ala Gly Gln Glu Phe Ala Arg Asn Asn Leu Met Gly Asp Asp Ile Phe
 225 230 235 240
 Cys Tyr Tyr Phe Lys Leu Phe Gln Glu Tyr Ala Asn Leu Gln Val Ser
 245 250 255
 Glu Pro Gln Ile Arg Glu Gly Met Lys Arg Val Glu Pro Gln Thr Glu
 260 265 270
 Asp Asp Leu Phe Pro Cys Thr Cys His Arg Lys Lys Thr Lys Asp Glu
 275 280 285

Leu

<210> 97
 <211> 492
 <212> DNA
 <213> Homo sapiens

<400> 97
 ctaggttctg ggaagatggc gaaggtctca gagctttacg atgtcacttg ggaagaaatg 60
 agagataaaa tgagaaaatg gagagaagaa aactcaagaa atagtggagca aattgtggaa 120
 gttggagaag aattaattaa tgaatatgct tctaagctgg gagatgatat ttggatcata 180
 tatgaacagg tgatgattgc agcactagac tatggtcggg atgacttggc attgttttgt 240
 cttcaagagc tgagaagaca gttccctggc agtcacagag tcaagcgatt aacaggcatg 300
 agatttgaag ccatggaaag atatgatgat gctatacagc tatatgatag gattttacaa 360
 gaagatccaa ctaacactgc tgcaagaaag cgtaagattg ccattcgaaa agcccagggg 420
 aaaaatgtgg aggccattcg ggagctgaat gagtatctgg aacaatttgt tggagaccaa 480
 gaagcctggc at 492

<210> 98
 <211> 159
 <212> PRT
 <213> Homo sapiens

<400> 98
 Met Ala Lys Val Ser Glu Leu Tyr Asp Val Thr Trp Glu Glu Met Arg
 1 5 10 15
 Asp Lys Met Arg Lys Trp Arg Glu Glu Asn Ser Arg Asn Ser Glu Gln
 20 25 30
 Ile Val Glu Val Gly Glu Glu Leu Ile Asn Glu Tyr Ala Ser Lys Leu
 35 40 45

Gly Asp Asp Ile Trp Ile Ile Tyr Glu Gln Val Met Ile Ala Ala Leu
 50 55 60
 Asp Tyr Gly Arg Asp Asp Leu Ala Leu Phe Cys Leu Gln Glu Leu Arg
 65 70 75 80
 Arg Gln Phe Pro Gly Ser His Arg Val Lys Arg Leu Thr Gly Met Arg
 85 90 95
 Phe Glu Ala Met Glu Arg Tyr Asp Asp Ala Ile Gln Leu Tyr Asp Arg
 100 105 110
 Ile Leu Gln Glu Asp Pro Thr Asn Thr Ala Ala Arg Lys Arg Lys Ile
 115 120 125
 Ala Ile Arg Lys Ala Gln Gly Lys Asn Val Glu Ala Ile Arg Glu Leu
 130 135 140
 Asn Glu Tyr Leu Glu Gln Phe Val Gly Asp Gln Glu Ala Trp His
 145 150 155

<210> 99
 <211> 85
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (20)

<220>
 <221> unsure
 <222> (27)

<400> 99
 ccttggttaa taaaccatgn tgatttntta aacttgccaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa aaaaaa 85

<210> 100
 <211> 313
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (68)

<220>
 <221> unsure
 <222> (108)

<220>
 <221> unsure
 <222> (137)

<220>
 <221> unsure
 <222> (288)

<400> 100

```

ccttggaatta tctgaactaa aaattggaca gattgatcag ctggtaaaaa atctacttcc 60
tggaatttngt aaaggcaaaa acatttcttc ccattggcat acatcccntg tctctgcaca 120
atccttctttt gaaaatnaat atggtaactt aaatatattt agtacattac gttcctcttg 180
cttgatcga catcattcaa aagctcttca aagcatttct tcaaatcttc agtactggcc 240
agttttcata cagtctcggg gttttaaaac tttgaaatca aggacacnac gtctccagtc 300
tacctccgag aga 313

```

<210> 101

<211> 964

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (395)

<400> 101

```

gcttttccgg agcagagttg gagaatcttg tgaaccaggc tgcattaaaa gcagctgttg 60
atggaaaaga aatggttacc atgaaggagc tggagttttc caaagacaaa attctaattg 120
ggcctgaaag aagaagtgtg gaaattgata acaaaaacaa aaccatcaca gcatacatg 180
aatctgggtca tgccattatt gcatattaca caaaagatgc aatgcctatc aacaaagcta 240
caatcatgcc acggggggcca acacttggac atgtgtccct gttacctgag aatgacagat 300
ggaatgaac tagagcccag ctgcttgcac aaatggatgt tagtatggga ggaagagtgg 360
cagaggagct tatatttggg accgaccata ttacnacagg tgcttccagt gattttgata 420
atgccactaa aatagcaaaag cggatrgyta ccaaatttgg aatgagtga aagcttggag 480
ttatgacctc cagtataca ggggaaacta agtccagaaa cccaatctgc catcgaacaa 540
gaaataagaa tccttctaag ggactcatat gaacgagcaa aacatatctt gaaaactcat 600
gcaaaggagc ataagaatct cgcagaagct ttattgacct atgagacttt ggatgccaaa 660
gagattcaaa ttgttctkga ggggaaaaag ttggaagtga gatgataact ctctkgatat 720
ggatgctgc tggttttatt gcaagaatat aagtagcatt gcagtagtct acttttacia 780
cgctttcccc tcattcttga tgtggtgtaa ttgaagggtg tgaaatgctt tgtcaatcat 840
ttgtcacatt tatccagttt gggttattct cattaggaca cctattgcaa attagcatcc 900
catggcaaat atattttgaa aaaataaaga actatcagga ttgaaaaaaa aaaaaaaaaa 960
aaaa 964

```

<210> 102

<211> 166

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (125)..(126)

<400> 102

```

Met Val Thr Met Lys Glu Leu Glu Phe Ser Lys Asp Lys Ile Leu Met
  1             5             10             15

Gly Pro Glu Arg Arg Ser Val Glu Ile Asp Asn Lys Asn Lys Thr Ile
          20             25             30

Thr Ala Tyr His Glu Ser Gly His Ala Ile Ile Ala Tyr Tyr Thr Lys
          35             40             45

Asp Ala Met Pro Ile Asn Lys Ala Thr Ile Met Pro Arg Gly Pro Thr
          50             55             60

Leu Gly His Val Ser Leu Leu Pro Glu Asn Asp Arg Trp Asn Glu Thr

```

<400> 104
Met Pro Val Ala Thr Glu Leu Val Ile Val Ser Arg Ile Tyr Gln Tyr
1 5 10 15

Ile Glu Gln Ile Ile Met Phe Cys Phe Val Leu Phe Leu Phe Leu Tyr
 20 25 30
 Phe Arg Asn Ser Thr Ala Thr Tyr Lys Ser Ser Leu Glu Leu Ser Gly
 35 40 45
 Tyr Leu Lys Ser Glu Ala Ser Thr Phe Leu Arg Thr Lys His Arg Asn
 50 55 60
 Asp Glu Met Ser Tyr Lys Tyr Pro Phe Ile Leu Phe His Asn Thr Tyr
 65 70 75 80
 Ile Asp Leu Leu Tyr Val
 85

<210> 105
 <211> 479
 <212> DNA
 <213> Homo sapiens

<400> 105
 ctggcttctt cggagctgtg taktcttaat ttgctttgct gctggctggc tgacagctga 60
 tgggggactc ctcaggacgg actcccttcc agatgcaccc atctccatcc ttctcaactc 120
 cccaaccttt gtcctcccca ctcttcgctc gcgcggcggt ctgagaccac caggaccagt 180
 ttcaggggtt tccttctcca gcgagacttg gcagaacagg ctttaaaagc aaaggaggca 240
 gcggaagact gtgaattcct ttggacaatt gatgatattt atcattgtgc ccagtttcta 300
 caaataaaaag atgggtggat tattttctcg atggaggaca aaaccttcaa ctgtagaagt 360
 tctagaaagt atagataagg aaattcaagc attggaagaa tttagggaaa aaaaaaaaaa 420
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 479

<210> 106
 <211> 33
 <212> PRT
 <213> Homo sapiens

<400> 106
 Met Gly Gly Leu Phe Ser Arg Trp Arg Thr Lys Pro Ser Thr Val Glu
 1 5 10 15
 Val Leu Glu Ser Ile Asp Lys Glu Ile Gln Ala Leu Glu Glu Phe Arg
 20 25 30
 Glu

<210> 107
 <211> 333
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (70)

<220>
 <221> unsure
 <222> (156)

<220>
 <221> unsure
 <222> (184)

<220>
 <221> unsure
 <222> (207)

<220>
 <221> unsure
 <222> (308)

<400> 107
 ctgacttact aatactaaag aagtgggggg agctcgagag ccagacggcc agacaggcag 60
 acccctccan aggcccgcca ggtgggcatg gtccccatt ttctttaagg cagcacctgg 120
 agtggagaga ggccactccc tctccagccc ccgatntgga cccgggggag gggaggctga 180
 tgcntttggc cccggcctgg ccaaaanagc ccatccccag ggcagtttca ggtgccggct 240
 gggccctgaa tgtcaaggat aatatatagc ccgctcctgg gtccctggagc tgtggccctt 300
 tgtactcntg ttgtgtccat tgtgtgtgtg cgt 333

<210> 108
 <211> 611
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (62)

<220>
 <221> unsure
 <222> (185)

<220>
 <221> unsure
 <222> (192)

<220>
 <221> unsure
 <222> (249)

<220>
 <221> unsure
 <222> (290)

<400> 108
 gtccattgtg tgtgtgctgt gggacagagg cctggaaatg cggaggacta tacagagaag 60
 gnagggtttg tgaaggccag gcagggttgg aggcggggg tgtgagagga gaggccata 120
 gggctgagtg gggtcgggtg aggcagaggt cagaaacaga agagctgcag ttgctggagc 180
 tgggntgaga antgggctgc ctctgccat cccccgtct cctccccctt tcccccttgg 240
 gccccctnt gctcagaatc tgaagtagt ccctctcag caatttcatt tcttgaacac 300
 tgactcacac cttttaggca cctactgtgt gcatagcatt ccaccaggac tcatctccct 360
 tccttctcag ggggtcccg gcccgcacta gctttgcct aactccttca tcaaaagacc 420
 cccgccagc tccccacacc tcatacgag ccacatctgc cctattctcc atgctttcca 480
 gcttgctgc ccttctcat ctctccctgc ctgtgcagac ctccaccctt ctttcttcca 540
 cccctccatc cccaatgct ttagacatt ccattcattc cgtctcatcg tgcgtggtct 600
 ctgacgtcc a 611

<210> 109
 <211> 47

<212> PRT

<213> Homo sapiens

<400> 109

Met Leu Ser Ser Leu Pro Ala Leu Pro His Leu Ser Leu Pro Val Gln
 1 5 10 15

Thr Ser Thr Leu Ser Ser Thr Pro Pro Ser Pro Asn Ala Cys Arg
 20 25 30

Pro Ser Ile His Ser Val Ser Ser Cys Val Val Ser Asp Arg Pro
 35 40 45

<210> 110

<211> 274

<212> DNA

<213> Homo sapiens

<400> 110

atccaggcg tggggagacc attggcattt gggaaccatt ttccttcgaa cggcttcccc 60
 ttgagctgag cattctgctt gctgcagtag acgggtcgcc ttttgcccat accgaaattt 120
 tctgaaatta aatcgacac cccaccatt tcctctcccc tgggatctgg aggaacatca 180
 tacatagtag gtgaatcgtt ttgtagagtg aagaatgcta atgtaaagca aatagtcacc 240
 cacgttccct tgtaaatcca aaaaaaaaaa aaaa 274

<210> 111

<211> 1646

<212> DNA

<213> Homo sapiens

<400> 111

catcgggtgg actagctggg atctccgcat tggatttggg gctgattacc actgcttgcc 60
 tattattatt gttgtgttta ctactattat ttttttttac ccaagggaga aagacaaaaa 120
 aacggtggga tttatttaac atgatcttgg caaacgcctt ctgcctcttc ttctttctag 180
 acgagacctt ccgctctttg gccagccctt cctccctgca gggccccgag ctccacggct 240
 ggcgcccccc agtggactgt gtccgggcca atgagctgtg tgccgcccga tccaactgca 300
 gctctcgcta ccgcactctg cggcagtgcc tggcaggccg cgaccgcaac accatgctgg 360
 ccaacaagga gtgccaggcg gccttgaggg tcttgacagga gagccccgtg tacgactgcc 420
 gctgcaagcg gggcatgaag aaggagctgc agtgctcgca gatctactgg agcatccacc 480
 tggggctgac cgaggggtgag gagttctacg aagcctcccc ctatgagccg gtgacctccc 540
 gcctctcgga catcttcagg cttgcttcaa tcttctcagg gacaggggca gaccgggtgg 600
 tcagcgccaa gagcaaccat tgcctggatg ctgccaaaggc ctgcaacctg aatgacaact 660
 gcaagaagct gcgctcctcc tacatctcca tctgcaaccg cgagatctcg cccaccgagc 720
 gctgcaaccg ccgcaagtgc cacaaggccc tgcgccagtt ctccgaccgg gtgcccagcg 780
 agtacacctt ccgcatgctc ttctgtcctt gccaaagacca ggcgtgcyct gagcgccgcc 840
 ggcaaacctt cctgccagc tgctcctatg aggacaagga gaagcccaac tgcttgagcc 900
 tgcgtggcgt gtgccggact gaccacctgt gtcggtcccc gctggccgac ttccatgcca 960
 attgtcgagc ctccctaccg acggtcacca gctgccctgc ggacaattac caggcggtgc 1020
 tgggctctta tgctggcatg attgggtttg acatgacacc taactatgtg gactccagcc 1080
 ccactggcat cgtggtgtcc ccctgggtgca gctgtcgtgg cagcggggaa atggaggagg 1140
 agtgtgagaa gtctctcagg gacttcaccg agaaccctat cctccggaac gccatccagg 1200
 ccttttgcaa cggcacggac gtgaacgtgt ccccaaaagg cccctcgttc caggccaccc 1260
 agggccctcg ggtggagaag acgccttctt tgccagatga cctcagtgc agtaccagct 1320
 tggggagcaa gtgtcatcacc acctgcacct ctgtccagga gcaggggctg aaggccaaca 1380
 actccaaaga gttaagcatg tgcttcacag agctcacgac aaatatcatc ccaggaggta 1440
 acaaggtgat caaacctaac tcaggcccca gcagagccag accgtcggct gccttgaccg 1500
 tgctgtctgt cctgatgctg aaacaggcct tgtaggctgt gggaaccgag tcagaagatt 1560
 tttgaaagct acgcagacaa gaacagccgc ctgacgaaat ggaaacacac acagacacac 1620
 acacaccttg caaaaaaaaaa aaaaaa 1646

<210> 112

<211> 464

<212> PRT

<213> Homo sapiens

<400> 112

Met Ile Leu Ala Asn Val Phe Cys Leu Phe Phe Phe Leu Asp Glu Thr
 1 5 10 15

Leu Arg Ser Leu Ala Ser Pro Ser Ser Leu Gln Gly Pro Glu Leu His
 20 25 30

Gly Trp Arg Pro Pro Val Asp Cys Val Arg Ala Asn Glu Leu Cys Ala
 35 40 45

Ala Glu Ser Asn Cys Ser Ser Arg Tyr Arg Thr Leu Arg Gln Cys Leu
 50 55 60

Ala Gly Arg Asp Arg Asn Thr Met Leu Ala Asn Lys Glu Cys Gln Ala
 65 70 75 80

Ala Leu Glu Val Leu Gln Glu Ser Pro Leu Tyr Asp Cys Arg Cys Lys
 85 90 95

Arg Gly Met Lys Lys Glu Leu Gln Cys Leu Gln Ile Tyr Trp Ser Ile
 100 105 110

His Leu Gly Leu Thr Glu Gly Glu Glu Phe Tyr Glu Ala Ser Pro Tyr
 115 120 125

Glu Pro Val Thr Ser Arg Leu Ser Asp Ile Phe Arg Leu Ala Ser Ile
 130 135 140

Phe Ser Gly Thr Gly Ala Asp Pro Val Val Ser Ala Lys Ser Asn His
 145 150 155 160

Cys Leu Asp Ala Ala Lys Ala Cys Asn Leu Asn Asp Asn Cys Lys Lys
 165 170 175

Leu Arg Ser Ser Tyr Ile Ser Ile Cys Asn Arg Glu Ile Ser Pro Thr
 180 185 190

Glu Arg Cys Asn Arg Arg Lys Cys His Lys Ala Leu Arg Gln Phe Phe
 195 200 205

Asp Arg Val Pro Ser Glu Tyr Thr Tyr Arg Met Leu Phe Cys Ser Cys
 210 215 220

Gln Asp Gln Ala Cys Ala Glu Arg Arg Arg Gln Thr Ile Leu Pro Ser
 225 230 235 240

Cys Ser Tyr Glu Asp Lys Glu Lys Pro Asn Cys Leu Asp Leu Arg Gly
 245 250 255

Val Cys Arg Thr Asp His Leu Cys Arg Ser Arg Leu Ala Asp Phe His
 260 265 270

Ala Asn Cys Arg Ala Ser Tyr Gln Thr Val Thr Ser Cys Pro Ala Asp
 275 280 285

Asn Tyr Gln Ala Cys Leu Gly Ser Tyr Ala Gly Met Ile Gly Phe Asp
 290 295 300
 Met Thr Pro Asn Tyr Val Asp Ser Ser Pro Thr Gly Ile Val Val Ser
 305 310 315 320
 Pro Trp Cys Ser Cys Arg Gly Ser Gly Asn Met Glu Glu Glu Cys Glu
 325 330 335
 Lys Phe Leu Arg Asp Phe Thr Glu Asn Pro Cys Leu Arg Asn Ala Ile
 340 345 350
 Gln Ala Phe Gly Asn Gly Thr Asp Val Asn Val Ser Pro Lys Gly Pro
 355 360 365
 Ser Phe Gln Ala Thr Gln Ala Pro Arg Val Glu Lys Thr Pro Ser Leu
 370 375 380
 Pro Asp Asp Leu Ser Asp Ser Thr Ser Leu Gly Thr Ser Val Ile Thr
 385 390 395 400
 Thr Cys Thr Ser Val Gln Glu Gln Gly Leu Lys Ala Asn Asn Ser Lys
 405 410 415
 Glu Leu Ser Met Cys Phe Thr Glu Leu Thr Thr Asn Ile Ile Pro Gly
 420 425 430
 Ser Asn Lys Val Ile Lys Pro Asn Ser Gly Pro Ser Arg Ala Arg Pro
 435 440 445
 Ser Ala Ala Leu Thr Val Leu Ser Val Leu Met Leu Lys Gln Ala Leu
 450 455 460

<210> 113
 <211> 355
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (133)

<220>
 <221> unsure
 <222> (151)

<220>
 <221> unsure
 <222> (196)

<220>
 <221> unsure
 <222> (228)

<400> 113
 ctctcatttt tcctttcgaa gtgatatcca ctggaacgag atcaaattctg taagctggag 60
 atacaacttc cactacataa gatccagaag gtatatcatg aaccacaaaa ctccatctg 120
 tcttaaggaa acngacgtgc tcttctccgt ntaccagcac tcgggccgcc gagatccagt 180

```
cctgaggctt caccnttggg acaactgcac gccctcaat cttgaagnga tctcctatgc 240
cgaccccaact ccctcccgat ccctcagcag cagccccggg caccctccgag ttctggacat 300
cccccgatag cagcagcagc agcaggacgg gaaagaagcc ccacagagcg gccgc 355
```

<210> 114
 <211> 587
 <212> DNA
 <213> Homo sapiens

```
<400> 114
ggctggacag acttttctaat gaaccaatg gttatgatga tggttcttcc tttattgata 60
tttgtgcttc tgcctaaagt ggtcaacaca agtgatcctg acatgagacg ggaaatggag 120
cagtcaatga atatgctgaa ttccaacccat gagttgcctg atgtttctga gttcatgaca 180
agactcttct cttcaaaatc atctggcaaa tctagcagcg gcagcagtaa aacaggcaaa 240
agtggggctg gcaaaaggag gtagtcaggc cgtccaragc tggcatttgc acaaacacgg 300
caacactggg tggcatccaa gtcttggaac accgtgtgaa gcaactacta taaacttgag 360
tcaccccgac gttgatctct tacaactgtg tatgttaact ttttagcaca tggtttgtac 420
ttggtacacg agaaaaccca gctttcatct tttgtctgta tgaggccaat attgatgtca 480
ctgaattaat tacagtgtcc tatagaaaat gccattaata aattatatga actactatac 540
attatgtata ttaattaaaa catcttaatc cagaaaaaaa aaaaaaa 587
```

<210> 115
 <211> 81
 <212> PRT
 <213> Homo sapiens

```
<400> 115
Met Asn Pro Met Val Met Met Met Val Leu Pro Leu Leu Ile Phe Val
  1             5             10             15

Leu Leu Pro Lys Val Val Asn Thr Ser Asp Pro Asp Met Arg Arg Glu
             20             25             30

Met Glu Gln Ser Met Asn Met Leu Asn Ser Asn His Glu Leu Pro Asp
             35             40             45

Val Ser Glu Phe Met Thr Arg Leu Phe Ser Ser Lys Ser Ser Gly Lys
             50             55             60

Ser Ser Ser Gly Ser Ser Lys Thr Gly Lys Ser Gly Ala Gly Lys Arg
             65             70             75             80

Arg
```

<210> 116
 <211> 601
 <212> DNA
 <213> Homo sapiens

```
<400> 116
gatgtacctt ggcaccgcgc agcctgcggc ccccaacacc acctcccccg agctcaacct 60
gtccccaccg ctccctgggca ccgccctggc caatgggaca ggtgagctct cggagcacca 120
gcagtacgtg atcggcctgt tcctctcgtg cctctacacc atcttctctt tccccatcgg 180
ctttgtgggc aacatcctga tcctgggtgt gaacatcagc ttccgcgaga agatgacat 240
ccccgacctg tacttcatca acctggcggt gccggacctc atcctggtgg ccgactccct 300
cattgaggtg ttcaacctgc acgagcggta ctacgacatc gccgtcctgt gcaccttcat 360
gtcgctcttc ctgcaggtca acatgtacag cagcgtcttc ttcctcacct ggatgagctt 420
cgaccgctac atcgccctgg ccagggccat gcgctgcagc ctgttccgca ccaagcacca 480
```

cgcccggtg agctgtggcc tcatctggat ggcacccgtg tcagccacgc tgggtgccctt 540
 caccgacgtg cacctgcagc acaccgacga ggctgcttc tgtttcgcg atgtccggga 600
 g 601

<210> 117
 <211> 200
 <212> PRT
 <213> Homo sapiens

<400> 117
 Met Tyr Leu Gly Thr Ala Gln Pro Ala Ala Pro Asn Thr Thr Ser Pro
 1 5 10 15
 Glu Leu Asn Leu Ser His Pro Leu Leu Gly Thr Ala Leu Ala Asn Gly
 20 25 30
 Thr Gly Glu Leu Ser Glu His Gln Gln Tyr Val Ile Gly Leu Phe Leu
 35 40 45
 Ser Cys Leu Tyr Thr Ile Phe Leu Phe Pro Ile Gly Phe Val Gly Asn
 50 55 60
 Ile Leu Ile Leu Val Val Asn Ile Ser Phe Arg Glu Lys Met Thr Ile
 65 70 75 80
 Pro Asp Leu Tyr Phe Ile Asn Leu Ala Val Ala Asp Leu Ile Leu Val
 85 90 95
 Ala Asp Ser Leu Ile Glu Val Phe Asn Leu His Glu Arg Tyr Tyr Asp
 100 105 110
 Ile Ala Val Leu Cys Thr Phe Met Ser Leu Phe Leu Gln Val Asn Met
 115 120 125
 Tyr Ser Ser Val Phe Phe Leu Thr Trp Met Ser Phe Asp Arg Tyr Ile
 130 135 140
 Ala Leu Ala Arg Ala Met Arg Cys Ser Leu Phe Arg Thr Lys His His
 145 150 155 160
 Ala Arg Leu Ser Cys Gly Leu Ile Trp Met Ala Ser Val Ser Ala Thr
 165 170 175
 Leu Val Pro Phe Thr Ala Val His Leu Gln His Thr Asp Glu Ala Cys
 180 185 190
 Phe Cys Phe Ala Asp Val Arg Glu
 195 200

<210> 118
 <211> 419
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (80)

<220>

<221> unsure

<222> (178)

<400> 118

```

acgcctgagc gtcctccatc ttccaggatg gcagcaatgg cgctgtgcgg cctcaccagg 60
cccacgagga gcagcagcgn tcggcccggg gcagcaggaa ggcccttttg tggagcgccc 120
gccgtctgct ccgggggtggg tcagtcaactg cttgttgaca tcaacatggc aattgcantc 180
atgtggactg ggaccgtgcg agctgccgtg tgggttagtc ggggtgccagg acaatgaaat 240
actccagcac gtgtggctga cgaatttgtt ttacagaaa taacagctgg ggacaactgc 300
ggtgatgatg taaaaacctt ccataaaat gtaagaaaag ctgatgaggc tggtagcgtt 360
cagcctttgt caataaacct gtcatgtgcg gaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 419

```

<210> 119

<211> 714

<212> DNA

<213> Homo sapiens .

<220>

<221> unsure

<222> (646)

<220>

<221> unsure

<222> (649)

<400> 119

```

agccattggg acaggaaatg ccaaacaaca cccagataag gttgctgaag ccataattga 60
tgccattgaa gactttgtcc agaaaggatc agcccagtct gtgaaaaaag ttaaagtgtg 120
tatctttctg cctcaagtac tggatgtgtt ttatgccaac atgaagaaaa gagaaggagc 180
tcagctttct tcccaacagy ctartswtsy ytwmyttky akcatttttg ggcttttcaa 240
agcaatctcc ccaaaaaaag aatcatttgg ttttggaata gaaaacagaa tcagcaactt 300
ttcgggtgtg tgggtgaaat gtcacgtgtg tggaatacgc tatctcctgg ctacaagacc 360
tgattgaaaa agaacagtgt ctttacacca gtgaagatga gtgcatcaaa gactttgatg 420
aaaaggagta tcaggagtgt aatgagctgc agaagaagtt aaatattaac atttccctgg 480
accataagag acctttgatt aaggttttgg gaattagcag agatgtgatg caggctagag 540
atgaaattga ggcgatgatc aagagagttc gattggccaa agaacaggaa tcccgggcag 600
attgtatcag tgagtttata gaatggcagt ataatacaca taacantntt cattgtttta 660
acaaatgac caatctgaaa ttagaggatg caaggagaga aaaaaaaaaa aaaa 714

```

<210> 120

<211> 159

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (141) .. (142)

<400> 120

```

Phe Leu Gly Phe Ser Lys Gln Ser Pro Gln Lys Lys Asn His Leu Val
  1                      5                      10                      15

```

```

Leu Glu Lys Lys Thr Glu Ser Ala Thr Phe Arg Val Cys Gly Glu Asn
      20                      25                      30

```

```

Val Thr Cys Val Glu Tyr Ala Ile Ser Trp Leu Gln Asp Leu Ile Glu
    35                      40                      45

```

```

Lys Glu Gln Cys Pro Tyr Thr Ser Glu Asp Glu Cys Ile Lys Asp Phe
    50                      55                      60

```


Asp Glu Lys Glu Tyr Gln Glu Leu Asn Glu Leu Gln Lys Lys Leu Asn
 65 70 75 80
 Ile Asn Ile Ser Leu Asp His Lys Arg Pro Leu Ile Lys Val Leu Gly
 85 90 95
 Ile Ser Arg Asp Val Met Gln Ala Arg Asp Glu Ile Glu Ala Met Ile
 100 105 110
 Lys Arg Val Arg Leu Ala Lys Glu Gln Glu Ser Arg Ala Asp Cys Ile
 115 120 125
 Ser Glu Phe Ile Glu Trp Gln Tyr Asn Asp Asn Asn Xaa Xaa His Cys
 130 135 140
 Phe Asn Lys Met Thr Asn Leu Lys Leu Glu Asp Ala Arg Arg Glu
 145 150 155

<210> 121
 <211> 2681
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (2656)

<400> 121
 cataacttcc aatcttcttg gcctggcact aaggacctac tacagactgg cactgaaata 60
 tcttccctac cttatcattc cttgtgcttc tccacaagcc cactcttccc ctccatcata 120
 catgtgccga cctttccctgt ctcttttact ttgcagcacc aaatgcttcc tactttgttg 180
 tctaggagga acacatgtca cttttgtaag ctgctcgaaa gcaggggcca caccttcac 240
 cttgttttcc acacaacacc aagcacttag tagacacca ataaatcatt gctgaatgaa 300
 tgtattcagc ctggaattgc actaggattt tttggccaac acattgtatt cttaactgat 360
 accagacttc caatcaaata aaatccttaa gcctttttca tagtctttaa ttaaactact 420
 tctcttccat tatttccctt tgcctacttt tgaactgata ttcagaactt ttctgttaat 480
 gtttaatttt catccattat tcttgtctgt acagatcttt ttgatttttg acctctttat 540
 ctagtttttt tttttttttt ttggcttccc gttgtgttat ccacaggcac aatgggtata 600
 ttgactatcg ttcattgaag ttgttgatac aaatgttgaa caggagaaaa accagtagct 660
 tgccaacttg gcacctcatt ctctagcttg acagtaatct cttatctagt agttttttaga 720
 tatggttatt taatcaaata tcaactagct ctaattttgt tataattcat tcatgtatac 780
 agacaatgaa gaatcatctt tcttcattca acacatttat tgggaaccaa tcattgtcca 840
 aggcaatatg ctaggcattg tgtaatagaa agcgattaag cttttacccc tgcctctttt 900
 ggcttgcaaa gagcaggtaa gattcatgca cagacaaatg tagtaccagg taaatagtga 960
 ggccaacac aagaggcagg tcaaagggtt ttggattcag aggatggaga agctctttct 1020
 aggtggtgta aatatttgca gaatgagtag gctcataagt ggaggtgagg tacaggctaa 1080
 gctgctgtaa gaaagagtcc aaaatgcaaa ggatttacca caatgcagtt aatttttctc 1140
 acaacggtcg aggtaggcag gtggtccaag tccagttaat cagccctcct caacacaagg 1200
 cttcccttg taagctcagg atgaccgctc cagttctcat catctcccag gcagaaaaac 1260
 gggagaaaga gaactagacg taccaccagc tcattttaag gacatagcca ggcagggtgc 1320
 agtgactcac atctgtaatc caagcacttt gggagggtcg aggttgagg atagcttgaa 1380
 gccaggagtt tgagaccagc ttggataaga aagcaagacc ttgtctctac aaaaaattta 1440
 aaaatttaac atgtgtggtg gtgtgcacct atagtcccag ctactcagga ggctgagggtg 1500
 gaaggatcac ttgagcacag gagttggagg atacagttag ccatgatcac accacctcac 1560
 tccagcctgg gcaacatagt gagaccccat ctcttttaaa aacaaaacaa aaaaacatca 1620
 ccaggaagtt gcttacatgt cttctattcc tataacatcg gcctgagctt agtcacatgg 1680
 ccacaccag cagaaagaac tctaggaaat atggctttat gctaggtaac cccaaaccca 1740
 gctaaaactg ttgcttttga agaagggtga aacagacaat gtggaggaga attaccagtc 1800

```

tgccacaaag agaaagaatt ctatgtgagg aaaacgctct agaagagggg gctgattaat 1860
tagttatatc tcagccgaga ggatgttatg agaaacacag atttgggttt aatagaaaaa 1920
cttagaaccc ttcaagaagg aatgggcgga cttgagagtg tcctccgctt ctgaaaatga 1980
tcaagaaaaa tggtaagaat gtacacctca aggggtggatg gtaaagggtca gcctttaagc 2040
ctcaatggcg gtgtggttgt aggtggggtg ggggtggagg ccccgaaagac aggcagacta 2100
ccacagtagc cccgatggaa gcagtggagaa tgaactggaa ggagcggctg tgggagcgac 2160
aacgtgatga gaataaaccc ggcttggctc tgccctgtgc acacacaggt gagctgtgtg 2220
ctcccggatg tgtcagctgg tatatgcgtc tgtcagaagg cagctgggga gcactgttag 2280
ctcagagact aagaggcaga cccaggaagc ccttctttgc attggtcagg gtttgctgca 2340
ttttcccaag cccaggaat ggaacccagt tcttcttctt tctgtgtaaa ataataagca 2400
tcacaatagg ctgtgctcat gagaacgcat tttgtttcca cagaaatggt ttttctcact 2460
ctgtcttgat tttgatttct gttaactca gtaaacacat taccaaattt taaaataagg 2520
tgacttgttt tcccctaactc acagttcacc aaagggtattt catctgtttg ttctgaaaat 2580
gcagctgctg tctagattta tgtgtgctct gacaagaaat gttttgtgta acaataaaaa 2640
tcatttcctt tgatgnaaaa aaaaaaaaaa aaaaaaaaaa a
2681

```

<210> 122

<211> 132

<212> PRT

<213> Homo sapiens

<400> 122

```

Met Glu Ala Val Arg Met Asn Trp Lys Glu Arg Leu Trp Glu Arg Gln
 1             5             10             15
Arg Asp Glu Asn Lys Pro Gly Leu Ala Leu Pro Cys Ala His Thr Gly
          20             25             30
Glu Leu Cys Ala Pro Gly Cys Val Ser Trp Tyr Met Arg Leu Ser Glu
          35             40             45
Gly Ser Trp Gly Ala Leu Leu Ala Gln Arg Leu Arg Gly Arg Pro Arg
          50             55             60
Lys Pro Phe Phe Ala Leu Val Arg Val Cys Cys Ile Phe Pro Ser Pro
          65             70             75             80
Gly Asn Gly Thr Gln Phe Phe Phe Phe Leu Cys Lys Ile Ile Ser Ile
          85             90             95
Thr Ile Gly Cys Ala His Glu Asn Ala Phe Cys Phe His Arg Asn Val
          100            105            110
Phe Ser His Ser Val Leu Ile Leu Ile Ser Val Lys Leu Ser Lys His
          115            120            125
Ile Thr Lys Phe
          130

```

<210> 123

<211> 1585

<212> DNA

<213> Homo sapiens

<400> 123

```

ctaagctatt tgattctagg tctagaatgt tatctcttat tagaggatat gttaattttc 60
ctgcatttta ttcaatttatt aacttaacat ctctgattgc ctaccatgtg tcaggctctg 120
tactaaggat tgaggaccga aagatgaaca aaacatgggg cctaattcaa agatttcaca 180
atctggagag aaagtcagcc acatacaaaa aattataagg tagaatgtgc tataaaaaat 240

```

```

gatttaggta cagtgggaagt ttagaaaggc ttactataag atgtcgtatt tgaattgggt 300
ctgagcatga attagctctc aggtgaggga gggggttaaa gaaaactcta atgaagggaac 360
tctagtatgt gtgaaggcaa ggaggctggt gtgttttaggt tgcagcaaac caagtacaac 420
caggcttgga cttgagtgtc tggaaagaca ggaagatgcc atgctgagaa aaactgcccc 480
tgccaagctg agaaatgttc agcagaaaca taagggtgagc tgcataatgtc attttaaagt 540
ttctaggagc cacattttta aaaaatcaaa attaacaagt caaaaaataa aaagcaatgg 600
ggggagatta aatgcatatt actaagttaa agaagccaat caaaaaaggc tacatacctg 660
tatgattcca actatatgac attctgaaaa aggccaaact atagagacag taaatgatca 720
gtggttgcca ggagttaggg aggaggagg gatgaacagg cagagcacag aagattttta 780
gagcagtaga actattttgt atgatattat aatattgtag atacatatca ttataaattt 840
gtccaaaccc atagaatgta caccaagagt gaactctaag gtaactgtg gactttgggt 900
gataatgatg tgtcagtgtg gattgatcag ttgtaacaaa tgtaccattc tgggtgggaa 960
tgttgataat ggagtaggct atgcatgtgt ggggcagtgg gtatatggga aatatctata 1020
ccttctggct agttttgtct tgaacttgat ctaaaaata gcctactaag aaacacaagt 1080
caaattaatt ttaataatac attttattta acccaattta tcagaaatac taatatatta 1140
acatgtaatt gatataaaag ttattaacta gatattttac tttttttggt actgagtcct 1200
tgaaatctgg tctgtatttt acatttacag tacatctcaa ttcacattag ccacatttca 1260
gatactcagt acatacatga gtacctatgg ctagtggctg ctgtgttgga caggcaggt 1320
cttgaacact ggacttcctt gactcagaag cctcaattct cagccacagt gatatcctgc 1380
tccctaagta ctataatgat aaacacaaga ggagaggagc tttcagatga tcatctaatt 1440
ccatgacgtt agctgtgtgt ctccacactg cccggtggct ccagtctgaa gcacttaggc 1500
agtgtgtctc aacagaaata caatgagagc caatacgtga taagtgtcct atgggccaca 1560
ttgaaacagt aaaaaaaaaa aaaaa 1585

```

<210> 124

<211> 63

<212> PRT

<213> Homo sapiens

<400> 124

```

Met Leu Ser Leu Ile Arg Gly Tyr Val Asn Phe Pro Ala Phe Tyr Ser
  1             5             10             15

```

```

Phe Ile Asn Leu Thr Ser Leu Ile Ala Tyr His Val Ser Gly Ser Val
      20             25             30

```

```

Leu Arg Ile Glu Asp Pro Lys Met Asn Lys Thr Trp Gly Leu Ile Gln
  35             40             45

```

```

Arg Phe His Asn Leu Glu Arg Lys Ser Ala Thr Tyr Lys Lys Leu
  50             55             60

```

<210> 125

<211> 625

<212> DNA

<213> Homo sapiens

<400> 125

```

gatccaccac gttctgcctg gcttctctcca tccccagagg cactaaaagc agtatatttaa 60
ggttgggtgtc ttactccctg gaagcctgaa atgggtggaa tagcggtaag gcttgagtaa 120
aactagggga cagagggtct tatttgtcga ttttatttta taatttgacc acagcatctg 180
aactccctct ctccctgaa taagtatttt tcccacattt ttggatatat gtatggtaga 240
caattttttt ttaagacaca gagataaatg ttttctgtct ttggttacct ttcttttccc 300
ctttaaaggg aattagctat agaactgctt tgtaaagatg cttcttgata ttttactttt 360
gttcttttcc cctaactcatt cctttttctc cccactctc cagaaggcat aacccttctc 420
tccacacccc ctaccccccac ccccgctcta ggctcccatc ctttccatca agaccttcat 480
tagcttatga tatttgtgtc cgagatgtta taacaaggac tctgtcatgt atataagcta 540
tttcttgatc cattttaaag gaattgtaca ttgtgtagga aaaaaaaaaa aaaaaaaaaa 600
aaaaaaaaaa aaaaaaaaaa aaaaa 625

```

<220>

<221> UNSURE

<222> (83)

<220>

<221> UNSURE

<222> (480)

<220>

<221> UNSURE

<222> (482)

<220>

<221> UNSURE

<222> (490)

<400> 128

Met Lys Pro Lys Leu Met Tyr Gln Glu Leu Lys Val Pro Ala Glu Glu
 1 5 10 15

Pro Ala Asn Glu Leu Pro Met Asn Glu Ile Glu Ala Trp Lys Ala Ala
 20 25 30

Glu Lys Lys Ala Arg Trp Val Leu Leu Val Leu Ile Leu Ala Val Val
 35 40 45

Gly Phe Gly Ala Leu Met Thr Gln Leu Phe Leu Trp Glu Tyr Gly Asp
 50 55 60

Leu His Leu Phe Gly Pro Asn Gln Arg Pro Ala Pro Cys Tyr Asp Pro
 65 70 75 80

Cys Glu Xaa Val Leu Val Glu Ser Ile Pro Glu Gly Leu Asp Phe Pro
 85 90 95

Asn Ala Ser Thr Gly Asn Pro Ser Thr Ser Gln Ala Trp Leu Gly Leu
 100 105 110

Leu Ala Gly Ala His Ser Ser Leu Asp Ile Ala Ser Phe Tyr Trp Thr
 115 120 125

Leu Thr Asn Asn Asp Thr His Thr Gln Glu Pro Ser Ala Gln Gln Gly
 130 135 140

Glu Glu Val Leu Arg Gln Leu Gln Thr Leu Ala Pro Lys Gly Val Asn
 145 150 155 160

Val Arg Ile Ala Val Ser Lys Pro Ser Gly Pro Gln Pro Gln Ala Asp
 165 170 175

Leu Gln Ala Leu Leu Gln Ser Gly Ala Gln Val Arg Met Val Asp Met
 180 185 190

Gln Lys Leu Thr His Gly Val Leu His Thr Lys Phe Trp Val Val Asp
 195 200 205

Gln Thr His Phe Tyr Leu Gly Ser Ala Asn Met Asp Trp Arg Ser Leu
 210 215 220

Thr Gln Val Lys Glu Leu Gly Val Val Met Tyr Asn Cys Ser Cys Leu
 225 230 235 240

Ala Arg Asp Leu Thr Lys Ile Phe Glu Ala Tyr Trp Phe Leu Gly Gln
 245 250 255
 Ala Gly Ser Ser Ile Pro Ser Thr Trp Pro Arg Phe Tyr Asp Thr Arg
 260 265 270
 Tyr Asn Gln Glu Thr Pro Met Glu Ile Cys Leu Asn Gly Thr Pro Ala
 275 280 285
 Leu Ala Tyr Leu Ala Ser Ala Pro Pro Pro Leu Cys Pro Ser Gly Arg
 290 295 300
 Thr Pro Asp Leu Lys Ala Leu Leu Asn Val Val Asp Asn Ala Arg Ser
 305 310 315 320
 Phe Ile Tyr Val Ala Val Met Asn Tyr Leu Pro Thr Leu Glu Phe Ser
 325 330 335
 His Pro His Arg Phe Trp Pro Ala Ile Asp Asp Gly Leu Arg Arg Ala
 340 345 350
 Thr Tyr Glu Arg Gly Val Lys Val Arg Leu Leu Ile Ser Cys Trp Gly
 355 360 365
 His Ser Glu Pro Ser Met Arg Ala Phe Leu Leu Ser Leu Ala Ala Leu
 370 375 380
 Arg Asp Asn His Thr His Ser Asp Ile Gln Val Lys Leu Phe Val Val
 385 390 395 400
 Pro Ala Asp Glu Ala Gln Ala Arg Ile Pro Tyr Ala Arg Val Asn His
 405 410 415
 Asn Lys Tyr Met Val Thr Glu Arg Ala Thr Tyr Ile Gly Thr Ser Asn
 420 425 430
 Trp Ser Gly Asn Tyr Phe Thr Glu Thr Ala Gly Thr Ser Leu Leu Val
 435 440 445
 Thr Gln Asn Gly Arg Gly Gly Leu Arg Ser Gln Leu Glu Ala Ile Phe
 450 455 460
 Leu Arg Asp Trp Asp Ser Pro Tyr Ser His Asp Leu Asp Thr Ser Xaa
 465 470 475 480
 Asp Xaa Val Gly Asn Ala Cys Arg Leu Xaa
 485 490

<210> 129

<211> 6254

<212> DNA

<213> Homo sapiens

<400> 129

agggctcctgg gcagtcgcgc cagagctgag cggaggggcgc ggcgcgagaa cgaatctttg 60
 tgacattctc tctcagcatt ctttatcccc tgtttgctga agacttcgac caaagctggt 120
 cttagctggt ggcattctcc tgagaaaagg atagcttcag aaatcagaaa aacatttggg 180
 aggtgtctag cccagtyggac cttctgaaga gcaatgctaa gaagacgttt gggttaaaga 240

```

attaaaagga agaacaactt aagagcttct tcaaagtctc ccgcatgaaa attacttaaa 300
cggtgcacac aacgtttcac aaaatctttt gtgaaagaag aaaaggaaat tcagtgtgtg 360
agttccagca ggagtttaagc taatgcagct taacatcatg ccgacaaaaga agcgcttatac 420
tgccggcaga gtgcccttga ttctcttctt gtgccagatg attagtgcac tggaaagtacc 480
tcttgatcca aaacttcttg aagacttggg acagcctcca accatcacc aacagtctcc 540
aaaagattac attattgacc ctccgggagaa tattgtaatc cagtgtgaag ccaaagggaa 600
accgccccca agcttttctt ggaccctgaa tgggactcat tttgacatcg ataaagacc 660
tctgggtcacc atgaagcctg gcacaggaac gctcataatt aacatcatga gcgaaggaa 720
agctgagacc tatgaaggag tctatcagtg tacagcaagg aacgaacgag gagctgcagt 780
ttctaataac attgtgttcc gcccatccag atcaccattg tggaccaaag aaaaacttga 840
accaatcaca cttcaaagtg gtcagtcttt agtacttccc tgcagacccc caattggatt 900
accaccacct ataataatctt ggatggataa ttcttttcaa agacttccac aaagtgcagt 960
agtttctcaa ggtttgaatg gggaccttta tttttccaat gtccctccag aggcaccccg 1020
cgaagactat atctgttatg ctagatttaa tcatactcaa accatacagc agaagcaacc 1080
tatttctgtg aaggtgattt cagctaaatc aagtagagag aggccacca catttttaac 1140
tccagaaggc aatgcaagta acaaagagga attaagagga aatgtgcttt cactggagt 1200
cattgcagaa ggactgccta cccaatttat ttactgggca aaggaagatg gaatgctacc 1260
caaaaacagg acagtttata agaactttga gaaaaccttg cagatcattc atgtttcaga 1320
agcagactct ggaaattacc aatgtatagc aaaaaatgca ttaggagcca tccaccatac 1380
catttctgtt agagttaaag cggctccata ctggatcaca gccctcaaa atcttgtgt 1440
gtccccagga gaggatggga ccttgatctg cagagctaag ggcaaccca aaccagaat 1500
tagctgggta acaaatggag tcccaataga aattgcccct gatgaccca gcagaaaat 1560
agatggcgat accattatctt ttccaatgt tcaagaaaga tcaagtgcag tatatcagt 1620
caatgcctct aatgaatatg gatatttact ggcaaacgca tttgtaaatg tgctggctga 1680
gccaccacga atcctcacac ctgcaaacac actctaccag gtcattgcaa acaggcctgc 1740
tttactagac tgtgccttct ttgggtctcc tctcccaacc atcagatggg ttaaaggagc 1800
taaaggaagt gctcttcata aagatatcta tgttttacct gaaaatggaa ctttggaaat 1860
tcctgtggcc caaaaaggag gtacaggaac ttatacgtgt gttgcaagga ataaattag 1920
gatggcaaa aatgaagttc acttagaaat caaagatgct acatggatcg ttaaacagcc 1980
cgaatatgca gttgtgcaaa gagggagcat ggtgtccttt gaatgcaaag tgaaacatga 2040
tcacacctta tccctcactg tctgtgggt gaaggacaac agggaaactgc ccagtgatga 2100
aaggttcact gttgacaagg atcatctagt ggtagctgat gtcagtgcag atgacagcgg 2160
gacctacacg tgtgtggcca acaccactct ggacagcgtc tccgccagcg ctgtgcttag 2220
cgttgtgtct cctactccaa ctccagctcc cgtttacgat gtcccaatc ctcccttga 2280
cttagaactg acagatcaac ttgacaaaag tgttcagctg tcatggaccc caggcgatga 2340
caacaatagc ccattacaa aattcatcat cgaatatgaa gatgcaatgc acaagccagg 2400
gctgtggcac caccaaactg aagtttcttg aacacagacc acagcccagc tgaagctgtc 2460
tccttacgtg aactactcct tccgcgtgat ggcagtgaac agcattggga agagcttgcc 2520
cagcgaggcc tctgagcagt atttgacgaa agcctcagaa ccagataaaa accccacagc 2580
tgtggaagga ctgggatcag agcctgataa tttgtgtgatt acgtggaagc ctttgaatgg 2640
tttcgaatct atagggccag gccttcagta caaagttagc tggcgccaga aagatggtga 2700
tgatgaatgg acatctgttg ttgtggcaaa tgtatccaaa tatattgtct caggcacgcc 2760
aacctttgtt ccatacctga tcaaagtcca ggccctgaat gacatggggg ttgccccga 2820
gccagctgta gtcatgggac attctggaga agacctccca atgggtggctc ctgggaacgt 2880
gcgtgtgaat gtggtgaaca gtaccttagc cgagggtgcac tgggacccag tacctctgaa 2940
aagcatccga ggacacctac aaggctatcg gatttactat tgggaagacc agagttcatc 3000
taaaagaaac agacgtcaca ttgagaaaaa gatcctcacc ttccaaggca gcaagactca 3060
tggcatgttg ccggggctag agcccttttag ccactacaca ctgaatgtcc gagtgttcaa 3120
tgggaaaggg gagggcccag ccagccctga cagagtcttt aatactccag aaggagtccc 3180
cagtgtctcc tcgtctttga agattgtgaa tccaacactg gactctctca ctttggatg 3240
ggatccaccg agccaccgga atggcatctt gacagagtac acctaaagt atcagccaat 3300
taacagcaca catgaattag gccctctggg agatttgaaa attcctgcca acaagacacg 3360
gtggacttta aaaaatttaa atttcagcac tcgatataag ttttattct atgcacaaac 3420
atcagcagga tcaggaagtc aaattacaga ggaagcagta acaactgtgg atgaagctgg 3480
tattcttcca cctgatgtag gtgcaggcaa agcgtatggc agccggcagg tggatattgc 3540
aactcagggc tggttcattg gtctgatgtg tgctgtgtct ctcttatct taattttgt 3600
gattgtttgc ttcacagaa gaaacaaggg tggtaaatac ccagttaaag aaaaggaaga 3660
tgcccatgct gacctgaaa tccagcctat gaaggaagat gatgggacat ttggagaata 3720
cagtgtgca gaagaccaca agcctttgaa aaaaggaagt cgaactcctt cagacaggac 3780
tgtgaaaaaa gaagatagtg acgacagcct agttgactat ggagaagggg ttaatggcca 3840

```

```

gttcaatgag gatggctcct ttattggaca atacagtggg aagaaagaga aagagccggc 3900
tgaaggaaac gaaagctcag aggcaccttc tcctgtcaac gccatgaatt cctttgttta 3960
atttttaagc tctttgcaa tattccattt ctctagaatg tttatcctaa gcacttggtt 4020
gtcagccctc tcatactatg aacatatggg tagagagtat attttctgct gtatgttagt 4080
attatgagaa tagttacagc aaaaacataa ctcagtcaaa tgatatgtta atatgaactg 4140
gaatgcaaag tgcatacttt ttcattcaaa atgggtattc ttgatttcct cagaactgat 4200
aaaaaataat gcaacatcac caacagatcc tgttatttcc tctgcaggat acagttcaat 4260
atgatgcatg aaaaatgctc cacatttaaa ggacataccc gtgtatgtta tgaaaacatg 4320
gtttgatact ttgtttatac taccctcagc tgaaccccta tatatgaatt ccgttttcat 4380
tgtcaagaat gttactgttag tattctctag aacttcaatg tctttgtgga cattgttgtg 4440
aaattgggtg ctatgtatag ctgtcgttag tcttttggg agactgttag gaacagtttg 4500
tacagtatat acttgctaaa tgagttcatt atgacagtca cattgctgat gcttactgag 4560
aactattacc tactcttggc tcctgttact ccgtaggctt cttaatcttc caggcattac 4620
agcagcacag tgttctactt ttacatcat ttctatgttc ggtgtgtttt aggcataaac 4680
aatgtgtatt gcagtgcatt tcggcatttg tgccatactg aaagaatcaa aaacaaatca 4740
tccaaattaa atttcaaaca ttatttcaga gaacacaggg caagacacat acagtgcctt 4800
cagatattaa gcattccaca acatcgtgca ttctgtatca gctggtccag tccattcttg 4860
gtccttagatt actgtcattg tctaaaagta acttttaaaa agcagagttc atgaaaactg 4920
caatgctggg aaaagaagga aacatgaaaa taaaaataag acagtttatt agaaatagca 4980
tttctcata agcataaaaa gaaatctttg ttgccaactg aagcacatgg gattttgtgg 5040
tcttttatgg tttctataac attcagtaag aaagatgtca acatgctaga aaattaattt 5100
taaaactaag ttattccaac actaaaagca tacaacagca tgccaacagt aatatattat 5160
tctccaagac ttacctatg taagtgttca aaactctgca gcattaaaca acgtgtatgc 5220
aaattgttat ggatacattt cagaatctaa gaaatcaggg aagtgttaa aaggccaacg 5280
gtccaaggga ttacatctgc agtttaaaaa gtaaatatat attctatcgt attcataaac 5340
aatatctatc aaatgggtta cctccaaata tgaaaatcta taacaacctt tgggtgaagg 5400
aatgctcagt ttcatattgc aataaattgg tttctcataa cttgcatcaa gttaaatttt 5460
aagtaaagct ttttatatgt agatatattg ttgaatttgt aaatacactt aaaatgtaga 5520
tgctatatgc ttataggtgt tacatacaaa taaacatgca atgtttatgt tgtactgtat 5580
aagaggtaag ctaattaatg cagtgaatgg gattggaaag catctactta aatatctatt 5640
gggttcccc cttccccccac cttttttgct gtgaaactga aatagtgaac ttttctacgt 5700
attgacagca gatttttcga tgaaatcttc agagctttgc ctatggggca cagtaggcct 5760
agtaacctgg catgtttgat atatgtagg aaagcataat ttaaagtaat cccagggtaaa 5820
gatggcccta aatactttca tgtctctata ttcatttttc acagatccac ctgtctcttg 5880
aaaatataaa aagacaaaac aggtttgcct tggcatcaga gagcacaag attaaaagt 5940
actttaaat tgccaatatt ttgggagaac aataaaacta cattttttcy tyttccatc 6000
tggtagatgc gaaatttatc tgtgcatgaa agggctcact ctgtaatagt gcacagattt 6060
ggtattaaaa attaaatgtg gttttaaaag ttcttytytc ttttgaatt tatgttccca 6120
attgagtgtg aatgtccaag taatgggtga tgtaatgga caggcaaatg tgactggatt 6180
tccctcaaaa aagtaactat taaacagtyt tgatctcaaa aaaaaaaaaa aaaaaaaaaa 6240
aaaaaaaaa aaaa 6254

```

<210> 130

<211> 1192

<212> PRT

<213> Homo sapiens

<400> 130

```

Met Gln Leu Asn Ile Met Pro Thr Lys Lys Arg Leu Ser Ala Gly Arg
  1                      5                      10                      15

```

```

Val Pro Leu Ile Leu Phe Leu Cys Gln Met Ile Ser Ala Leu Glu Val
      20                      25                      30

```

```

Pro Leu Asp Pro Lys Leu Leu Glu Asp Leu Val Gln Pro Pro Thr Ile
      35                      40                      45

```

```

Thr Gln Gln Ser Pro Lys Asp Tyr Ile Ile Asp Pro Arg Glu Asn Ile
      50                      55                      60

```


Val Ile Gln Cys Glu Ala Lys Gly Lys Pro Pro Pro Ser Phe Ser Trp
 65 70 75 80
 Thr Arg Asn Gly Thr His Phe Asp Ile Asp Lys Asp Pro Leu Val Thr
 85 90 95
 Met Lys Pro Gly Thr Gly Thr Leu Ile Ile Asn Ile Met Ser Glu Gly
 100 105 110
 Lys Ala Glu Thr Tyr Glu Gly Val Tyr Gln Cys Thr Ala Arg Asn Glu
 115 120 125
 Arg Gly Ala Ala Val Ser Asn Asn Ile Val Val Arg Pro Ser Arg Ser
 130 135 140
 Pro Leu Trp Thr Lys Glu Lys Leu Glu Pro Ile Thr Leu Gln Ser Gly
 145 150 155 160
 Gln Ser Leu Val Leu Pro Cys Arg Pro Pro Ile Gly Leu Pro Pro Pro
 165 170 175
 Ile Ile Phe Trp Met Asp Asn Ser Phe Gln Arg Leu Pro Gln Ser Glu
 180 185 190
 Arg Val Ser Gln Gly Leu Asn Gly Asp Leu Tyr Phe Ser Asn Val Leu
 195 200 205
 Pro Glu Asp Thr Arg Glu Asp Tyr Ile Cys Tyr Ala Arg Phe Asn His
 210 215 220
 Thr Gln Thr Ile Gln Gln Lys Gln Pro Ile Ser Val Lys Val Ile Ser
 225 230 235 240
 Ala Lys Ser Ser Arg Glu Arg Pro Pro Thr Phe Leu Thr Pro Glu Gly
 245 250 255
 Asn Ala Ser Asn Lys Glu Glu Leu Arg Gly Asn Val Leu Ser Leu Glu
 260 265 270
 Cys Ile Ala Glu Gly Leu Pro Thr Pro Ile Ile Tyr Trp Ala Lys Glu
 275 280 285
 Asp Gly Met Leu Pro Lys Asn Arg Thr Val Tyr Lys Asn Phe Glu Lys
 290 295 300
 Thr Leu Gln Ile Ile His Val Ser Glu Ala Asp Ser Gly Asn Tyr Gln
 305 310 315 320
 Cys Ile Ala Lys Asn Ala Leu Gly Ala Ile His His Thr Ile Ser Val
 325 330 335
 Arg Val Lys Ala Ala Pro Tyr Trp Ile Thr Ala Pro Gln Asn Leu Val
 340 345 350
 Leu Ser Pro Gly Glu Asp Gly Thr Leu Ile Cys Arg Ala Asn Gly Asn
 355 360 365
 Pro Lys Pro Arg Ile Ser Trp Leu Thr Asn Gly Val Pro Ile Glu Ile
 370 375 380

Ala Pro Asp Asp Pro Ser Arg Lys Ile Asp Gly Asp Thr Ile Ile Phe
 385 390 395 400
 Ser Asn Val Gln Glu Arg Ser Ser Ala Val Tyr Gln Cys Asn Ala Ser
 405 410 415
 Asn Glu Tyr Gly Tyr Leu Leu Ala Asn Ala Phe Val Asn Val Leu Ala
 420 425 430
 Glu Pro Pro Arg Ile Leu Thr Pro Ala Asn Thr Leu Tyr Gln Val Ile
 435 440 445
 Ala Asn Arg Pro Ala Leu Leu Asp Cys Ala Phe Phe Gly Ser Pro Leu
 450 455 460
 Pro Thr Ile Glu Trp Phe Lys Gly Ala Lys Gly Ser Ala Leu His Glu
 465 470 475 480
 Asp Ile Tyr Val Leu His Glu Asn Gly Thr Leu Glu Ile Pro Val Ala
 485 490 495
 Gln Lys Asp Ser Thr Gly Thr Tyr Thr Cys Val Ala Arg Asn Lys Leu
 500 505 510
 Gly Met Ala Lys Asn Glu Val His Leu Glu Ile Lys Asp Ala Thr Trp
 515 520 525
 Ile Val Lys Gln Pro Glu Tyr Ala Val Val Gln Arg Gly Ser Met Val
 530 535 540
 Ser Phe Glu Cys Lys Val Lys His Asp His Thr Leu Ser Leu Thr Val
 545 550 555 560
 Leu Trp Leu Lys Asp Asn Arg Glu Leu Pro Ser Asp Glu Arg Phe Thr
 565 570 575
 Val Asp Lys Asp His Leu Val Val Ala Asp Val Ser Asp Asp Ser
 580 585 590
 Gly Thr Tyr Thr Cys Val Ala Asn Thr Thr Leu Asp Ser Val Ser Ala
 595 600 605
 Ser Ala Val Leu Ser Val Val Ala Pro Thr Pro Thr Pro Ala Pro Val
 610 615 620
 Tyr Asp Val Pro Asn Pro Pro Phe Asp Leu Glu Leu Thr Asp Gln Leu
 625 630 635 640
 Asp Lys Ser Val Gln Leu Ser Trp Thr Pro Gly Asp Asp Asn Asn Ser
 645 650 655
 Pro Ile Thr Lys Phe Ile Ile Glu Tyr Glu Asp Ala Met His Lys Pro
 660 665 670
 Gly Leu Trp His His Gln Thr Glu Val Ser Gly Thr Gln Thr Thr Ala
 675 680 685
 Gln Leu Lys Leu Ser Pro Tyr Val Asn Tyr Ser Phe Arg Val Met Ala
 690 695 700

Val Asn Ser Ile Gly Lys Ser Leu Pro Ser Glu Ala Ser Glu Gln Tyr
 705 710 715 720
 Leu Thr Lys Ala Ser Glu Pro Asp Lys Asn Pro Thr Ala Val Glu Gly
 725 730 735
 Leu Gly Ser Glu Pro Asp Asn Leu Val Ile Thr Trp Lys Pro Leu Asn
 740 745 750
 Gly Phe Glu Ser Asn Gly Pro Gly Leu Gln Tyr Lys Val Ser Trp Arg
 755 760 765
 Gln Lys Asp Gly Asp Asp Glu Trp Thr Ser Val Val Val Ala Asn Val
 770 775 780
 Ser Lys Tyr Ile Val Ser Gly Thr Pro Thr Phe Val Pro Tyr Leu Ile
 785 790 795 800
 Lys Val Gln Ala Leu Asn Asp Met Gly Phe Ala Pro Glu Pro Ala Val
 805 810 815
 Val Met Gly His Ser Gly Glu Asp Leu Pro Met Val Ala Pro Gly Asn
 820 825 830
 Val Arg Val Asn Val Val Asn Ser Thr Leu Ala Glu Val His Trp Asp
 835 840 845
 Pro Val Pro Leu Lys Ser Ile Arg Gly His Leu Gln Gly Tyr Arg Ile
 850 855 860
 Tyr Tyr Trp Lys Thr Gln Ser Ser Ser Lys Arg Asn Arg Arg His Ile
 865 870 875 880
 Glu Lys Lys Ile Leu Thr Phe Gln Gly Ser Lys Thr His Gly Met Leu
 885 890 895
 Pro Gly Leu Glu Pro Phe Ser His Tyr Thr Leu Asn Val Arg Val Val
 900 905 910
 Asn Gly Lys Gly Glu Gly Pro Ala Ser Pro Asp Arg Val Phe Asn Thr
 915 920 925
 Pro Glu Gly Val Pro Ser Ala Pro Ser Ser Leu Lys Ile Val Asn Pro
 930 935 940
 Thr Leu Asp Ser Leu Thr Leu Glu Trp Asp Pro Pro Ser His Pro Asn
 945 950 955 960
 Gly Ile Leu Thr Glu Tyr Thr Leu Lys Tyr Gln Pro Ile Asn Ser Thr
 965 970 975
 His Glu Leu Gly Pro Leu Val Asp Leu Lys Ile Pro Ala Asn Lys Thr
 980 985 990
 Arg Trp Thr Leu Lys Asn Leu Asn Phe Ser Thr Arg Tyr Lys Phe Tyr
 995 1000 1005
 Phe Tyr Ala Gln Thr Ser Ala Gly Ser Gly Ser Gln Ile Thr Glu Glu
 1010 1015 1020

Ala Val Thr Thr Val Asp Glu Ala Gly Ile Leu Pro Pro Asp Val Gly
 1025 1030 1035 1040
 Ala Gly Lys Ala Met Ala Ser Arg Gln Val Asp Ile Ala Thr Gln Gly
 1045 1050 1055
 Trp Phe Ile Gly Leu Met Cys Ala Val Ala Leu Leu Ile Leu Ile Leu
 1060 1065 1070
 Leu Ile Val Cys Phe Ile Arg Arg Asn Lys Gly Gly Lys Tyr Pro Val
 1075 1080 1085
 Lys Glu Lys Glu Asp Ala His Ala Asp Pro Glu Ile Gln Pro Met Lys
 1090 1095 1100
 Glu Asp Asp Gly Thr Phe Gly Glu Tyr Ser Asp Ala Glu Asp His Lys
 1105 1110 1115 1120
 Pro Leu Lys Lys Gly Ser Arg Thr Pro Ser Asp Arg Thr Val Lys Lys
 1125 1130 1135
 Glu Asp Ser Asp Asp Ser Leu Val Asp Tyr Gly Glu Gly Val Asn Gly
 1140 1145 1150
 Gln Phe Asn Glu Asp Gly Ser Phe Ile Gly Gln Tyr Ser Gly Lys Lys
 1155 1160 1165
 Glu Lys Glu Pro Ala Glu Gly Asn Glu Ser Ser Glu Ala Pro Ser Pro
 1170 1175 1180
 Val Asn Ala Met Asn Ser Phe Val
 1185 1190

 <210> 131
 <211> 4253
 <212> DNA
 <213> Homo sapiens

 <400> 131
 gcgatcagaa gactgaggca gtgctgcagg tgaacatggc tgtagcctac caatccttca 60
 ttttggtcga cttcataatt gaacctgctc actgccaccc agtcgttcac atggcaaaacc 120
 ccagtgttcc ttaagtgggc tgaaaagaat ggaatcctag acctgtcttg ttagagtara 180
 agctcattga gctgcctgcc ctcccatatt gcttgtggac ctatgaatgg tgctctttaa 240
 aatgtcagtg tagagggact gggaatcctg tattcctgcc ctatcattct gtaactttga 300
 gaagcatgtg ggctcagtc aractactta tttttagttc tgtagaaca ggatttttaa 360
 gaaaccatct gaacagaaga aataaagcgg aaggtaagct cagctcatgt gcccatcaag 420
 ctgcataaat ctccatctgc tctgggaaac attcatggga acagaaggat tcttcttctt 480
 gagaatgttg ttttcattat ataatcatag atacagcatg tgggtgattt ctgtgtacaa 540
 atactttacc catgagaaat gtatagctgt tagatatgtg tgggctcaca catatatgca 600
 atagatgcta tacatagcga tgtgcatgtg tatttgtata ttttatgtt catattttaa 660
 catatgaaac atgaaataga agtataatat tatttttgta tccaaatagt taacagagca 720
 gatgccctta gattctaata tgttgttcag tcctgaagat agccaagtta aatataaaaa 780
 caggagaaga ggaataataa aaktccattg agaaggaata gagagaaaga acrggtcttg 840
 tgaatgagaa kttcatttgg atcaacacag catttcttar aatgagtctg cactgtgctt 900
 tttgcarara craatgggct cctgaggctc tccacggctt ttgatggcar atgaktgtgc 960
 tctgtttcar atgaktcaag ctccagctct ggaaggcaga tgtcctcctc cgatgggtggg 1020
 ccaccgggccc agtcaracac agacagctcc cgtggaggag agcgaactcg acaccatgcc 1080
 agacattgag agtgataaaa acatcatccg gaccaagatg ttcctttacc tgtcagattt 1140
 gtccaggaag gaccggagaa ttgtcagcaa aaaatataaa atttattttt ggaacatcat 1200

```

caccattgct gtgttttacg cgctgcccgt gatccagctg gtcattacct atcagacagt 1260
ggtaaatgtc actggcaacc aggacatctg ttactacaac ttcctctgtg ctcacccttt 1320
gggcgtcctg agtgccttca acaacattct cagcaatctg ggccacgtgc tctctgggctt 1380
cctcttcctg ctgatagtct tgcgcgcgca catcctccat cggagagccc tggaaagccaa 1440
ggacatcttt gctgtggagt acgggattcc caaacacttt ggtctcttct acgctatggg 1500
cattgcattg atgatggaag ggggtgctcag tgcttgctac catgtctgcc ctaattattc 1560
caacttccaa ttcgacacct ccttcatgta catgatcgct ggccctgtgca tgctgaagct 1620
ctatcagacc cgccaccaccg acatcaatgc cagcgccctac tctgcctatg cctcctttgc 1680
tgtgttcate atggtcaccg tcttgggagt ggtgtttgga aaaaatgacg tatggttctg 1740
ggtcatcttc tctgcaatcc acgttctggc ctgcctagcc ctcagcacce aaatatatta 1800
tatgggtcgt ttcaagatag atgtgtctga cacagatttg ggaattttcc ggccgggctgc 1860
catggtgttc tacacagact gtatccagca gtgtagccga cctctatata tggatagaat 1920
ggtgttgctg gttgtgggga atctgggtta ctggctcttc gccctctttg gattgatata 1980
ccgccccagg gactttgctt cctacatgct gggcatcttc atctgtaacc ttttgctgta 2040
cctggccttt tacatcatca tgaagctccg cagctctgaa aaggctcctc cagtcccgct 2100
cttctgcate gtggccaccg ctgtgatgtg ggtgcccgc ctatattttt tcttccagaa 2160
tctcagcagc tgggagggaa ctccggccga atcccgggag aagaaccgag agtgcatctt 2220
gctggatttc ttcgatgacc atgacatctg gcaactcttc tctgctactg ctctgttttt 2280
ctcattcttg gttttgttaa ctttggatga tgacctgat gtggttcgga gagaccagat 2340
ccctgtcttc tgaacctcca acattaagag aggggagggga gcgatcaatc ttggtgctgt 2400
ttcacaaaaa ttacagtgc cagacaaaag taaccactgc cagatgctcc actcaccctc 2460
tgtagagcca actctgcatt cacacaggaa ggagaggggc tgcgggagat ttaaacctgc 2520
aagaaaggag cgagaagggg agccatgttt tgaggacaga cgaaaacctg agggagctgag 2580
aaacacttgc tcttccatc tgcagctttg ggagtgcac agggataggc actgcatcca 2640
agtcaactca ccatcttggg gtccctccca cctcacgga gacttgccag caatggcaga 2700
atgtctgtgc acacttccct ccagtgtgca cctgcccag aaaggccagc agcttggtgact 2760
tctgcccag aaactgtgtt ggcccccttc acacctctgc aacacctgct gctccagcaa 2820
gaggatgtga ttctttagaa tatggcgggg aggtgacccc aggcctgccc ctactgggat 2880
agatgtttta atggcaccag ctagtccact ccagaaagaa actctgtata tttccccag 2940
gtttctgatg ccacagaag ggctcaggag tgggtttgtt cacacattcc tcttaacaag 3000
taactgtcac tgggaccgag tcttgggtgc ttacatactt ctctgtgtct tcatctcact 3060
gacctgtgtg gacctcatca ctctgactct gccttctctg aaaggccctg tcaactccaca 3120
gatgtctggc cagcttcaag gcagaaggaa aaacaggaaa agctctttta acagcagcag 3180
gaacaagaga aatgactaac catactaaaa gactggtaac agcagcagca gccagacagg 3240
cctcacctta aggacttggg ctgccagagc aaattcagca gagcttattt ggccctccat 3300
tcacacagct cagttctgtg cccacatcac ctttggggaa gaaatcagca ttctaatacag 3360
ggacactact tcaggagtcc tccacagcga gtccgtcatc tgtcacttta tgtagatcag 3420
ggttctagac ttcttccctg aggttctcag aagcagctct caggatgaac gtattgtcct 3480
cttccccctt tcttgcaag tgacagcta atctaattgt gtctctcggg tgcacctgac 3540
attctctccc cagtaagggt ttggcaagct cagcatctgg gttccactct cacactgtct 3600
ggcagctctg tgcttgagaa gttctacatt gaccaggccc ccttgttgcc tggagtatga 3660
cgtaatcaga aatatagact ataaatgtgc acatgcgtat gtatttgctt gtgaaattaa 3720
agtcacctct tgcctctgct ttctgatca ttctgttagag aaatggatca ggcatttttt 3780
taaattatta ttctttctct aaactatttg cattgtgttc aaaaacccat tttagaagtt 3840
tgaacagcaa gcttttccctg attttaaaaa cacaaagtgt ctttcaatga aatatattgt 3900
gattttttta aagtcccaaa atgtgtactt agccttctgt tattccttat tctttaagca 3960
gtgttggctt ccattgacca tatgaaggcc accaattaaa tggttgtgtt aatccaacat 4020
gtaaaaaaact ttttggcagg gcacagtggc tcacgcctgt aatcccaaca ctttgggagg 4080
ctgaggcagg aggatcamtt gagcccagga gattgacgcc gcagtgaact atgattgtgc 4140
ccctgcactc cagcctggat gacagagtga gaccccatyt yttaaaaaat aaaaaaaaat 4200
aaaaatttga acctgtttca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 4253

```

<210> 132

<211> 479

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (13)..(14)

<220>

<221> UNSURE

<222> (21)

<400> 132

```

Met Gly Ser Val Gly Ser Pro Arg Leu Leu Met Ala Xaa Xaa Leu Ser
 1           5           10           15

Leu Phe Gln Met Xaa Gln Ala Pro Val Leu Glu Gly Arg Cys Pro Pro
          20           25           30

Pro Met Val Gly His Arg Ala Ser Gln Thr Gln Thr Ala Pro Val Glu
          35           40           45

Glu Ser Asp Phe Asp Thr Met Pro Asp Ile Glu Ser Asp Lys Asn Ile
          50           55           60

Ile Arg Thr Lys Met Phe Leu Tyr Leu Ser Asp Leu Ser Arg Lys Asp
          65           70           75           80

Arg Arg Ile Val Ser Lys Lys Tyr Lys Ile Tyr Phe Trp Asn Ile Ile
          85           90           95

Thr Ile Ala Val Phe Tyr Ala Leu Pro Val Ile Gln Leu Val Ile Thr
          100          105          110

Tyr Gln Thr Val Val Asn Val Thr Gly Asn Gln Asp Ile Cys Tyr Tyr
          115          120          125

Asn Phe Leu Cys Ala His Pro Leu Gly Val Leu Ser Ala Phe Asn Asn
          130          135          140

Ile Leu Ser Asn Leu Gly His Val Leu Leu Gly Phe Leu Phe Leu Leu
          145          150          155          160

Ile Val Leu Arg Arg Asp Ile Leu His Arg Arg Ala Leu Glu Ala Lys
          165          170          175

Asp Ile Phe Ala Val Glu Tyr Gly Ile Pro Lys His Phe Gly Leu Phe
          180          185          190

Tyr Ala Met Gly Ile Ala Leu Met Met Glu Gly Val Leu Ser Ala Cys
          195          200          205

Tyr His Val Cys Pro Asn Tyr Ser Asn Phe Gln Phe Asp Thr Ser Phe
          210          215          220

Met Tyr Met Ile Ala Gly Leu Cys Met Leu Lys Leu Tyr Gln Thr Arg
          225          230          235          240

His Pro Asp Ile Asn Ala Ser Ala Tyr Ser Ala Tyr Ala Ser Phe Ala
          245          250          255

Val Val Ile Met Val Thr Val Leu Gly Val Val Phe Gly Lys Asn Asp
          260          265          270

Val Trp Phe Trp Val Ile Phe Ser Ala Ile His Val Leu Ala Ser Leu
          275          280          285

```

Ala Leu Ser Thr Gln Ile Tyr Tyr Met Gly Arg Phe Lys Ile Asp Val
 290 295 300

Ser Asp Thr Asp Leu Gly Ile Phe Arg Arg Ala Ala Met Val Phe Tyr
 305 310 315 320

Thr Asp Cys Ile Gln Gln Cys Ser Arg Pro Leu Tyr Met Asp Arg Met
 325 330 335

Val Leu Leu Val Val Gly Asn Leu Val Asn Trp Ser Phe Ala Leu Phe
 340 345 350

Gly Leu Ile Tyr Arg Pro Arg Asp Phe Ala Ser Tyr Met Leu Gly Ile
 355 360 365

Phe Ile Cys Asn Leu Leu Leu Tyr Leu Ala Phe Tyr Ile Ile Met Lys
 370 375 380

Leu Arg Ser Ser Glu Lys Val Leu Pro Val Pro Leu Phe Cys Ile Val
 385 390 395 400

Ala Thr Ala Val Met Trp Ala Ala Ala Leu Tyr Phe Phe Phe Gln Asn
 405 410 415

Leu Ser Ser Trp Glu Gly Thr Pro Ala Glu Ser Arg Glu Lys Asn Arg
 420 425 430

Glu Cys Ile Leu Leu Asp Phe Phe Asp Asp His Asp Ile Trp His Phe
 435 440 445

Leu Ser Ala Thr Ala Leu Phe Phe Ser Phe Leu Val Leu Leu Thr Leu
 450 455 460

Asp Asp Asp Leu Asp Val Val Arg Arg Asp Gln Ile Pro Val Phe
 465 470 475

<210> 133
 <211> 462
 <212> DNA
 <213> Homo sapiens

<400> 133
 gtggaaggag tggataataa aatgagtcag tgcaccagct ccaccattcc tagctcaagt 60
 caagagaaag accctaaaat taagacagag acaagtgaag agggatcttg agacttggat 120
 aatctagatg ctattcttgg tgatctgact agttctgact tttacaataa ttccatatcc 180
 tcaaatggta gtcattctgg gactaagcaa cagggtgttc aaggaactaa ttctctgggt 240
 ttgaaaagtt cacagtctgt gcagtctatt cgtcctccat ataaccgagc agtgtctctg 300
 gatagccctg tttctgttgg ctcaagtcct ccagtaaaaa atatcagtgc tttcccatg 360
 ttaccaaagc aaccatgtt gggtgggaat ccaagaatga tggatagtca rgaaaattat 420
 ggctcaagta tgggagactg gggcttacca aactcaaagg cc 462

<210> 134
 <211> 147
 <212> PRT
 <213> Homo sapiens

<400> 134
 Met Ser Gln Cys Thr Ser Ser Thr Ile Pro Ser Ser Ser Gln Glu Lys
 1 5 10 15

Asp Pro Lys Ile Lys Thr Glu Thr Ser Glu Glu Gly Ser Gly Asp Leu
 20 25 30
 Asp Asn Leu Asp Ala Ile Leu Gly Asp Leu Thr Ser Ser Asp Phe Tyr
 35 40 45
 Asn Asn Ser Ile Ser Ser Asn Gly Ser His Leu Gly Thr Lys Gln Gln
 50 55 60
 Val Phe Gln Gly Thr Asn Ser Leu Gly Leu Lys Ser Ser Gln Ser Val
 65 70 75 80
 Gln Ser Ile Arg Pro Pro Tyr Asn Arg Ala Val Ser Leu Asp Ser Pro
 85 90 95
 Val Ser Val Gly Ser Ser Pro Pro Val Lys Asn Ile Ser Ala Phe Pro
 100 105 110
 Met Leu Pro Lys Gln Pro Met Leu Gly Gly Asn Pro Arg Met Met Asp
 115 120 125
 Ser Gln Glu Asn Tyr Gly Ser Ser Met Gly Asp Trp Gly Leu Pro Asn
 130 135 140
 Ser Lys Ala
 145

 <210> 135
 <211> 119
 <212> DNA
 <213> Homo sapiens

 <400> 135
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 119

 <210> 136
 <211> 3316
 <212> DNA
 <213> Homo sapiens

 <400> 136
 ggtaagatgg cggctgtgag tctgcggctc ggcgacttgg tgtgggggaa actcggccga 60
 tatctctcctt ggccaggaaa gattgttaat ccaccaagg acttgaagaa acctcgcgga 120
 aagaaatgct tctttgtgaa attttttgga acagaagatc atgcctggat caaagtggaa 180
 cagctgaagc catatcatgc tcataaagag gaaatgataa aaattaacaa gggtaaacga 240
 ttccagcaag cggtagatgc tgtcgaagag ttctcagga gagccaaagg gaaagaccag 300
 acgtcatccc acaattcttc tgatgacaag aatcgacgta attccagtga ggagagaagt 360
 aggccaaact caggtgatga gaagcgcaaa cttagcctgt ctgaaggga ggtgaagaag 420
 aacatgggag aaggaaagaa gaggggtgtct tcaggctctt cagagagagg ctccaatcc 480
 cctctgaaaa gagcccaaga gcaaagtccc cggaagcggg gtcggccccc aaaggatgag 540
 aaggatctca ccatcccga gtctagtacc gtgaagggga tgatggccgg accgatggcc 600
 gcgtttaaat ggcagccaac cgcaagcgag cctgttaaag atgcagatcc tcatttccat 660
 catttctcgc taagccaaac agagaagcca gctgtctgtt accaggcaat cagcaagaag 720
 ttgaaaatat gtgaagagga aactggtccc acctccatcc aggcagctga cagcacagcc 780
 gtgaatggca gcatcacacc cacagacaaa aagataggat ttttgggect tggctctcatg 840
 ggaagtggaa tcgtctccaa cttgctaaaa atgggtcaca cagtgtactgt ctggaaccgc 900
 actgcagaga aagagggggc ccgtctggga agaaccctcg ctgaagtcgt ctcaacctgc 960


```

gacatcactt tcgcctgcgt gtcggatccc aaggcgccca aggacctggt gctggggcccc 1020
agtgggtgtc tgcaagggat ccgccctggg aagtgtctacg tggacatgtc aacagtggac 1080
gctgacaccg tctactgagct ggcccagggtg attgtgtcca ggggggggcg ctttctggaa 1140
gccccgtct cagggaaatca gcagctgtct aatgacggga tgttggtgat cttagcggct 1200
ggagacaggg gcttatatga ggactgcagc agctgcttcc aggcgatggg gaagacctcc 1260
ttcttcctag gtgaagtggg caatgcagcc aagatgatgc tgatcgtgaa catggtccaa 1320
gggagcttca tggccactat tgccgagggg ctgacctggtg cccaggtgac aggccagtcc 1380
cagcagacac tcttggacat cctcaatcag ggacagttgg ccagcatctt cctggaccag 1440
aagtgcacaa atatcctgca aggaaacttt aagcctgatt tctacctgaa atacattcag 1500
aaggatctcc gcttagccat tgcgctgggt gatgcggtca accatccgac tcccatggca 1560
gctgcagcaa atgaggtgta caaagagcc aaggcgctgg accagtctga caacgatatg 1620
tccgccgtgt accgagccta catacactaa gctgtcgaca ccccgccctc accctccaa 1680
tccccctct gacccctct tccctacatg ggttcggggg cctgggagtt cattctggac 1740
cagccacctt atctccattt ccttttatac agactttgag acttgccatc agcacagcac 1800
acagcagcac ccttcccttg aggcgggtgg ggaggggaca agtgtcagca ggattggcgt 1860
gtgggaaagc tcttgagctg ggcactggcc ccccgagcga ggtggctgtg tgttcacaca 1920
cacacacaca cacacacaca cacaggtctt cggccagga tagaagctgc ccagaaactg 1980
ctgectgggt ttttttcttc cgagcttgtc ttatctcaaa ccccttccag tcaaggaact 2040
agaatcagca acgagagttg gaagccttcc cacagcttcc cccagagcga agaggctgta 2100
gtcatgtccc catccccac tggattccct acaaggagag gccttgggccc cagatgagcc 2160
agtacagact ccagacagag gggcccttgg ggccttccaa cctcaggtga tgagctgaga 2220
aagatgttca cgtctaagcg tccagtgtgc acccagcgct ccatagacgc ctttgtgaac 2280
tgaaaagaga ctggcagagt cccgagaaga tggggccctg gctttccagg gagtgcagca 2340
agcagccggc ctgcagaccc agcctgacca acgatgagca tttcttaggc tcagctcttg 2400
atacggaaac gagtgtcttc actccagcca gcatcatggt cttcggtgct tcccgggccc 2460
ggggtctgtc gggagggaag agaactgggc ctgacctacc tgaactgact ggccctccga 2520
ggtgggtctg ggacatccta gaggccctac atttgcctt ggatagggga ccgggggggg 2580
cttgggaatg tgcaaaaaaa aaagttaacc aagggatgtc agttttttat ccctctgcat 2640
gggttggatt ttccaaaatc ataatttgca gaaggaaggc cagcatttac gatgcaatat 2700
gtaattatat ataggggtgg cactactagg cggggtcctt ccccccctac agctttggcc 2760
cctttcagag attagaaact gggttagagg attgcagaag acgagtgggg ggagggcagg 2820
gaagatgcct gtcgggtttt tagcacagtt catttcactg ggattttgaa gcattttctg 2880
ctggacacaa agcctgttct agtctggcg gaacacactg ggggtggggg cgggggaaga 2940
tgcggtaatg aaaccggtta gtcaattttg tcttaatat gttgacaatt ctgtaaagt 3000
cctttttatg aatattttctg tttaagctat ttcaccttcc ttttgaaatc cttccctttt 3060
aaggagaaaa tgtgcactt gtgaaaaagc ttgtaagaaa gcccctccct tttttcttta 3120
aacctttaa tgacaaatct aggtaatata ggttgtaaat ttttattttt gctttgtttt 3180
taatgaacat ttgtctttca gaataggatt gtgtgataat gtttaaatgg caaaaacaaa 3240
acatgatttt gtgcaattaa caaagctact gcaagaaaaa taaaacactt cttggttaaca 3300
caaaaaaaaa aaaaaa 3316

```

<210> 137

<211> 547

<212> PRT

<213> Homo sapiens

<400> 137

```

Met Ala Ala Val Ser Leu Arg Leu Gly Asp Leu Val Trp Gly Lys Leu
  1             5             10             15

```

```

Gly Arg Tyr Pro Pro Trp Pro Gly Lys Ile Val Asn Pro Pro Lys Asp
          20             25             30

```

```

Leu Lys Lys Pro Arg Gly Lys Lys Cys Phe Phe Val Lys Phe Phe Gly
  35             40             45

```

```

Thr Glu Asp His Ala Trp Ile Lys Val Glu Gln Leu Lys Pro Tyr His
  50             55             60

```

```

Ala His Lys Glu Glu Met Ile Lys Ile Asn Lys Gly Lys Arg Phe Gln

```

65	70	75	80
Gln Ala Val Asp Ala Val Glu Glu Phe Leu Arg Arg Ala Lys Gly Lys	85	90	95
Asp Gln Thr Ser Ser His Asn Ser Ser Asp Asp Lys Asn Arg Arg Asn	100	105	110
Ser Ser Glu Glu Arg Ser Arg Pro Asn Ser Gly Asp Glu Lys Arg Lys	115	120	125
Leu Ser Leu Ser Glu Gly Lys Val Lys Lys Asn Met Gly Glu Gly Lys	130	135	140
Lys Arg Val Ser Ser Gly Ser Ser Glu Arg Gly Ser Lys Ser Pro Leu	145	150	155
Lys Arg Ala Gln Glu Gln Ser Pro Arg Lys Arg Gly Arg Pro Pro Lys	165	170	175
Asp Glu Lys Asp Leu Thr Ile Pro Glu Ser Ser Thr Val Lys Gly Met	180	185	190
Met Ala Gly Pro Met Ala Ala Phe Lys Trp Gln Pro Thr Ala Ser Glu	195	200	205
Pro Val Lys Asp Ala Asp Pro His Phe His His Phe Leu Leu Ser Gln	210	215	220
Thr Glu Lys Pro Ala Val Cys Tyr Gln Ala Ile Thr Lys Lys Leu Lys	225	230	235
Ile Cys Glu Glu Glu Thr Gly Ser Thr Ser Ile Gln Ala Ala Asp Ser	245	250	255
Thr Ala Val Asn Gly Ser Ile Thr Pro Thr Asp Lys Lys Ile Gly Phe	260	265	270
Leu Gly Leu Gly Leu Met Gly Ser Gly Ile Val Ser Asn Leu Leu Lys	275	280	285
Met Gly His Thr Val Thr Val Trp Asn Arg Thr Ala Glu Lys Glu Gly	290	295	300
Ala Arg Leu Gly Arg Thr Pro Ala Glu Val Val Ser Thr Cys Asp Ile	305	310	315
Thr Phe Ala Cys Val Ser Asp Pro Lys Ala Ala Lys Asp Leu Val Leu	325	330	335
Gly Pro Ser Gly Val Leu Gln Gly Ile Arg Pro Gly Lys Cys Tyr Val	340	345	350
Asp Met Ser Thr Val Asp Ala Asp Thr Val Thr Glu Leu Ala Gln Val	355	360	365
Ile Val Ser Arg Gly Gly Arg Phe Leu Glu Ala Pro Val Ser Gly Asn	370	375	380
Gln Gln Leu Ser Asn Asp Gly Met Leu Val Ile Leu Ala Ala Gly Asp			

```

385              390              395              400
Arg Gly Leu Tyr Glu Asp Cys Ser Ser Cys Phe Gln Ala Met Gly Lys
              405              410              415
Thr Ser Phe Phe Leu Gly Glu Val Gly Asn Ala Ala Lys Met Met Leu
              420              425              430
Ile Val Asn Met Val Gln Gly Ser Phe Met Ala Thr Ile Ala Glu Gly
              435              440              445
Leu Thr Leu Ala Gln Val Thr Gly Gln Ser Gln Gln Thr Leu Leu Asp
              450              455              460
Ile Leu Asn Gln Gly Gln Leu Ala Ser Ile Phe Leu Asp Gln Lys Cys
465              470              475              480
Gln Asn Ile Leu Gln Gly Asn Phe Lys Pro Asp Phe Tyr Leu Lys Tyr
              485              490              495
Ile Gln Lys Asp Leu Arg Leu Ala Ile Ala Leu Gly Asp Ala Val Asn
              500              505              510
His Pro Thr Pro Met Ala Ala Ala Ala Asn Glu Val Tyr Lys Arg Ala
              515              520              525
Lys Ala Leu Asp Gln Ser Asp Asn Asp Met Ser Ala Val Tyr Arg Ala
              530              535              540
Tyr Ile His
545

```

```

<210> 138
<211> 1097
<212> DNA
<213> Homo sapiens

```

```

<400> 138
attggttcaa gaagaaaatt tgagcccagg cacccaaaca ccttcaaag ataaagcaag 60
tatgttgcaa gaatactcca aatacctcca acaggctttt gaaaaatcca ctaatgcaag 120
ttttactctt ggacacgggt tccaatttgt cagtttgtct tcacctctcc acaaccacac 180
tttggtttcca gaaaaacaaa tatacactac gtctcctttg gagtgtggtt tcggccaatc 240
tgttacctca gtgttgccat cttcattgcc aaagcctcct tttgggatgt tgtttggatc 300
tcagccaggt ctttatttgt ctgctttgga tgctacacat cagcagttga caccttccca 360
ggagctggat gatctgatag attctcagaa gaacttagag acttcatcag ccttcagtc 420
ctcatctcag aaattgacta gccagaagga acagaaaaac ttagagtctt caacaggctt 480
tcagattcca tctcaggagt tagctagcca gatagatcct cagaaagaca tagagcctag 540
aacaacgtat cagattgaga actttgcaca agcgtttggt tctcagttta agtcgggcag 600
cagggtgcca atgaccttta tctaactc taatggagaa gtggaccata gagtaaggac 660
ttcagtgtca gatttctcag ggtatacaaa tatgatgtct gatgtaagt agccatgtag 720
tacaagagta aagacaccca ccagccagag ttacaggtaa ggtcccaaaa gtggccaggc 780
tggaggtttt ttaatgtaat tttgttttat tttgagaaca ctgccattgg aatgttttta 840
cacgatccta ttaagaataa tgtgatgccc tttcaatgca acttttcata ttagtttat 900
tttgtagcgt ttaatttagc tctgtttgta ttatgatttt taatcaaaat caatagatta 960
aaaatagttt gacattcaaa gtgacaatgt ttagcaatca aatttacatg tatagattgt 1020
cagggaatag cccaatgtt ttaaacgcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1080
aaaaaaaaaa aaaaaaa
1097

```

```

<210> 139

```

<211> 232
 <212> PRT
 <213> Homo sapiens

<400> 139
 Met Leu Gln Glu Tyr Ser Lys Tyr Leu Gln Gln Ala Phe Glu Lys Ser
 1 5 10 15
 Thr Asn Ala Ser Phe Thr Leu Gly His Gly Phe Gln Phe Val Ser Leu
 20 25 30
 Ser Ser Pro Leu His Asn His Thr Leu Phe Pro Glu Lys Gln Ile Tyr
 35 40 45
 Thr Thr Ser Pro Leu Glu Cys Gly Phe Gly Gln Ser Val Thr Ser Val
 50 55 60
 Leu Pro Ser Ser Leu Pro Lys Pro Pro Phe Gly Met Leu Phe Gly Ser
 65 70 75 80
 Gln Pro Gly Leu Tyr Leu Ser Ala Leu Asp Ala Thr His Gln Gln Leu
 85 90 95
 Thr Pro Ser Gln Glu Leu Asp Asp Leu Ile Asp Ser Gln Lys Asn Leu
 100 105 110
 Glu Thr Ser Ser Ala Phe Gln Ser Ser Ser Gln Lys Leu Thr Ser Gln
 115 120 125
 Lys Glu Gln Lys Asn Leu Glu Ser Ser Thr Gly Phe Gln Ile Pro Ser
 130 135 140
 Gln Glu Leu Ala Ser Gln Ile Asp Pro Gln Lys Asp Ile Glu Pro Arg
 145 150 155 160
 Thr Thr Tyr Gln Ile Glu Asn Phe Ala Gln Ala Phe Gly Ser Gln Phe
 165 170 175
 Lys Ser Gly Ser Arg Val Pro Met Thr Phe Ile Thr Asn Ser Asn Gly
 180 185 190
 Glu Val Asp His Arg Val Arg Thr Ser Val Ser Asp Phe Ser Gly Tyr
 195 200 205
 Thr Asn Met Met Ser Asp Val Ser Glu Pro Cys Ser Thr Arg Val Lys
 210 215 220
 Thr Pro Thr Ser Gln Ser Tyr Arg
 225 230

<210> 140
 <211> 775
 <212> DNA
 <213> Homo sapiens

<400> 140
 gtgtcacata ccactcttgt aggtgtcctc aataatcccc ttttcccaca aaatacacag 60
 ggtgtattat ctttctcttt attcaccccc actttgctga actgaagtta attacatagc 120
 ctttcttcta acctccttag taatgaacct tcacataaag tgtatttaca gcgtctgtgg 180

```

tagccagccc ttcctcctct actttctagg aggggatagc caataactag gaatttaagt 240
acagattttt tttttctttg aaataaatgg ccagagtttc tccatttttag aattttgttg 300
tcctccttaa tcatctgctt acctagtcac tactcaatct gcagaaactt cataaaggaa 360
aagtgtctga ttgtttttac aaataacagt ttgtagggaa aatatgacaa acctcaacta 420
tgggagttgt ccacaataca aaattttgaa aaaacattac atagtataa tatcatactt 480
ggttgtagg cttgttgctt cccacatca gaggcattca atgatttatc ttttgtaatt 540
gctgtgaact tttttaata agccatttag tgtgaaattg tcatgtatca aatggctatt 600
ggaaatggac tttactcaat tttaattcca ctgcactcta gccggagtga cagagtaaga 660
ctctgtctca aaaataaata aataaataaa taaataaata aataaataaa taaataaaaa 720
ataataatac aagttttcat aaggaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 775

```

<210> 141

<211> 44

<212> PRT

<213> Homo sapiens

<400> 141

```

Met Thr Asn Leu Asn Tyr Gly Ser Cys Pro Gln Tyr Lys Ile Leu Lys
  1             5             10            15

```

```

Lys His Tyr Ile Val Ile Ile Ser Tyr Leu Val Val Arg Leu Val Ala
      20             25             30

```

```

Ser Pro His Gln Arg His Leu Met Ile Tyr Leu Leu
      35             40

```

<210> 142

<211> 2060

<212> DNA

<213> Homo sapiens

<400> 142

```

gaaaaagaag acaaagctca ctttcaggcg gaggtgcagc acctgcgaga ggacaacctg 60
aggctacagg aggagtccca gaacgcctcg gacaagctga agaagttcac agaatgggtc 120
ttcaacacca tagacatgag ctagggaagg ctgaggagga caggagaagg gccagacac 180
tcctccagtg gagtgcctcg cagcccttat tcctccata gaaagcatcc tcagagcacc 240
ttccctgggt tcctactctg ccccttttcg gggagtgcac aacacaatag ttgcagatca 300
acaatcatca cctgcctttt gttagaaaaga aaaacaaaaa aagtaataaa aaatttttaa 360
cagtaaaata aaagtttaac tgctaaaatg tgaatgtctt tatttttttg cacaatatct 420
ttatctgtta tgtatttaag aagaaactgg gccttgagcc agggcgcccc ctggcccatc 480
cgcctctatt cccatcagct ttcttatcaa cttcaggtaa cccaagcttt cccttggtat 540
tctaacaagt atcattatcc cttagaaaag aatgttttta taacttggtt ggggagtaga 600
gagggatatt tccttacctt cttccctaaa atgcctggag agggagtgc tttgagaaaa 660
tgcctacctc ccttgaatga ctctgtcatg agctagtgtc gtctgtacct gtccctccaga 720
gatcagcagg accggagtta aatatttaac agcaagtctg taaaccagag cagctctgac 780
agtgcctgca ggccacaccc cttctcagtc ctgcattgtg aggtcatttc ctgctctctc 840
ctttcccccga gaagatggtc ccacttgtgc tgcaagactc ttttttggtt ggcttaattg 900
agccccacta aattggaatc aatctctctt acagcttcct ggctccaaac attaatgat 960
ttcagaatcc ccccaacta aaacctatc tgtctgcatt ttgaatgcat ttgggtcaaa 1020
agtatacgtt ttaaagattt ttaaagataa aaatgtggca caactgggtt ttttagcttg 1080
ctgaaaaatga ccatatctct aaattaatct ttctctccag agcaagactt caccagtatt 1140
tgtaacttag agaagctaag tgaatgttta attgtgaatt ttaatcattg cttgttagga 1200
ataatgactg tgatactaga atgggctttt gaaacctgca tgtcccagtg tgaaatttca 1260
gcacggcatt ttctgcatcc ttctatggcc atccaaagga ttccgctgca gaaattattg 1320
atgtgtctatt tttgtgtctt tgtgatgcag gctgcttttg gccctgggt cactcttcca 1380
aggctgtctg agagcacaga gacatggggc tggccagtgt tgactgacct gaggagaccc 1440
ctttgtttgt tgcttccata actgtcacta aaccgacccc tctgcccctt cagtggcaac 1500
tctggtctaa gggaacattc agcactctag cgcatctga ttggaagttc cctcacccaa 1560
gtaatctcaa ttccttctc tctccatccc tgaaagaaac aggatggatt ttcctctctt 1620

```

```

ctccctgcta cattcactac cagattttta tgctacagtt tcattcttga ttgtgatttc 1680
tccatggaat tttttttttc tggtagacatt tctatcatgg aaataggaag atttcggagt 1740
gctttgtgaa gatttcaatt gtctgtctct tctctctctt gacttgtagt aaggagattg 1800
tacattgcct gatattctct tgtaaatgag aaatattgct aacatccaag cattctgaag 1860
tcttgcttat cctcttgagt ttagttctca ttttgtttta cattttgttt ggggacttgg 1920
ggcaagctat ttattagagt tttgcaacag agttcttggt tgaagcctct aaagactacc 1980
tgtaaaattc aaagaataaa attcatttta aacgctcttt taaaaaaaaa aaaaaaaaaa 2040
aaaaaaaaa aaaaaaaaaa
2060

```

<210> 143
 <211> 62
 <212> PRT
 <213> Homo sapiens

```

<400> 143
Met Thr Val Ile Leu Glu Trp Ala Phe Glu Thr Cys Met Ser Gln Cys
  1             5             10             15
Glu Ile Ser Ala Arg His Phe Leu His Pro Phe Met Ala Ile Gln Arg
          20             25             30
Ile Pro Leu Gln Lys Leu Leu Met Cys Tyr Phe Cys Cys Leu Val Met
          35             40             45
Gln Ala Ala Leu Gly Pro Trp Val Thr Leu Pro Arg Leu Leu
          50             55             60

```

<210> 144
 <211> 1160
 <212> DNA
 <213> Homo sapiens

```

<400> 144
gattattttc agtaggcaga catctaactg gaattcttgc cttgttgccc aggctggagt 60
gtaatggcac aatctcggct tactgcaacc tctgcctcct ggattcaagt gattctctcg 120
cctcagcctc ccaagtagct gggattacag ccctgaaaac cactcgcttg cagagcgctg 180
gatcagcaat gcctactagt tcttcattca aacaccggat taaagagcag gaagactaca 240
tccgagattg gactgctcat cgagaagaga tagccaggat cagccaagat cttgctctca 300
ttgctcggga gatcaacgat gtagcaggag agatagattc agtgacttca tcaggcactg 360
cccctagtac cacagtaagc actgctgcca ccaccctgg ctctgccata gacactagag 420
aagagttggg tgatcgtggt tttgatgaaa gcctcaactt ccaaaagatt cctccattag 480
ttcattccaa aacaccagaa ggaaacaacg gtcgatctgg tgatccaaga cctcaagcag 540
cagagcctcc cgatcactta acaattacaa ggccggagaac ctggagcagg gatgaagtca 600
tgaggagataa tctgctgctg tcatccgtct ttcagttctc targaagata agacaatcta 660
tagataagac agctggaaaag atcagaatat tatttaaaga caaagatcgg aattgggatg 720
acatagaaaag caaattaaga gccgaaagtg aagtccttat tgtgaaaacc tcgagcatgg 780
agatttcttc tatcttacag gaactgaaaa gagtagaaaa gcagctacaa gcaatcaatg 840
ctatgattga tcttgatgga actttggagg ctctgaacaa catgggattt cccagtgcga 900
tggtgccatc tccaccgaaa cagaagtcca gccctgtgaa taaccaccac agcccgggtc 960
agacaccaac acttggccaa ccagaagcta gggctcttca tctgctgctg gtttcagccg 1020
cagctgaatt tgagaatgct gaatctgagg ctgatttcag tatacatttc aatagagtca 1080
accctgatgg ggaagaggaa gatgttacag taacataaat gactttctct tgattgttga 1140
aaaaaaaaa aaaaaaaaaa
1160

```

<210> 145
 <211> 309
 <212> PRT
 <213> Homo sapiens

<220>

<221> UNSURE

<222> (152)

<400> 145

```

Met Pro Thr Ser Ser Ser Phe Lys His Arg Ile Lys Glu Gln Glu Asp
 1           5           10           15

Tyr Ile Arg Asp Trp Thr Ala His Arg Glu Glu Ile Ala Arg Ile Ser
          20           25           30

Gln Asp Leu Ala Leu Ile Ala Arg Glu Ile Asn Asp Val Ala Gly Glu
          35           40           45

Ile Asp Ser Val Thr Ser Ser Gly Thr Ala Pro Ser Thr Thr Val Ser
          50           55           60

Thr Ala Ala Thr Thr Pro Gly Ser Ala Ile Asp Thr Arg Glu Glu Leu
          65           70           75           80

Val Asp Arg Val Phe Asp Glu Ser Leu Asn Phe Gln Lys Ile Pro Pro
          85           90           95

Leu Val His Ser Lys Thr Pro Glu Gly Asn Asn Gly Arg Ser Gly Asp
          100           105           110

Pro Arg Pro Gln Ala Ala Glu Pro Pro Asp His Leu Thr Ile Thr Arg
          115           120           125

Arg Arg Thr Trp Ser Arg Asp Glu Val Met Gly Asp Asn Leu Leu Leu
          130           135           140

Ser Ser Val Phe Gln Phe Ser Xaa Lys Ile Arg Gln Ser Ile Asp Lys
          145           150           155           160

Thr Ala Gly Lys Ile Arg Ile Leu Phe Lys Asp Lys Asp Arg Asn Trp
          165           170           175

Asp Asp Ile Glu Ser Lys Leu Arg Ala Glu Ser Glu Val Pro Ile Val
          180           185           190

Lys Thr Ser Ser Met Glu Ile Ser Ser Ile Leu Gln Glu Leu Lys Arg
          195           200           205

Val Glu Lys Gln Leu Gln Ala Ile Asn Ala Met Ile Asp Pro Asp Gly
          210           215           220

Thr Leu Glu Ala Leu Asn Asn Met Gly Phe Pro Ser Ala Met Leu Pro
          225           230           235           240

Ser Pro Pro Lys Gln Lys Ser Ser Pro Val Asn Asn His His Ser Pro
          245           250           255

Gly Gln Thr Pro Thr Leu Gly Gln Pro Glu Ala Arg Ala Leu His Pro
          260           265           270

Ala Ala Val Ser Ala Ala Ala Glu Phe Glu Asn Ala Glu Ser Glu Ala
          275           280           285

Asp Phe Ser Ile His Phe Asn Arg Val Asn Pro Asp Gly Glu Glu Glu

```

290

295

300

Asp Val Thr Val Thr
305

<210> 146

<211> 1536

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (317)

<400> 146

```

gaagagagaa aatcagcctg tctgctctct ccttggctca acaaggcctc taacagtctt 60
ctgtcctcta ttctgcacac ggcataatttg ggaacgagaa acaaaagttt tcccaaatga 120
agagaactca cttgtttatt gtggggattt attttctgtc ctcttgagg gcagaagagg 180
ggcttaattt ccccatatat gatgggaagg accgagtggg aagtctttcc gagaagaact 240
tcaagcaggt tttaaagaaa tatgacttgc tttgcctcta ctaccatgag ccggtgtctt 300
cagataaggt cagcnaaaa cagttccaac tgaaagaaaat cgtgcttgag cttgtggccc 360
acgtccttga acataaagct ataggctttg tgatggttga tgccaagaaa gaagccaagc 420
ttgccaagaa actgggtttt gatgaagaag gaagcctgta tattcttaag ggtgatcgca 480
caatagagtt tgatggcgag tttgcagctg atgtcttggg ggagttcctc ttggatctaa 540
ttgaagaccc agtgagatc atcagcagca aactggaagt ccaagccttc gaacgcattg 600
aagactacat caaactcatt ggctttttca agagtgagga ctcaagaatac tacaaggctt 660
ttgaagaagc agctgaacac ttccagcctt acatcaaatt ctttgccacc tttgacaaag 720
gggttgcaaa gaaattatct ttgaagatga atgaggttga cttctatgag ccatttatgg 780
atgagcccat tgccatcccc aacaaacctt acacagaaga ggagctggtg gagtttgtga 840
aggaacacca aaggtgctg agatggcatg tgggggctgg gggcctgggg tctggggaat 900
ggagaggagc ctctctgtgc taacatttca gacctgcaa gagcaacaac ctagttagta 960
ccccagcagt acagaactca gtatgatggc tttgttgatc agtaatgact agcagggatg 1020
ttattacttc tgaatctaag tctgcacctg caagcagagt ttgataaatc cctcagtcag 1080
caaatccctt caaagccagg gcaagatata aataaaattc tatacttagga atgagagcaa 1140
tttagtgaaa gttcccatat accaataaacc atgcccagtg ctttagggaa actattttat 1200
ctaacttcca accttaggga gtaattatta ttatcccaat tttacagatc aaggaatttg 1260
actcaatagt taagtaactt agccaaggat gaacactcta tgcatagaac ttctgggaga 1320
gaaatgcttg ataccactta gtgtagctcc agcatggatc agcaaacttt ttctgtaaag 1380
aacaaaatgg taaatatttc aggttctgtg ggccagatgg cgtctgtagc aactacttaa 1440
ctgctggctgt ggcagtaaa cagccatgga tcattgtataa acaaatgggt gtggctgtgt 1500
accagtaaaa gtttatattg gaaaaaaaa aaaaaa 1536

```

<210> 147

<211> 268

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (67)

<400> 147

```

Met Lys Arg Thr His Leu Phe Ile Val Gly Ile Tyr Phe Leu Ser Ser
 1           5           10          15

```

```

Cys Arg Ala Glu Glu Gly Leu Asn Phe Pro Thr Tyr Asp Gly Lys Asp
      20           25           30

```

```

Arg Val Val Ser Leu Ser Glu Lys Asn Phe Lys Gln Val Leu Lys Lys

```


35 40 45
 Tyr Asp Leu Leu Cys Leu Tyr Tyr His Glu Pro Val Ser Ser Asp Lys
 50 55 60
 Val Thr Xaa Lys Gln Phe Gln Leu Lys Glu Ile Val Leu Glu Leu Val
 65 70 75 80
 Ala His Val Leu Glu His Lys Ala Ile Gly Phe Val Met Val Asp Ala
 85 90 95
 Lys Lys Glu Ala Lys Leu Ala Lys Lys Leu Gly Phe Asp Glu Glu Gly
 100 105 110
 Ser Leu Tyr Ile Leu Lys Gly Asp Arg Thr Ile Glu Phe Asp Gly Glu
 115 120 125
 Phe Ala Ala Asp Val Leu Val Glu Phe Leu Leu Asp Leu Ile Glu Asp
 130 135 140
 Pro Val Glu Ile Ile Ser Ser Lys Leu Glu Val Gln Ala Phe Glu Arg
 145 150 155 160
 Ile Glu Asp Tyr Ile Lys Leu Ile Gly Phe Phe Lys Ser Glu Asp Ser
 165 170 175
 Glu Tyr Tyr Lys Ala Phe Glu Glu Ala Ala Glu His Phe Gln Pro Tyr
 180 185 190
 Ile Lys Phe Phe Ala Thr Phe Asp Lys Gly Val Ala Lys Lys Leu Ser
 195 200 205
 Leu Lys Met Asn Glu Val Asp Phe Tyr Glu Pro Phe Met Asp Glu Pro
 210 215 220
 Ile Ala Ile Pro Asn Lys Pro Tyr Thr Glu Glu Glu Leu Val Glu Phe
 225 230 235 240
 Val Lys Glu His Gln Arg Cys Leu Arg Trp His Val Gly Ala Gly Gly
 245 250 255
 Leu Gly Ser Gly Glu Trp Arg Gly Ala Ser Leu Cys
 260 265

<210> 148

<211> 1009

<212> DNA

<213> Homo sapiens

<400> 148

gaattcggcc ttcatgcgcc tgcaggaaag aatctgacat catcacactg tgttttcctt 60
 aacttgacag gaagtcaact tcaagcagat tgacttgaaa cgggatctca tttaggaagc 120
 ataagtgtcc aatcaaaaac tgtgtatttt tttaaatttg gaaaatactc aagttccagt 180
 tgcttatcat tctccttcac tttctgaaaa cctggcaatc ccatgtggac ttctggtaga 240
 atgagcaatg caaagaactg gcttggaactt ggcattgtcct tgtacttctg ggggctgatg 300
 gaccttacga ccaccgttct ctcggacacc ccaacaccac aaggtgaatt agaagcactc 360
 ctgtcagaca agccacagtc acatcagcgg accaagarga gctgggtttg gaaccagttt 420
 ttcgtttctg aagagtacac tgggaccgac cctttgtatg tcggcaaggt aagaaatgcc 480
 aagtagaaat gaccggggtg gtggatattg aaattgaata tgaattgagt atcaaagttg 540

```

acctagcctt tatytgagac ctgagaaaaa ctagaacaag tggtagctta cttgacacct 600
agctaaaaatg taacttctgc tttgtcagag accagtctga aaggaaagat ttatttcctt 660
gtccatgtct ctggtatgaa tgggaaaaag tgggaattgg gatttggagg aaaaggctca 720
gacctgcaa gagctattca agtcctaaaa gaggcagcag cagctgtctg ggaatgacag 780
aatgggggag agggaaaactt ggaaatacaa gaagagtaca gagttttttg ctttgtgttt 840
ttatgggggt tttttcagag catttcqaga ggtattgcct gaggtgcaaa ttaaatgaaa 900
aaaataaaaa taaaacattt ttcatttcag aagatcctag catgtgcttt aggatagttg 960
gagacaataa atatatttat aaatgttaaa aaaaaaaaaa aaaaaaaaaa 1009

```

<210> 149

<211> 87

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (59)

<400> 149

```

Met Trp Thr Ser Gly Arg Met Ser Asn Ala Lys Asn Trp Leu Gly Leu
  1              5              10             15

```

```

Gly Met Ser Leu Tyr Phe Trp Gly Leu Met Asp Leu Thr Thr Thr Val
      20              25              30

```

```

Leu Ser Asp Thr Pro Thr Pro Gln Gly Glu Leu Glu Ala Leu Leu Ser
      35              40              45

```

```

Asp Lys Pro Gln Ser His Gln Arg Thr Lys Xaa Ser Trp Val Trp Asn
      50              55              60

```

```

Gln Phe Phe Val Leu Glu Glu Tyr Thr Gly Thr Asp Pro Leu Tyr Val
      65              70              75             80

```

```

Gly Lys Val Arg Asn Ala Lys
      85

```

<210> 150

<211> 2546

<212> DNA

<213> Homo sapiens

<400> 150

```

aaaagaaacc aaggaaattt gtatgataag gcaggtaaag tgaggaaaca tgcaactgaa 60
caggaaaaaa ctgaagaggg attaggccct aatataaaaa gcattgtcac catgttgatg 120
ctgatgctat tgatgatgtt tgctgtccac tgtacctggg tcacaagcaa tgcctactct 180
agtccaagtg tagtcctggc ctcatacaat catgatggca ccaggaatat cttagatgat 240
tttagagaag cttacttttg gctaaggcaa aatacagatg aacatgcacg agtaatgtct 300
tggtgggatt atggctatca gatagctgga atggctaata gaactacgtt ggtggataat 360
aacacctgga ataacagcca catagcactg gtgggaaaag ctatgtcttc taatgaaaca 420
gcagcctata aaatcatgag gactctagat gtagattatg ttttggttat ttttgagggg 480
gttattggct attctgggtg tgatatcaac aaatttctct ggatgggttag gatagctgaa 540
ggagaacatc ccaaagacat tcgggaaagt gactatttta cccacaggg agaattccgt 600
gtagacaaaag caggatcccc tactttgttg aattgcctta tgtataaaat gtcatactac 660
agatttgag aaatgcagct ggattttcgt acacccccag gttttgaccg aacacgtaat 720
gctgagattg gaaataagga cattaaattc aaacatttgg aagaagcctt tacatcagaa 780
cactggcctt ttaggatata taaagtaaaa gcacctgata acagggagac attagatcac 840
aaacctcgag tcaccaacat tttcccaaaa cagaagtatt tgtcaaagaa gactaccaaa 900
aggaagcgtg gctacattaa aaataagctg gtttttaaga aaggcaagaa aatatctaag 960

```

```

aagactgttt aaatgcactg ttctggttcc taacttgaag cagttgtcct tgtgagaacc 1020
ggctctttgcc tttagctcat gtcgtgtttc acagcaaaga gggtagacaa ccatcactgg 1080
tccagggttaa tgtacaaaat ttcttggaac tgcctgatta aaaaaataaa attggcttgt 1140
tgagaacagc tgttttcgat ttctaattgt aagcaagaca gagcactgct gtaaattgtct 1200
agcagcagat ttttttttta ttggtacata ttatccttca aatctgagaa tttggactaa 1260
ctgcacaaaa gaacctctta atttgggtccc tggcacatgc atacttgtca atgtttttat 1320
tctctttacaa gacctgcatt ttatttgaat taccgaata gcaatatgta aaatacaagt 1380
gacaaaatgt gatgagagct tcttgaaccg gtaaaactagt acaggtctga gaaagacata 1440
ttaagaagaat cattatactt ccttgaatta tttttatctt catgtttctc taatgcaaag 1500
aatgtttcat caaatgtata ttttctgttg cttactgttt gctctgagaa gaagctgctg 1560
tttcaaagat ggacctctga gtagctaatt gattcaagta gtttttttat gttgacacat 1620
tattactgct gtttagcagtc gttttcacca ggtacttaca gagcagattt catacatcat 1680
tcattcaagg gctaaattta ttttttttgg aaatcatggc aactacacag gatgttgctt 1740
accaggacg agtttttggt tcttagtact gaagttagca ctatgtttac atgcaaaaaga 1800
ttaaggaaaa aacctttaa gtggacaggt atccaaagtt cattttctgt gactcatcaa 1860
agtgaacaaa gacttgaac aactttgcct ggactttttt cattttacaa cagttcatcc 1920
attcacaatg attttgttct ctgctccata ttttttaatc ccttaagcat ttgatgaaac 1980
actctttagt gctatatgca ttttcttact tttgttaaaa atgtgacaat tgtcaaaaaa 2040
tgcactaaaa tgtaaatgga gattgaacaa gttcactttc cagcttatag gcaactttat 2100
acagacttga acattttctc cagttgttta gtaaaagtga aagagaaagg gtttttcctg 2160
ccacaggata taactttttt ttatataaca agcataaacac accactgctt ttggtggaaa 2220
agtgcagaat agtatgtacc ttttatgaag aaaaatgtaa tttacaatat tcagtgagaa 2280
tgttactgct gattttcttt tccaaggtgt agaatactct ttgatttata gaattcattt 2340
ttgaccacaga tgaagggtcc tttacagaac aataaaatgg ctgaacattt tcacaaatag 2400
agtgtaacga agtctggatt tctgatacct tgtcatttgg gggattttat tttactttgt 2460
tgcttttaaaa ttcaatgcag agaagttgtt gactgtaggg gaaataaagt taattcaaat 2520
tttgaaaaaa aaaaaaaaaa aaaaaa 2546

```

<210> 151

<211> 286

<212> PRT

<213> Homo sapiens

<400> 151

```

Met Leu Met Leu Met Leu Leu Met Met Phe Ala Val His Cys Thr Trp
  1                      5                      10                      15

```

```

Val Thr Ser Asn Ala Tyr Ser Ser Pro Ser Val Val Leu Ala Ser Tyr
      20                      25                      30

```

```

Asn His Asp Gly Thr Arg Asn Ile Leu Asp Asp Phe Arg Glu Ala Tyr
      35                      40                      45

```

```

Phe Trp Leu Arg Gln Asn Thr Asp Glu His Ala Arg Val Met Ser Trp
      50                      55                      60

```

```

Trp Asp Tyr Gly Tyr Gln Ile Ala Gly Met Ala Asn Arg Thr Thr Leu
      65                      70                      75                      80

```

```

Val Asp Asn Asn Thr Trp Asn Asn Ser His Ile Ala Leu Val Gly Lys
      85                      90                      95

```

```

Ala Met Ser Ser Asn Glu Thr Ala Ala Tyr Lys Ile Met Arg Thr Leu
      100                      105                      110

```

```

Asp Val Asp Tyr Val Leu Val Ile Phe Gly Gly Val Ile Gly Tyr Ser
      115                      120                      125

```

```

Gly Asp Asp Ile Asn Lys Phe Leu Trp Met Val Arg Ile Ala Glu Gly
      130                      135                      140

```

Glu His Pro Lys Asp Ile Arg Glu Ser Asp Tyr Phe Thr Pro Gln Gly
 145 150 155 160
 Glu Phe Arg Val Asp Lys Ala Gly Ser Pro Thr Leu Leu Asn Cys Leu
 165 170 175
 Met Tyr Lys Met Ser Tyr Tyr Arg Phe Gly Glu Met Gln Leu Asp Phe
 180 185 190
 Arg Thr Pro Pro Gly Phe Asp Arg Thr Arg Asn Ala Glu Ile Gly Asn
 195 200 205
 Lys Asp Ile Lys Phe Lys His Leu Glu Glu Ala Phe Thr Ser Glu His
 210 215 220
 Trp Leu Val Arg Ile Tyr Lys Val Lys Ala Pro Asp Asn Arg Glu Thr
 225 230 235 240
 Leu Asp His Lys Pro Arg Val Thr Asn Ile Phe Pro Lys Gln Lys Tyr
 245 250 255
 Leu Ser Lys Lys Thr Thr Lys Arg Lys Arg Gly Tyr Ile Lys Asn Lys
 260 265 270
 Leu Val Phe Lys Lys Gly Lys Lys Ile Ser Lys Lys Thr Val
 275 280 285

<210> 152

<211> 4061

<212> DNA

<213> Homo sapiens

<400> 152

aagcaattga agaaattgca gcaggatgtg atggaaatga aaaaaacaaa ggctcgcta 60
 atgaaacaaa tgaaagaaga acaagagaaa gccagactga ctgagtctag aagaaacaga 120
 gagattgctc agttgaaaaa ggatcaacgt aaaagagatc atcmacttag actcttgaa 180
 gccccaaaaa gaaaccaaga agtggttcta cgtcgcaaaa ctgaagaggt tacggctctt 240
 cgtcggcaag taagacccat gtcagataaa gtggctggga aagttactcg gaagctgagt 300
 tcatctgatg cacctgctca ggacacaggt tccagtgcag ctgctgtcga aacagatgca 360
 tcaaggacag gagcccagca gaaaatgaga attcctgtgg cgagagtcca ggcttacca 420
 acgcccggcaa caaatggaaa caggaaaaaa tatcagagga aaggattgac tggccgagt 480
 tttatttcca agacagctcg catgaagtgg cagctccttg agcgcagggt cacagacatc 540
 atcatgcaga agatgaccat ttccaacatg gaggcagata tgaatagact cctcaagcaa 600
 cgggaggaac tcacaaaaag acgagagaaa ctttcaaaaa gaaggagaa gatagtcaag 660
 gagaatggag agggagataa aaatgtggct aatatcaatg aagagatgga gtcactgact 720
 gctaatatcg attacatcaa tgacagtatt tctgattgtc aggccaacat aatgcagatg 780
 gaagaagcaa aggaagaagg tgagacattg gatgttactg cagtcattaa tgctgcacc 840
 cttacagaag cccgatacct gctagatcac ttctgtcaa tgggcatcaa taagggtctt 900
 caggctgccc agaaagaggc tcaaattaaa gtactggaag gtcgactcaa acaaacagaa 960
 ataaccagtg ctacccaaaa ccagctctta ttccatattg tgaagagaa ggcagaatta 1020
 aatcctgagc tagatgcttt actaggccat gctttacaag atctagatag cgtaccatta 1080
 gaaaatgtag aggatagtag tgatgaggat gtccttttaa acagcccagg atcagaagga 1140
 agcacgctgt cttcagatct catgaagctt tgtggtgaag tgaaccta gaacaaggcc 1200
 cgaaggagaa ccaccactca gatggaattg ctgtatgcag atagcagtga actagcttca 1260
 gacactagta caggagatgc ctcttgctt ggccctctca cacctgttgc agaagggcaa 1320
 gagattggaa tgaatacaga gacaagtgg acttctgcta gggaaaaa gctctctccc 1380
 ccacctggct taccttctaa gataggcagc atttccaggc agtcattct atcagaaaaa 1440
 aaaattccag agccttctcc tgtaacaagg agaaaggcat atgagaaaagc agaaaaatca 1500

```

aaggccaagg aacaaaagca ctcagattct ggaacttcag aggctagtct ttcacctcct 1560
tcttccccac caagccggcc ccgtaatgaa ctgaatgttt ttaatcgtct tactgtttct 1620
cagggaaaaca catcagttca gcaggataag tctgatgaaa gtgactcctc tctctcggag 1680
gtacacagca gatcctccag aaggggcata atcaacccat ttcctgcttc aaaaggaatc 1740
agagcttttc cacttcagtg tattcacata gctgaagggc atacaaaagc tgtgctctgt 1800
gtggattcta ctgatgatct cctcttcact ggatcaaaaag atcgtacttg taaagtatgg 1860
aatctggtga ctgggcagga aataatgtca ctgggggggc atcccaacaa tgtcgtgtct 1920
gtaaaatact gtaattatac cagtttggtc ttactgtat caacatctta tattaagggtg 1980
tgggatatca gagattcagc aaagtgcatt cgaacactaa cgtcttcagg tcaagtact 2040
cttgagagatg cttgttctgc aagtaccagt cgaacagtag ctattccttc tggagagaa 2100
cagatcaatc aaattgcctt aaacccaact ggcaccttcc tctatgtctg ttctggaaat 2160
gctgtcagga tgtgggatct taaaagggtt cagtctacag gaaagttaac aggacacct 2220
ggccctgtta tgtgccttac tgtggatcag atttccagtg gacaagatct aatcatcact 2280
ggctccaagg atcattacat caaaatgttt gatgttacag aaggagctct tgggactgtg 2340
agtccacccc acaattttga acccctcat tatgatggca tagaagcact aaccattcaa 2400
ggggataacc tatttagtgg gtctagagat aatggaatca agaaatggga cttaactcaa 2460
aaagaccttc ttcagcaagt tccaaatgca cataaggatt gggctctgtg cctgggagt 2520
gtgccagacc acccagtttt gctcagtggc tgcagagggg gcattttgaa agtctggaac 2580
atggatactt ttatgccagt gggagagatg aagggtcatg atagtctat caatgccata 2640
tgtgttaatt ccaccacat tttactgca gctgatgac gaactgtgag aatttggaag 2700
gctcgcaatt tgcaagatgg tcagatctct gacacaggag atctggggga agatattgcc 2760
agtaattaaa catgaatgaa gatagggtgt aaactgaatg ctgtgataat actctgtatt 2820
ctttatggaa aatgttgtcc tgtacttact aggcataacg tatgaatcgg attaactgga 2880
aaatatactt gaattcaact gctgactata aatgggtatc taataaaaatt gtgtactatc 2940
ctgtgtgctt agttttaaga tcaaccaata gatacatatc ctacaattga tatattgctt 3000
tattcacact tttattgtgg ctgaattttt gtgcctatct ataaaaacaca ctttcaaatt 3060
atttgaatta ccaagacgtc tgcctttgtg acagtcagaa aacacacctg gaatacgtg 3120
cagcccacca ttaactcatt catgtagttt attcaagtga tttatgtatt taaactaaat 3180
attgaaaaatg ttagtcaaat tgtgggttgc ttgtcaggta tttatatcag tctgtagtgg 3240
attcccaaat ttcaaagctc ttttaatgta atggacaaaa ataagatatg agaataattat 3300
tgatgaattt tcataagggtg gaattgatct taactacta acagagaagg gtagacagt 3360
tgtgttaaat gtggcattt acttgattg accaaagtgt tgcagctcta ctatattctg 3420
tgcctcaggac taaaatgctg ttaatttttt tttttttttt tccagtgtg tgcataatt 3480
ctgtgatggg aaacattgtt gatgtcctaa cagaaatata ttttgatcta ttttctatg 3540
gagttgtttc tattatgacc atttaatttt gtttttattt aatagtagta tttccttccc 3600
ttttatctaa ttttttatat gctgctaaat atattttaaa tatactatgt ttgcgaacct 3660
tggtagctat gatgagagct attatcatct gtggtgggaa aagctatgta aataggtaga 3720
ttgtatagag agactatctt gtgttggtgc tgtatgaatt tttaaaagtt gttgactgga 3780
ttttgcaaaa ggatgtataa tttttctgtc tgctcagaat attaatgtt aaattctgca 3840
agtttaattt ttatgtatag ggtataacat ttgaaaatat tgccttatgt gattttttcc 3900
cctgaaaaata tttgcttgta aatgaaaact tagctagggc ttaaataaac atgttgctat 3960
gaaatkraaa aaaaaaaaaa aaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 4020
aaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 4061

```

<210> 153

<211> 910

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (43)

<400> 153

```

Met Lys Lys Thr Lys Val Arg Leu Met Lys Gln Met Lys Glu Glu Gln
  1             5             10             15

```

```

Glu Lys Ala Arg Leu Thr Glu Ser Arg Arg Asn Arg Glu Ile Ala Gln
      20             25             30

```

Leu Lys Lys Asp Gln Arg Lys Arg Asp His Xaa Leu Arg Leu Leu Glu
 35 40 45
 Ala Gln Lys Arg Asn Gln Glu Val Val Leu Arg Arg Lys Thr Glu Glu
 50 55 60
 Val Thr Ala Leu Arg Arg Gln Val Arg Pro Met Ser Asp Lys Val Ala
 65 70 75 80
 Gly Lys Val Thr Arg Lys Leu Ser Ser Ser Asp Ala Pro Ala Gln Asp
 85 90 95
 Thr Gly Ser Ser Ala Ala Ala Val Glu Thr Asp Ala Ser Arg Thr Gly
 100 105 110
 Ala Gln Gln Lys Met Arg Ile Pro Val Ala Arg Val Gln Ala Leu Pro
 115 120 125
 Thr Pro Ala Thr Asn Gly Asn Arg Lys Lys Tyr Gln Arg Lys Gly Leu
 130 135 140
 Thr Gly Arg Val Phe Ile Ser Lys Thr Ala Arg Met Lys Trp Gln Leu
 145 150 155 160
 Leu Glu Arg Arg Val Thr Asp Ile Ile Met Gln Lys Met Thr Ile Ser
 165 170 175
 Asn Met Glu Ala Asp Met Asn Arg Leu Leu Lys Gln Arg Glu Glu Leu
 180 185 190
 Thr Lys Arg Arg Glu Lys Leu Ser Lys Arg Arg Glu Lys Ile Val Lys
 195 200 205
 Glu Asn Gly Glu Gly Asp Lys Asn Val Ala Asn Ile Asn Glu Glu Met
 210 215 220
 Glu Ser Leu Thr Ala Asn Ile Asp Tyr Ile Asn Asp Ser Ile Ser Asp
 225 230 235 240
 Cys Gln Ala Asn Ile Met Gln Met Glu Glu Ala Lys Glu Glu Gly Glu
 245 250 255
 Thr Leu Asp Val Thr Ala Val Ile Asn Ala Cys Thr Leu Thr Glu Ala
 260 265 270
 Arg Tyr Leu Leu Asp His Phe Leu Ser Met Gly Ile Asn Lys Gly Leu
 275 280 285
 Gln Ala Ala Gln Lys Glu Ala Gln Ile Lys Val Leu Glu Gly Arg Leu
 290 295 300
 Lys Gln Thr Glu Ile Thr Ser Ala Thr Gln Asn Gln Leu Leu Phe His
 305 310 315 320
 Met Leu Lys Glu Lys Ala Glu Leu Asn Pro Glu Leu Asp Ala Leu Leu
 325 330 335
 Gly His Ala Leu Gln Asp Leu Asp Ser Val Pro Leu Glu Asn Val Glu
 340 345 350

Asp Ser Thr Asp Glu Asp Ala Pro Leu Asn Ser Pro Gly Ser Glu Gly
 355 360 365
 Ser Thr Leu Ser Ser Asp Leu Met Lys Leu Cys Gly Glu Val Lys Pro
 370 375 380
 Lys Asn Lys Ala Arg Arg Arg Thr Thr Thr Gln Met Glu Leu Leu Tyr
 385 390 395 400
 Ala Asp Ser Ser Glu Leu Ala Ser Asp Thr Ser Thr Gly Asp Ala Ser
 405 410 415
 Leu Pro Gly Pro Leu Thr Pro Val Ala Glu Gly Gln Glu Ile Gly Met
 420 425 430
 Asn Thr Glu Thr Ser Gly Thr Ser Ala Arg Glu Lys Glu Leu Ser Pro
 435 440 445
 Pro Pro Gly Leu Pro Ser Lys Ile Gly Ser Ile Ser Arg Gln Ser Ser
 450 455 460
 Leu Ser Glu Lys Lys Ile Pro Glu Pro Ser Pro Val Thr Arg Arg Lys
 465 470 475 480
 Ala Tyr Glu Lys Ala Glu Lys Ser Lys Ala Lys Glu Gln Lys His Ser
 485 490 495
 Asp Ser Gly Thr Ser Glu Ala Ser Leu Ser Pro Pro Ser Ser Pro Pro
 500 505 510
 Ser Arg Pro Arg Asn Glu Leu Asn Val Phe Asn Arg Leu Thr Val Ser
 515 520 525
 Gln Gly Asn Thr Ser Val Gln Gln Asp Lys Ser Asp Glu Ser Asp Ser
 530 535 540
 Ser Leu Ser Glu Val His Ser Arg Ser Ser Arg Arg Gly Ile Ile Asn
 545 550 555 560
 Pro Phe Pro Ala Ser Lys Gly Ile Arg Ala Phe Pro Leu Gln Cys Ile
 565 570 575
 His Ile Ala Glu Gly His Thr Lys Ala Val Leu Cys Val Asp Ser Thr
 580 585 590
 Asp Asp Leu Leu Phe Thr Gly Ser Lys Asp Arg Thr Cys Lys Val Trp
 595 600 605
 Asn Leu Val Thr Gly Gln Glu Ile Met Ser Leu Gly Gly His Pro Asn
 610 615 620
 Asn Val Val Ser Val Lys Tyr Cys Asn Tyr Thr Ser Leu Val Phe Thr
 625 630 635 640
 Val Ser Thr Ser Tyr Ile Lys Val Trp Asp Ile Arg Asp Ser Ala Lys
 645 650 655
 Cys Ile Arg Thr Leu Thr Ser Ser Gly Gln Val Thr Leu Gly Asp Ala
 660 665 670

Cys Ser Ala Ser Thr Ser Arg Thr Val Ala Ile Pro Ser Gly Glu Asn
 675 680 685
 Gln Ile Asn Gln Ile Ala Leu Asn Pro Thr Gly Thr Phe Leu Tyr Ala
 690 695 700
 Ala Ser Gly Asn Ala Val Arg Met Trp Asp Leu Lys Arg Phe Gln Ser
 705 710 715 720
 Thr Gly Lys Leu Thr Gly His Leu Gly Pro Val Met Cys Leu Thr Val
 725 730 735
 Asp Gln Ile Ser Ser Gly Gln Asp Leu Ile Ile Thr Gly Ser Lys Asp
 740 745 750
 His Tyr Ile Lys Met Phe Asp Val Thr Glu Gly Ala Leu Gly Thr Val
 755 760 765
 Ser Pro Thr His Asn Phe Glu Pro Pro His Tyr Asp Gly Ile Glu Ala
 770 775 780
 Leu Thr Ile Gln Gly Asp Asn Leu Phe Ser Gly Ser Arg Asp Asn Gly
 785 790 795 800
 Ile Lys Lys Trp Asp Leu Thr Gln Lys Asp Leu Leu Gln Gln Val Pro
 805 810 815
 Asn Ala His Lys Asp Trp Val Cys Ala Leu Gly Val Val Pro Asp His
 820 825 830
 Pro Val Leu Leu Ser Gly Cys Arg Gly Gly Ile Leu Lys Val Trp Asn
 835 840 845
 Met Asp Thr Phe Met Pro Val Gly Glu Met Lys Gly His Asp Ser Pro
 850 855 860
 Ile Asn Ala Ile Cys Val Asn Ser Thr His Ile Phe Thr Ala Ala Asp
 865 870 875 880
 Asp Arg Thr Val Arg Ile Trp Lys Ala Arg Asn Leu Gln Asp Gly Gln
 885 890 895
 Ile Ser Asp Thr Gly Asp Leu Gly Glu Asp Ile Ala Ser Asn
 900 905 910

<210> 154

<211> 372

<212> DNA

<213> Homo sapiens

<400> 154

aatcgacact gtccaacttg caatgacttc catggacttg tgcagaaaat catggagcta 60
 caggatattt tagccaaaac atcagccaag ctgtctcgag ctgaacagcg aatgaataga 120
 ttggatcagt gctattgtga aaggacttgc accatgaagg gaaccaccta ccgagaattt 180
 gagtcctgga tagacggctg taagaactgc acatgcctga atggaacat ccagtgtgaa 240
 actctaactt gcccaaatcc tgactgcccc cttaagtccg ctcttgcgta tgtggatggc 300
 aatgctgta aggaatgcaa atcgatatcc caatttcaag gacgaaccta ctttgaagga 360
 gaaagaaata ca 372

<210> 155
 <211> 761
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (108)

<220>
 <221> unsure
 <222> (191)

<220>
 <221> unsure
 <222> (268)

<220>
 <221> unsure
 <222> (299)

<400> 155
 ttctggagta tgtgttctct atgagtgcaa ggaccagacc atgaaatttg ttgagagtgc 60
 aggctgtcca gctttggatt gtccagagtc tcatcagata accttgtntc acagctgttg 120
 caaagtttgt aaaggttatg atttttgttt tgaaaggcat aactgcatgg agaattccat 180
 ctgcagaaat ntgaatgaca gggctgtttg tagctgtcga gatggtttta gggtttttcg 240
 agaggataat gcctactgtg aagacatnga tgagtgtgct gaagggcgcc attactgtng 300
 tgaaaataca atgtgtgtca acaccccggt ttcttttatg tgcactctga aaactggata 360
 catcagaatt gatgattatt catgtacaga acatgatgag tgtatcaca atcagcacag 420
 ctgtgatgaa aatgctttat gcttcaacac tgttgaggga cacaactgtg tttgcaagcc 480
 gggctataca gggaatggaa cgacatgcaa agcattttgc aaagatggct gtaggaatgg 540
 aggagcctgt attgccgcta atgtgtgtgc ctgcccacaa ggcttcactg gaccagctg 600
 tgaaacggac attgatgaat gctctgatgg ttttgttcaa tgtgacagtc gtgctaattg 660
 cattaacctg cctggatggt accactgtga gtgcagagat ggctaccatg acaatgggat 720
 gttttcacca agtggagaat cgtgtgaaga tattgatgag t 761

<210> 156
 <211> 240
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (23)

<220>
 <221> UNSURE
 <222> (51)

<220>
 <221> UNSURE
 <222> (76)

<220>
 <221> UNSURE
 <222> (87)

<400> 156
 Met Lys Phe Val Glu Ser Ser Gly Cys Pro Ala Leu Asp Cys Pro Glu
 1 5 10 15

Ser His Gln Ile Thr Leu Xaa His Ser Cys Cys Lys Val Cys Lys Gly
 20 25 30
 Tyr Asp Phe Cys Phe Glu Arg His Asn Cys Met Glu Asn Ser Ile Cys
 35 40 45
 Arg Asn Xaa Asn Asp Arg Ala Val Cys Ser Cys Arg Asp Gly Phe Arg
 50 55 60
 Val Phe Arg Glu Asp Asn Ala Tyr Cys Glu Asp Xaa Asp Glu Cys Ala
 65 70 75 80
 Glu Gly Arg His Tyr Cys Xaa Glu Asn Thr Met Cys Val Asn Thr Pro
 85 90 95
 Gly Ser Phe Met Cys Ile Cys Lys Thr Gly Tyr Ile Arg Ile Asp Asp
 100 105 110
 Tyr Ser Cys Thr Glu His Asp Glu Cys Ile Thr Asn Gln His Ser Cys
 115 120 125
 Asp Glu Asn Ala Leu Cys Phe Asn Thr Val Gly Gly His Asn Cys Val
 130 135 140
 Cys Lys Pro Gly Tyr Thr Gly Asn Gly Thr Thr Cys Lys Ala Phe Cys
 145 150 155 160
 Lys Asp Gly Cys Arg Asn Gly Gly Ala Cys Ile Ala Ala Asn Val Cys
 165 170 175
 Ala Cys Pro Gln Gly Phe Thr Gly Pro Ser Cys Glu Thr Asp Ile Asp
 180 185 190
 Glu Cys Ser Asp Gly Phe Val Gln Cys Asp Ser Arg Ala Asn Cys Ile
 195 200 205
 Asn Leu Pro Gly Trp Tyr His Cys Glu Cys Arg Asp Gly Tyr His Asp
 210 215 220
 Asn Gly Met Phe Ser Pro Ser Gly Glu Ser Cys Glu Asp Ile Asp Glu
 225 230 235 240

<210> 157

<211> 342

<212> DNA

<213> Homo sapiens

<400> 157

gcagaaaatt ttccctctaga tcagaatctt caagaatcag ttaggttcct cactgcaaga 60
 aataaaatgt caggcagtga atgaattata ttttaagaag taaagcaaag aagctataac 120
 atgttatgta cagtacactc tgaaaagaaa tctgaaacaa gttattgtaa tgataaaaat 180
 aatgcacagg catggttact taatattttc taacaggaaa agtcacccct atttccttgt 240
 ttactgcac ttaatattat ttggttgaat ttgttcagta taagttcggt ccttggtgcaa 300
 aattaaataa atatttttct taccttaaaa aaaaaaaaaa aa 342

<210> 158

<211> 1445

<212> DNA

<213> Homo sapiens

<400> 158

```

gtgcgcatgg ggacgctata gcaattcggt tgcgtgcctt cctctccttc gaagatgaca 60
aggcctacca tcgtttcttc ctgccttttg gccgtcaggc agttgggttg gaccgcctcc 120
aaccctcggg tcttcctgca atacagtggg tacaatttgt catggctact ctgagataag 180
accacttttt tatctgagct tctgtgacct gctcctggga ctttgctggc tcacggagac 240
actttcttat ggagcttcag tagcaaataa ggacatcatc tgctataacc tacaagcagt 300
tggacagata ttctacattt cctcatttct ctacaccgtc aattacatct ggtatttgta 360
cacagagctg aggatgaaac acacccaaag tggacagagc acatctccac tggatgata 420
ttatacttgt cgattttgtc aaatggcctt tgttttctca agcctgatac ctctgctatt 480
gatgacacct gtattctgtc tgggaaatac tagtgaatgt ttccaaaact tcagtcagag 540
ccacaattgt atcttgatgc actcaccacc atcagccatg gctgaacttc caccttctgc 600
caacacatct gtctgtagca cactttattt ttatgggtatc gccattttcc tgggcagctt 660
tgtactcagc ctcttaccac ttatgggtctt acttatccga gccagacat tgtataagaa 720
gttttgtaag tcaactggct ttctggggag tgaacagtgg gcagtgatc acattgtgga 780
ccaacgggtg cgcttctacc cagtggcctt cttttgctgc tggggcccag ctgtcattct 840
aatgatcata aagctgacta agccacagga caccaagctt cacatggccc tttatgttct 900
ccaggctcta acggcaacat ctccagggtct actcaactgt ggagtatatg gctggacgca 960
gcacaaattc caccaactaa agcaggaggc tcggcggtgat gcagataccc agacaccatt 1020
attatgctca cagaagagat tctatagcag gggcttaaact tcaactggaat ccacctgac 1080
ttttctgccc agtacttcta ccattttttg aaactacaat actggaacat ccagggaactg 1140
gagttattct acgctaattg attggaaaga atgttgggaa aggacatctt aaatcttttc 1200
taactatgcc ctaaactgca gaactcaaa gaaatatagt gccattgtta gtagtcattc 1260
tagatgaatt gggagtatct ctccagttat tcccagattc actagtgatc cttaaagtct 1320
ctattcaggg agaggaagac actttccatc tcagagatag actcgtgtta ccttgatgga 1380
tattggattt gtctaagtct cttctagaaa aaataaattc tagattatta aaaaaaaaaa 1440
aaaaa 1445

```

<210> 159

<211> 245

<212> PRT

<213> Homo sapiens

<400> 159

```

Met Lys His Thr Gln Ser Gly Gln Ser Thr Ser Pro Leu Val Ile Asp
 1             5             10            15

Tyr Thr Cys Arg Phe Cys Gln Met Ala Phe Val Phe Ser Ser Leu Ile
      20             25             30

Pro Leu Leu Leu Met Thr Pro Val Phe Cys Leu Gly Asn Thr Ser Glu
      35             40             45

Cys Phe Gln Asn Phe Ser Gln Ser His Asn Cys Ile Leu Met His Ser
      50             55             60

Pro Pro Ser Ala Met Ala Glu Leu Pro Pro Ser Ala Asn Thr Ser Val
      65             70             75             80

Cys Ser Thr Leu Tyr Phe Tyr Gly Ile Ala Ile Phe Leu Gly Ser Phe
      85             90             95

Val Leu Ser Leu Leu Thr Ile Met Val Leu Leu Ile Arg Ala Gln Thr
      100            105            110

Leu Tyr Lys Lys Phe Val Lys Ser Thr Gly Phe Leu Gly Ser Glu Gln
      115            120            125

Trp Ala Val Ile His Ile Val Asp Gln Arg Val Arg Phe Tyr Pro Val

```

130 135 140
 Ala Phe Phe Cys Cys Trp Gly Pro Ala Val Ile Leu Met Ile Ile Lys
 145 150 155 160
 Leu Thr Lys Pro Gln Asp Thr Lys Leu His Met Ala Leu Tyr Val Leu
 165 170 175
 Gln Ala Leu Thr Ala Thr Ser Gln Gly Leu Leu Asn Cys Gly Val Tyr
 180 185 190
 Gly Trp Thr Gln His Lys Phe His Gln Leu Lys Gln Glu Ala Arg Arg
 195 200 205
 Asp Ala Asp Thr Gln Thr Pro Leu Leu Cys Ser Gln Lys Arg Phe Tyr
 210 215 220
 Ser Arg Gly Leu Asn Ser Leu Glu Ser Thr Leu Thr Phe Pro Ala Ser
 225 230 235 240
 Thr Ser Thr Ile Phe
 245

<210> 160

<211> 3550

<212> DNA

<213> Homo sapiens

<400> 160

ccctcgata atggattact aaatgggata cacgctgtac cagttcgctc cgagccccgg 60
 ccgctgtgcc gtcgatgcac cgaaaagggt gaagtagaga aataaagtct ccccgctgaa 120
 ctactatgag gtcagaagcc ttgctgctat atttcacact gctacacttt gctggggctg 180
 gtttcccaga agattctgag ccaatcagta ttctgcatgg caactataca aaacagtatc 240
 cgggtgttgt gggccacaag ccaggacgga acaccacaca gaggcacagg ctggacatcc 300
 agatgattat gatcatgaac ggaaccctct acattgctgc tagggaccat atttatactg 360
 ttgatataga cacatcacac acsgaagaaa tttattgtag caaaaaactg acatggaaat 420
 ctagacaggc cgatgtagac acatgcagaa tgaagggaaa acataaggat gagtgccaca 480
 actttattaa agttcttcta aagaaaaacg atgatgcatt gtttgtctgt ggaactaatg 540
 ccttcaaccc ttctgcaga aactataaga tggatacatt ggaaccattc ggggatgaat 600
 tcagcggaaat ggccagatgc ccatatgatg ccaaacatgc caacgttgca ctgtttgcag 660
 atggaaaact atactcagtc acagtgcact acttccttgc cattgacgca gtcatttacc 720
 ggagtcttgg agaaagccct accctgcgga ccgtcaagca cgattcaaaa tgggtgaaag 780
 aaccatactt tgttcaagcc gtggattacg gagattatat ctacttcttc ttcagggaaa 840
 tagcagtgga gtataacacc atgggaaagg tagttttccc aagagtggtc caggtttgta 900
 agaatgatat gggaggatct caaagagtcc tggagaaaca gtggacgtcg ttctgaagg 960
 cgcgcttgaa ctgctcagtt cctggagact ctcatTTTTA tttcaacatt ctccaggcag 1020
 ttacagatgt gattcgtatc aacgggctgt atgttgcctt ggcaacgttt tctacacctt 1080
 ataacagcat ccctgggtct gcagtctgtg cctatgacat gcttgacatt gccagtgttt 1140
 ttactgggag attcaaggaa cagaagtctc ctgattccac ctggacacca gttcctgatg 1200
 aacgagttcc taagcccagg ccagggtgct gtgctggctc atcctcctta gaaagatatg 1260
 caacctcaa tgagtccctt gatgataccc tgaacttcat caagacgcac ccgctcatgg 1320
 atgaggcagt gccctccatc ttcaacaggc catggttccct gagaacaatg gtcagatacc 1380
 gccttaccaa aattgcagtg gacacagctg ctgggccata tcagaatcac actgtggttt 1440
 ttctgggatac agagaagga atcatcttga agtttttggc cagaatagga aatagtgttt 1500
 ttctaaatga cagccttttc ctggaggaga tgagtgttta caactctgaa aatgcagct 1560
 atgatggagt cgaagacaaa aggatcatgg gcatgcagct ggacagagca agcagctctc 1620
 tgtatgttgc gttctctacc tgtgtgataa aggttccctt tggccggtgt gaacgacatg 1680
 ggaagtgtaa aaaaacctgt attgcctcca gagaccata ttgtggatgg ataaaggaag 1740
 gtgggtcctg cagccattta tcaccaaca gcagactgac ttttgagcag gacatagagc 1800

```

gtggcaatac agatgggtctg ggggactgtc acaattcctt tgtggcactg aatggagtga 1860
ttcgggaaag ttacctcaaa ggccacgacc agctgggtcc cgtcacccctc ttggccattg 1920
cagtcatacct ggcttttcgtc atgggggccc tcttctcggg catcacccgtc tactgcgtct 1980
gtgatcatcg gcgcaaagac gtggctgtgg tgcagcgcaa ggagaaggag ctcacccact 2040
cgcgccgggg ctccatgagc agcgtcacca agctcagcgg cctcttttggg gacactcaat 2100
ccaaagaccc aaagccggag gccatcctca cgccactcat gcacaacggc aagctcgcca 2160
ctccccggcaa cacggccaag atgctcatta aagcagacca gcaccacctg gacctgacgg 2220
ccctccccac ccagagtgca accccaacgc tgcagcagaa gcggaagccc agccgcgga 2280
gccgcgagtg ggagaggaac cagaacctca tcaatgcctg cacaaaggac atgcccccca 2340
tgggctcccc tgtgattccc acggacctgc ccctgcgggc ctccccagc cacatcccc 2400
gcgtggtggt cctgccccac acgcagcagg gctaccagca tgagtacgtg gaccagccca 2460
aaatgagcga ggtggcccag atggcgctgg aggaccaggc cgccacactg gagtataaga 2520
ccatcaagga acatttcagc agcaagagtc ccaaccatgg ggtgaacctt gtggagaacc 2580
tggacagcct gcccccaaaa gtccacagc gggaggcctc cctgggtccc ccgggagcct 2640
ccctgtttca gaccggttta agcaagcggc tggaaatgca ccactccttt tctacgggg 2700
ttgactataa gaggagctac ccacgaact cgtcacgag aagccaccag gccaccactc 2760
tcaaaagaaa caacactaac tctccaatt cctctcacct ctccagaaac cagagctttg 2820
gcaggggaga caaccgcgg ccgcggcggc agagggtgga ctccatccag gtgcacagct 2880
cccagccatc tggccaggcc gtgactgtct cgaggcagcc cagcctcaac gcctacaact 2940
cactgacaag gtcggggctg aagcgtacgc cctcgctaaa gccggacgta cccccaaa 3000
catcctttgc tcccccttcc acatccatga agcccaatga tgcgtgtaca taatcccagg 3060
gggagggggg caggtgtcga accagcaggc aaggcgaggt gcccgctcag ctcagcaagg 3120
ttctcaactg cctcgagtac ccaccagacc aagaaggcct gcggcagagc cgaggacgct 3180
gggtcctcct ctctgggaca caggggtact cagaaaaact gggccgctg gtttgggtgaa 3240
ggtttgcaac ggcggggact cacccttcatt ctcttccttc actttcccc acaccctaca 3300
acaggtcgga cccacaaaag acttcagtta tcatcacaaa catgagccaa aagcacatac 3360
ataccccatc cccaccccc acacacacac acacatgcac acaacacata cacacacagc 3420
cacagaggtg aacagaaact gaaacatttt gtccacaact tcacgggacg tggccagact 3480
gggtttgcgt tccaacctgc aaaacacaaa tacatttttt aaaatcaaga aaatttataa 3540
aaaaaaaaa 3550

```

<210> 161

<211> 975

<212> PRT

<213> Homo sapiens

<400> 161

```

Met Arg Ser Glu Ala Leu Leu Leu Tyr Phe Thr Leu Leu His Phe Ala
  1             5             10             15

```

```

Gly Ala Gly Phe Pro Glu Asp Ser Glu Pro Ile Ser Ile Ser His Gly
      20             25             30

```

```

Asn Tyr Thr Lys Gln Tyr Pro Val Phe Val Gly His Lys Pro Gly Arg
  35             40             45

```

```

Asn Thr Thr Gln Arg His Arg Leu Asp Ile Gln Met Ile Met Ile Met
  50             55             60

```

```

Asn Gly Thr Leu Tyr Ile Ala Ala Arg Asp His Ile Tyr Thr Val Asp
  65             70             75             80

```

```

Ile Asp Thr Ser His Thr Glu Glu Ile Tyr Cys Ser Lys Lys Leu Thr
      85             90             95

```

```

Trp Lys Ser Arg Gln Ala Asp Val Asp Thr Cys Arg Met Lys Gly Lys
  100             105             110

```

```

His Lys Asp Glu Cys His Asn Phe Ile Lys Val Leu Leu Lys Lys Asn
  115             120             125

```

Asp Asp Ala Leu Phe Val Cys Gly Thr Asn Ala Phe Asn Pro Ser Cys
 130 135 140
 Arg Asn Tyr Lys Met Asp Thr Leu Glu Pro Phe Gly Asp Glu Phe Ser
 145 150 155 160
 Gly Met Ala Arg Cys Pro Tyr Asp Ala Lys His Ala Asn Val Ala Leu
 165 170 175
 Phe Ala Asp Gly Lys Leu Tyr Ser Ala Thr Val Thr Asp Phe Leu Ala
 180 185 190
 Ile Asp Ala Val Ile Tyr Arg Ser Leu Gly Glu Ser Pro Thr Leu Arg
 195 200 205
 Thr Val Lys His Asp Ser Lys Trp Leu Lys Glu Pro Tyr Phe Val Gln
 210 215 220
 Ala Val Asp Tyr Gly Asp Tyr Ile Tyr Phe Phe Phe Arg Glu Ile Ala
 225 230 235 240
 Val Glu Tyr Asn Thr Met Gly Lys Val Val Phe Pro Arg Val Ala Gln
 245 250 255
 Val Cys Lys Asn Asp Met Gly Gly Ser Gln Arg Val Leu Glu Lys Gln
 260 265 270
 Trp Thr Ser Phe Leu Lys Ala Arg Leu Asn Cys Ser Val Pro Gly Asp
 275 280 285
 Ser His Phe Tyr Phe Asn Ile Leu Gln Ala Val Thr Asp Val Ile Arg
 290 295 300
 Ile Asn Gly Arg Asp Val Val Leu Ala Thr Phe Ser Thr Pro Tyr Asn
 305 310 315 320
 Ser Ile Pro Gly Ser Ala Val Cys Ala Tyr Asp Met Leu Asp Ile Ala
 325 330 335
 Ser Val Phe Thr Gly Arg Phe Lys Glu Gln Lys Ser Pro Asp Ser Thr
 340 345 350
 Trp Thr Pro Val Pro Asp Glu Arg Val Pro Lys Pro Arg Pro Gly Cys
 355 360 365
 Cys Ala Gly Ser Ser Ser Leu Glu Arg Tyr Ala Thr Ser Asn Glu Phe
 370 375 380
 Pro Asp Asp Thr Leu Asn Phe Ile Lys Thr His Pro Leu Met Asp Glu
 385 390 395 400
 Ala Val Pro Ser Ile Phe Asn Arg Pro Trp Phe Leu Arg Thr Met Val
 405 410 415
 Arg Tyr Arg Leu Thr Lys Ile Ala Val Asp Thr Ala Ala Gly Pro Tyr
 420 425 430
 Gln Asn His Thr Val Val Phe Leu Gly Ser Glu Lys Gly Ile Ile Leu
 435 440 445

Lys Phe Leu Ala Arg Ile Gly Asn Ser Gly Phe Leu Asn Asp Ser Leu
 450 455 460
 Phe Leu Glu Glu Met Ser Val Tyr Asn Ser Glu Lys Cys Ser Tyr Asp
 465 470 475 480
 Gly Val Glu Asp Lys Arg Ile Met Gly Met Gln Leu Asp Arg Ala Ser
 485 490 495
 Ser Ser Leu Tyr Val Ala Phe Ser Thr Cys Val Ile Lys Val Pro Leu
 500 505 510
 Gly Arg Cys Glu Arg His Gly Lys Cys Lys Lys Thr Cys Ile Ala Ser
 515 520 525
 Arg Asp Pro Tyr Cys Gly Trp Ile Lys Glu Gly Gly Ala Cys Ser His
 530 535 540
 Leu Ser Pro Asn Ser Arg Leu Thr Phe Glu Gln Asp Ile Glu Arg Gly
 545 550 555 560
 Asn Thr Asp Gly Leu Gly Asp Cys His Asn Ser Phe Val Ala Leu Asn
 565 570 575
 Gly Val Ile Arg Glu Ser Tyr Leu Lys Gly His Asp Gln Leu Val Pro
 580 585 590
 Val Thr Leu Leu Ala Ile Ala Val Ile Leu Ala Phe Val Met Gly Ala
 595 600 605
 Val Phe Ser Gly Ile Thr Val Tyr Cys Val Cys Asp His Arg Arg Lys
 610 615 620
 Asp Val Ala Val Val Gln Arg Lys Glu Lys Glu Leu Thr His Ser Arg
 625 630 635 640
 Arg Gly Ser Met Ser Ser Val Thr Lys Leu Ser Gly Leu Phe Gly Asp
 645 650 655
 Thr Gln Ser Lys Asp Pro Lys Pro Glu Ala Ile Leu Thr Pro Leu Met
 660 665 670
 His Asn Gly Lys Leu Ala Thr Pro Gly Asn Thr Ala Lys Met Leu Ile
 675 680 685
 Lys Ala Asp Gln His His Leu Asp Leu Thr Ala Leu Pro Thr Pro Glu
 690 695 700
 Ser Thr Pro Thr Leu Gln Gln Lys Arg Lys Pro Ser Arg Gly Ser Arg
 705 710 715 720
 Glu Trp Glu Arg Asn Gln Asn Leu Ile Asn Ala Cys Thr Lys Asp Met
 725 730 735
 Pro Pro Met Gly Ser Pro Val Ile Pro Thr Asp Leu Pro Leu Arg Ala
 740 745 750
 Ser Pro Ser His Ile Pro Ser Val Val Val Leu Pro Ile Thr Gln Gln
 755 760 765

Gly Tyr Gln His Glu Tyr Val Asp Gln Pro Lys Met Ser Glu Val Ala
 770 775 780
 Gln Met Ala Leu Glu Asp Gln Ala Ala Thr Leu Glu Tyr Lys Thr Ile
 785 790 795 800
 Lys Glu His Phe Ser Ser Lys Ser Pro Asn His Gly Val Asn Leu Val
 805 810 815
 Glu Asn Leu Asp Ser Leu Pro Pro Lys Val Pro Gln Arg Glu Ala Ser
 820 825 830
 Leu Gly Pro Pro Gly Ala Ser Leu Phe Gln Thr Gly Leu Ser Lys Arg
 835 840 845
 Leu Glu Met His His Ser Phe Ser Tyr Gly Val Asp Tyr Lys Arg Ser
 850 855 860
 Tyr Pro Thr Asn Ser Leu Thr Arg Ser His Gln Ala Thr Thr Leu Lys
 865 870 875 880
 Arg Asn Asn Thr Asn Ser Ser Asn Ser Ser His Leu Ser Arg Asn Gln
 885 890 895
 Ser Phe Gly Arg Gly Asp Asn Pro Pro Pro Ala Pro Gln Arg Val Asp
 900 905 910
 Ser Ile Gln Val His Ser Ser Gln Pro Ser Gly Gln Ala Val Thr Val
 915 920 925
 Ser Arg Gln Pro Ser Leu Asn Ala Tyr Asn Ser Leu Thr Arg Ser Gly
 930 935 940
 Leu Lys Arg Thr Pro Ser Leu Lys Pro Asp Val Pro Pro Lys Pro Ser
 945 950 955 960
 Phe Ala Pro Leu Ser Thr Ser Met Lys Pro Asn Asp Ala Cys Thr
 965 970 975

<210> 162

<211> 1723

<212> DNA

<213> Homo sapiens

<400> 162

ctgcagactt tggggtcacc ggccagccac acaggcaccg ttttcagatg tccacttctc 60
 attgggtaca tcaatctttt aacttttgggg gtcacagttt tagccacctt tcgggggggtg 120
 actggagcag taggaggtgt ggggtcattt tatgaatata ataaaatgga gctgactatg 180
 gacrrrgact wagtgtgggg gagaggggac gatacagggt gtgtgtcttg gagtgccttg 240
 gggacaggga ccccccggtg gtccctatggc aggatgagaa rggagggact tggctcccc 300
 agagccccgt ggaagctact gttctctcca gtgtctcgag cgtagccaaa ataaggttg 360
 gaggtctccg gcctgtctgc tgtggtctga gctggctgca agcccagggt ggggagcgag 420
 tctgggaaga ttggctttga ctctctgttg ccagaggaga tgccatccca gcacggcccc 480
 cactgtagtc caggctcgtg gtggcagcgg gggcaagggg aggggcaagg ctgccccccac 540
 cccacgcacc aagtcacgcc aagtctcagc aggtaaaagc acgtgagcct agggcgagcg 600
 gagggagtc tgggtggcccc gcaggtcagg agggaaagca gggctcagag ggcacgtgg 660
 cccaggggca gggctctacc tgggggtcag gacaccttg gtcttgatga ttgattgatt 720
 gatagaatgg agctgggtct gagcctccca ggcttgagct cctgggagtt cttgtgcggt 780


```

gagctgggca gctcctgggt aggtccgggc accaagcagg ccctgatgtg gacagagtcc 840
catcagaggg agctgatgaa gaatgggtccc tgtaagtaag tcactagggt caacaactgc 900
ctggccgagc actcagcccg tggagctcag gccaacacca gagccccggt tttagggggc 960
aggagagcag gtgaccaatt atttggggag tcttgggtag aatttccgcc acacattctc 1020
cccagggctg caggggtctt ccgaggcagg gcggtggagc aggattcagg atgtggtggg 1080
aatagagtga ggggcagtgg gtgggcagac ctgggcgtca gaggtcctga tgggaaagga 1140
ggcaggggct acccagagag gggggctcgt gtggcacagc ccccaccgac tccgccgtcc 1200
ccctcccctg tgagcccccg gggtgtaca tactctactc catccccttg tccatccctg 1260
agaccacccc cgcgcgccct gcgtcgactt agcaaccacc tcataggccc acccaccctc 1320
ggatccgagc caaccatccc acatcacaaa ctttgggttg ggggacttta cgttcgttta 1380
atttctcatt ttgtacggag aaatattctt ttcaaaagcg tcttttgact gaagtaactt 1440
tcctggtgct gttgttaact cgttcctttt ttttaatttat tccccaccc caggcagccc 1500
tcctggttcc tactcaccct cccccctcc cccaccctcc gtcccatctg aaccatttgt 1560
ttcttttctt tccgtcagat ttggaaaaa ttctctctc ctccccgccc cctccacacc 1620
atcctcccsa atttaaatat agtcactgct acaagtaaca gatgcactgt gaagattcca 1680
gtattaataa aggtgtactg taattaacaa aaaaaaaaaa aaa 1723

```

<210> 163

<211> 101

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (49)

<220>

<221> UNSURE

<222> (51)

<220>

<221> UNSURE

<222> (81)

<400> 163

```

Val Phe Arg Cys Pro Leu Leu Ile Gly Tyr Ile Asn Leu Leu Thr Leu
  1           5           10           15

```

```

Gly Val Thr Val Leu Ala Thr Phe Arg Gly Val Thr Gly Ala Val Gly
          20           25           30

```

```

Gly Val Gly Ser Phe Tyr Glu Tyr Asn Lys Met Glu Leu Thr Met Asp
  35           40           45

```

```

Xaa Asp Xaa Val Trp Gly Arg Gly Asp Asp Thr Gly Cys Val Ser Gly
  50           55           60

```

```

Ser Ala Trp Gly Thr Gly Thr Pro Arg Trp Ser Tyr Gly Arg Met Arg
  65           70           75           80

```

```

Xaa Glu Gly Leu Gly Ser Pro Arg Ala Arg Trp Lys Leu Leu Phe Ser
          85           90           95

```

```

Pro Val Ser Arg Ala
          100

```

<210> 164

<211> 469

<212> DNA

<213> Homo sapiens

<400> 164

```
gcaacatata agccggccat attagagaga tggaaataaa gcttccttaa tgttgatat 60
gtctttgaag tacatccgtg catttttttt tagcatccaa ccattcctcc cttgtagttc 120
tcgccccctc aaatcacccct ctcccgtagc ccacccgact aacatctcag tctctgaaaa 180
tgcacagaga tgcttggtta cctcgccctg ccttcagcct cacggggctc agtctctttt 240
tctctttggt gccaccagga cggagcatgg aggtcacagt acctgccacc ctcaacgtcc 300
tcaatggctc tgacgcccgc ctgcccgtga ccttcaactc ctgctacaca gtgaaccaca 360
aacagttctc cctgaactgg acttaccagg agtgcaacaa ctgctctgag gagatgttcc 420
tccagttccg catgaagatc attaacctga agctggagcg gtttcaaga 469
```

<210> 165

<211> 96

<212> PRT

<213> Homo sapiens

<400> 165

```
Met His Arg Asp Ala Trp Leu Pro Arg Pro Ala Phe Ser Leu Thr Gly
  1              5              10              15

Leu Ser Leu Phe Phe Ser Leu Val Pro Pro Gly Arg Ser Met Glu Val
      20              25              30

Thr Val Pro Ala Thr Leu Asn Val Leu Asn Gly Ser Asp Ala Arg Leu
      35              40              45

Pro Cys Thr Phe Asn Ser Cys Tyr Thr Val Asn His Lys Gln Phe Ser
      50              55              60

Leu Asn Trp Thr Tyr Gln Glu Cys Asn Asn Cys Ser Glu Glu Met Phe
      65              70              75              80

Leu Gln Phe Arg Met Lys Ile Ile Asn Leu Lys Leu Glu Arg Phe Gln
      85              90              95
```

<210> 166

<211> 454

<212> DNA

<213> Homo sapiens

<400> 166

```
tggcttttgg ctacagagag ggaagggaaa gcctgaggcc ggcataaggg gaggccttgg 60
aacctgagct gccaatgcca gccctgtccc atctgcgccc acgatactcg ctctctctcc 120
aacaactcct ttggtgggga caaaagtgc aattgtaggc caggcacagt ggctcacgcc 180
tgtaatccca gcactttggg aggccaaggc ggggtgatta cctccatctg tttagtagaa 240
atggggcaaaa ccccatTTTT actaaaaata caagaattag ctgggcgtgg tggcgtgtgc 300
ctgtaatccc agctatttgg gaggctgagg caggagaatc gcttgagccc gggaagcaga 360
ggttgacagt aactgagata gtgatagtgc cactgcaatt cagcctgggt gacatagaga 420
gactccatct caaaaaaaaa aaaaaaaaaa aaaa 454
```

<210> 167

<211> 736

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (680)

<220>
 <221> unsure
 <222> (704)

<400> 167
 gttttttaa ac attatgttct acatgataaa tacatataat agtatgtcta tttaaataat 60
 taaatttgaa aaaaactaat caaatattat cataagtaat gataaaaacc acaatttctt 120
 ttgcagcaaa ctaataacac ctggatttct caatttatta agttgtactt acctgatgct 180
 gatgatgatt actgtattta cacattgtct cagagctcac tcttgccggag gttgtggcct 240
 cgaaaatgcc ttgttgcccc tctggaatct gtcttttcag cttcatctcc tcctcctcac 300
 ctccctgctgt ggtgcacaga tacctatagg caggctccat ctccctcctcc ccagctcctc 360
 ccctagtcca cagataccta taggcaggct tcatctcctc ctccccagct tctcccctag 420
 tgcacagata cctataggca ggctccatct cctcctcccc agctcctccc ctartgcaca 480
 gacacctata ggcaagctcc atctcctcct ctttagctag cctccccatc tcatcacaac 540
 gcatgtctgt gaccttttgt aatcatttac agtgccacac ggaaccctgt attttgcaca 600
 cagcaaaaca aacaatgttt agctttatct atggtatttg atgactgtaa atggaaataa 660
 atattgttct ttattttttt aaaaaaaaaa aaaaaaaaaa aaanaaaaaa aaaaaaaaaa 720
 aaaaaaaaaa aaaaaa 736

<210> 168
 <211> 114
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (100)

<400> 168
 Met Leu Met Met Ile Thr Val Phe Thr His Cys Leu Arg Ala His Ser
 1 5 10 15
 Cys Gly Gly Cys Gly Leu Glu Asn Ala Leu Leu Ser Leu Trp Asn Leu
 20 25 30
 Ser Phe Gln Leu His Leu Leu Leu Thr Ser Cys Cys Gly Ala Gln
 35 40 45
 Ile Pro Ile Gly Arg Leu His Leu Leu Leu Pro Ser Ser Ser Pro Ser
 50 55 60
 Ala Gln Ile Pro Ile Gly Arg Leu His Leu Leu Leu Pro Ser Phe Ser
 65 70 75 80
 Pro Ser Ala Gln Ile Pro Ile Gly Arg Leu His Leu Leu Leu Pro Ser
 85 90 95
 Ser Ser Pro Xaa Ala Gln Thr Pro Ile Gly Lys Leu His Leu Leu Leu
 100 105 110
 Phe Ser

<210> 169
 <211> 1427
 <212> DNA
 <213> Homo sapiens

<400> 169

```

gtagttacta actccaacac ctaatagcat tggtagaaag cttataaatg cagttattta 60
gcctcgacta agatttttct gatacctagt ttcacttttt aatgccctct gaaagttttt 120
tgatcagttg tttaatggga gatctgaaat gttaaactca gaccagaaag aagagaacct 180
gttttctaga aattagggtt ttaatccaag taagatgcaa gcttttgctt ttttaataac 240
ttgtatagct aaaaacttga cggtgaaaag ctctcagatc aaagctgac cttctgtcag 300
taatgattct aaaaataagc aagattttta tggggaatat attttatttc attcttatct 360
caaacctagg tactgtggtc gttttgagtt catttcgagg cattttcaat gtgcctcagg 420
ccacatccaa cctctycca gggccagatt taatgttcag cctcataaag gttatcatag 480
ttttaacatt taagtactat tttgcagtg gtatatacca aaatttgcta atagtaagat 540
aaccttagtt atatatcatt cacgttagtt ctatcttgga ggcaataaac atttcttggt 600
caagaaattc atgttctatc ttggaggcaa taaacaaaca ttttttgctt aaaattaggg 660
ctaccctatt gtccttatgt cttttctctga tctgtggtca aacatttttc ttagtcattt 720
agaaattttc tatgttggtt taaattttct ttaaacttag aatggagtat gtgaccaata 780
ctttcctttg gaatggtatg gacatttgaa atagagccca ttctttataa agtataaaat 840
atgtttaatg ctagtatttt taactaaact tttgagaaac tagattcaca tgctgttgta 900
agaaataata cagagacctc tttcgtgtac ctttcacttt gtttccaca cagtgaacat 960
ctttcaaaac tgcatataca tatcataccc aggatactga cactggtata gctaagatag 1020
agaacgtttc cacacagaac tttttctagc acagggatcc ctcatcttgc ttttgatgac 1080
cataccact tcaactccat cctactccc ttcttaacct ttggcaacca taatctgttc 1140
tccattttta tagttttttt tttttcattt caataaagct gtataactgg aatcataata 1200
atatgtaacc ttttgggatt ggcttttttt catttagcat gattttcttg aggttaatcc 1260
agcttattat gtgtatcaag tctattgaca ggtacttttt agtgtgaata gaatccata 1320
gtatagatgt accacagttt gtttaactgt tcacctgctg agagacattg ggccagtttt 1380
tggctactat aaataaagtt gctataaaca aaaaaaaaa aaaaaaa 1427

```

<210> 170

<211> 79

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (45)

<400> 170

```

Met Ile Leu Lys Ile Ser Lys Ile Leu Met Gly Asn Ile Phe Tyr Phe
 1             5             10             15
Ile Leu Ile Ser Asn Leu Gly Thr Val Val Val Leu Ser Ser Phe Arg
          20             25             30
Gly Ile Phe Asn Val Pro Gln Ala Thr Ser Asn Leu Xaa Pro Gly Pro
          35             40             45
Asp Leu Met Phe Ser Leu Ile Lys Val Ile Ile Val Leu Thr Phe Lys
          50             55             60
Tyr Tyr Phe Ala Val Gly Ile Tyr Gln Asn Leu Leu Ile Val Arg
          65             70             75

```

<210> 171

<211> 572

<212> DNA

<213> Homo sapiens

<400> 171

```

tgcagattct gtggttatac tcactctca tcccaaagaa tgaaatttac cactctctc 60
ttcttggcag ctgtagcagg ggccttggtc tatgctgaag atgcctctc tgactcgacg 120

```

```

ggtgctgata ctgcccagga agctgggacc tetaagccta atgaagagat ctcagggtcca 180
gcagaaccag cttcaccccc agagacaacc acaacagccc aggagacttc ggcggcagca 240
gttcagggga cagccaaggt cacctcaagc aggcaggaac taaaccccct gaaatccata 300
gtggagaaaa gtatcttact aacagaacaa gcccttgcaa aagcaggaaa aggaatgcac 360
ggaggcgtgc caggtggaaa acaattcatc gaaaatggaa gtgaatttgc acaaaaatta 420
ctgaagaaat tcagtctatt aaaacccatgg gcatgagaag ctgaaaagaa tgggatcatt 480
ggacttaaaag ccttaaatac ccttgtagcc cagagytatt aaaacgaaag catccaaaaa 540
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aa 572

```

<210> 172

<211> 138

<212> PRT

<213> Homo sapiens

<400> 172

```

Met Lys Phe Thr Thr Leu Leu Phe Leu Ala Ala Val Ala Gly Ala Leu
  1             5             10            15

Val Tyr Ala Glu Asp Ala Ser Ser Asp Ser Thr Gly Ala Asp Pro Ala
             20             25            30

Gln Glu Ala Gly Thr Ser Lys Pro Asn Glu Glu Ile Ser Gly Pro Ala
  35             40            45

Glu Pro Ala Ser Pro Pro Glu Thr Thr Thr Thr Ala Gln Glu Thr Ser
  50             55            60

Ala Ala Ala Val Gln Gly Thr Ala Lys Val Thr Ser Ser Arg Gln Glu
  65             70            75            80

Leu Asn Pro Leu Lys Ser Ile Val Glu Lys Ser Ile Leu Leu Thr Glu
             85             90            95

Gln Ala Leu Ala Lys Ala Gly Lys Gly Met His Gly Gly Val Pro Gly
 100             105            110

Gly Lys Gln Phe Ile Glu Asn Gly Ser Glu Phe Ala Gln Lys Leu Leu
 115             120            125

Lys Lys Phe Ser Leu Leu Lys Pro Trp Ala
 130             135

```

<210> 173

<211> 1223

<212> DNA

<213> Homo sapiens

<400> 173

```

ccttgttcca cgtagctggc aaggtcttca ttcacttgcc actgctagtc ttccaaccct 60
ctcggacttt ctttgctctt gtcttgtttt ggggtgactg gatcatgaca cttctttttc 120
ttggcactac cggcagtcct gttcagaatg agcaaggctt tgtggagtgc aaaattttctg 180
ggcctctgca gtacatgtgg tggtagcatg tggtagggcct gatttggatc agtgaattta 240
ttctagcatg tcagcagatg acagtggcag gagctgtggt aacatactat ttactagggt 300
ataaaaggaa tttgccattt acacctattt tggcatcagt aaatcgccct atycgttacc 360
acctaggtag ggtggcaaaa ggtcttttca ttatcacatt agtcaaaatt ccgcgaatga 420
tccttatgta tattcacagt cagctcaaag gaaaggaaaa tgcttgtgca cgatgtgtgc 480
tgaatcttgc catttggtgc ctttggtgtc ttgaaaagtg cctaaattat ttaaatcaga 540
atgcatacac agccacagct atcaacagca ccaacttctg cacctcagca aaggatgcct 600
ttgtcattct ggtggagaat gcttgcgag tggctaccat caacacagta ggagatttta 660

```

```

tgttattcct tggcaagggt ctgatagtct gcagcacagg tttagctggg attatgctgc 720
tcaactacca gcaggactac acagtatggg tgctgcctct gatcatcgtc tgcctctttg 780
ctttcctagt cgctcattgc ttcctgtcta tttatgaaat ggtagtggat gtattattct 840
kgkgtttttc cattgawaca aaatacaatg atgggmgcc tggcagagaa ttctatatgg 900
ataaagtgtc gatggagttt gtggaaaaca gtaggaaagc aatgaaagaa gctggttaagg 960
gaggcgctgc tgattccaga gagctaaagc cgatgctgaa gaaaagggtga ctggtctcat 1020
gagccctgaa gaatgaactc agaggagggt gtttacctga ggttctccca ctcaccagct 1080
gttgagagtc tgcgattatg aagagcagga tcttattact tcaatgaaag catgtaacaa 1140
gtttctcaaa ccaccaacag ccaagtggat ttggtacagt gcggctgtct aataaataat 1200
caaaagcaaa aaaaaaaaaa aaa                                     1223

```

```

<210> 174
<211> 301
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> UNSURE
<222> (246)..(247)

```

```

<220>
<221> UNSURE
<222> (251)

```

```

<220>
<221> UNSURE
<222> (258)

```

```

<400> 174
Met Thr Leu Leu Phe Leu Gly Thr Thr Gly Ser Pro Val Gln Asn Glu
  1              5              10              15

Gln Gly Phe Val Glu Phe Lys Ile Ser Gly Pro Leu Gln Tyr Met Trp
      20              25              30

Trp Tyr His Val Val Gly Leu Ile Trp Ile Ser Glu Phe Ile Leu Ala
  35              40              45

Cys Gln Gln Met Thr Val Ala Gly Ala Val Val Thr Tyr Tyr Phe Thr
  50              55              60

Arg Asp Lys Arg Asn Leu Pro Phe Thr Pro Ile Leu Ala Ser Val Asn
  65              70              75              80

Arg Leu Ile Arg Tyr His Leu Gly Thr Val Ala Lys Gly Ser Phe Ile
      85              90              95

Ile Thr Leu Val Lys Ile Pro Arg Met Ile Leu Met Tyr Ile His Ser
  100             105             110

Gln Leu Lys Gly Lys Glu Asn Ala Cys Ala Arg Cys Val Leu Lys Ser
  115             120             125

Cys Ile Cys Cys Leu Trp Cys Leu Glu Lys Cys Leu Asn Tyr Leu Asn
  130             135             140

Gln Asn Ala Tyr Thr Ala Thr Ala Ile Asn Ser Thr Asn Phe Cys Thr
  145             150             155             160

Ser Ala Lys Asp Ala Phe Val Ile Leu Val Glu Asn Ala Leu Arg Val

```

165	170	175
Ala Thr Ile Asn Thr Val Gly Asp Phe Met Leu Phe Leu Gly Lys Val		
180	185	190
Leu Ile Val Cys Ser Thr Gly Leu Ala Gly Ile Met Leu Leu Asn Tyr		
195	200	205
Gln Gln Asp Tyr Thr Val Trp Val Leu Pro Leu Ile Ile Val Cys Leu		
210	215	220
Phe Ala Phe Leu Val Ala His Cys Phe Leu Ser Ile Tyr Glu Met Val		
225	230	235
Val Asp Val Leu Phe Xaa Xaa Phe Ala Ile Xaa Thr Lys Tyr Asn Asp		
245	250	255
Gly Xaa Pro Gly Arg Glu Phe Tyr Met Asp Lys Val Leu Met Glu Phe		
260	265	270
Val Glu Asn Ser Arg Lys Ala Met Lys Glu Ala Gly Lys Gly Gly Val		
275	280	285
Ala Asp Ser Arg Glu Leu Lys Pro Met Leu Lys Lys Arg		
290	295	300

<210> 175

<211> 2460

<212> DNA

<213> Homo sapiens

<400> 175

```

gagaagcgcg atgacggcta cgaggccgct gccagctcca aaactagctc aggcgatgcc 60
tcctcactca gcatcgaggg agactaacia actccgggca aagttggggc tgaaaccctt 120
ggagggttaat gccatcaaga aggaggcggg caccaaggag gagcccgtga cagctgatgt 180
catcaaccct atggccttgc gacagcgaga ggagctgcgg gagaagctgg cggctgcca 240
ggagaagcgc ctgctgaacc aaaagctggg gaagataaag accctaggag aggatgaccc 300
ctggctggac gacactgcag cctggatcga gaggagccgg cagctgcaga aggagaagga 360
cctggcagag aagagggcca agttactgga ggagatggac caaaagtgtg gtgtcagcac 420
tctggtggag gaggagtctg ggcagaggcg gcaggacctg tacagtgcgc gggacctgca 480
gggcctcact gtggagcatg ccattgattc cttccgagaa ggggagacaa tgattcttac 540
cctcaaggac aaaggcgtgc tgcaggagga ggaggacgtg ctggtgaacg tgaacctggt 600
ggataaggag cgggcagaga aaaatgtgga gctgcggaag aagaagcctg actacctgcc 660
ctatgccgag gacgagagcg tggacgacct ggcgagcaa aaacctcgct ctatcctgtc 720
caagtatgac gaaaagcttg aaggggagcg gccacattcc ttccgcttgg agcagggcgg 780
cacggctgat ggcctgcggg agcgggagct ggaggagatc cgggccaagc tgcggctgca 840
ggctcagtc ctgagcacag tggggccccg gctggcctcc gaatacctca cgcctgagga 900
gatggtgacc tttaaaaaga ccaagcggag ggtgaagaaa atccgcaaga aggagaagga 960
ggtagtagtg cgggcagatg acttgctgcc tctcggggac cagactcagg atggggactt 1020
tggttccaga ctgcggggac ggggtcgcgg ccgagtgtcc gaagtggagg aggagaagga 1080
gcctgtgcct cagccccctg cgtcgagaca caccagtg gagaacatgg acatcagtga 1140
tgaggaggaa ggtggagctc caccgcccgg gtccccgag gtgctggagg aggacgaggc 1200
ggagctggag ctgcagaagc agctggagaa gggacgcccg ctgcgacagt tacagcagct 1260
acagcagctg cgagacagtg gcgagaaggt ggtggagatt gtgaagaagc tggagtctcg 1320
ccagcggggc tgggaggagg atgaggatcc cgagcggaa gggggccatc tgttcaacgc 1380
cacgtccgag ttctgcgcga ccttggggga gatccccacc tacgggctgg ctggcaatcg 1440
cgaggagcag gaggagctca tggactttga acgggatgag gacgctcag ccaacgggtg 1500
ctccgaatct gacggggagg agaacatcgg ctggagcacg gtgaacctgg acgaggagaa 1560
gcacgacgag gatgtgaggg ccacgccgct ggggggtggg cgtttggggg tgctcaagct 1620

```

ggagatgagc accgggctcg gtgtccagag cctcagcctc ctcattccaga gtgggctctg 1680
 cagacctccc agggcgatct gaggagtaaa tgaggaaatt aaatgttgtg gagggctggt 1740
 gcctggcagg tggtagaccag tgggtggggc tgagaagagc cggtagggcc tgctaaccac 1800
 ccccgccacg tgtcccgtag ttctctgctt cctccaccac catcctggac gaggaaccga 1860
 tcgtgaatag ggggttgga gctgccctgc tcctgtgtca gaacaaaggg ctgctggaga 1920
 ccacagtgcga gaaggtggcc cgggtgaagg cccccaacaa gtcgctgccc tcagccgtgt 1980
 actgcatcga ggataagatg gccatcgatg acaagtacag ccggagggag gaataccgag 2040
 gcttcacaca ggacttcaag gagaaggacg gtacaaaccc gacgttaaga tcgaatacgt 2100
 ggatgagacg ggccggaaac tcacacccaa ggaggctttc cggcagctgt cgcaccgctt 2160
 ccattggcaag ggctcaggca agatgaagac agagcggcgg atgaagaagc tggacgagga 2220
 ggcgctcctg aagaagatga gctccagsga cagcccytg ggcaccgtgg cctgytcca 2280
 ggagaagcag aaggtcaga agaccccta catygtgttc agcggcagcg gcaagagcat 2340
 gaacgcgaac accatcacca agtgacagcg ccctcccgcc cggccctgc ctcaaccttc 2400
 atattaaata aagctccctc cttaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2460

<210> 176

<211> 563

<212> PRT

<213> Homo sapiens

<400> 176

Met	Thr	Ala	Thr	Arg	Pro	Leu	Pro	Ala	Pro	Lys	Leu	Ala	Gln	Ala	Met
1				5					10					15	
Pro	Pro	His	Ser	Ala	Ser	Arg	Glu	Thr	Asn	Lys	Leu	Arg	Ala	Lys	Leu
			20					25					30		
Gly	Leu	Lys	Pro	Leu	Glu	Val	Asn	Ala	Ile	Lys	Lys	Glu	Ala	Gly	Thr
	35					40						45			
Lys	Glu	Glu	Pro	Val	Thr	Ala	Asp	Val	Ile	Asn	Pro	Met	Ala	Leu	Arg
	50					55					60				
Gln	Arg	Glu	Glu	Leu	Arg	Glu	Lys	Leu	Ala	Ala	Ala	Lys	Glu	Lys	Arg
65					70				75						80
Leu	Leu	Asn	Gln	Lys	Leu	Gly	Lys	Ile	Lys	Thr	Leu	Gly	Glu	Asp	Asp
			85					90						95	
Pro	Trp	Leu	Asp	Asp	Thr	Ala	Ala	Trp	Ile	Glu	Arg	Ser	Arg	Gln	Leu
		100						105					110		
Gln	Lys	Glu	Lys	Asp	Leu	Ala	Glu	Lys	Arg	Ala	Lys	Leu	Leu	Glu	Glu
	115					120						125			
Met	Asp	Gln	Lys	Phe	Gly	Val	Ser	Thr	Leu	Val	Glu	Glu	Glu	Phe	Gly
130					135						140				
Gln	Arg	Arg	Gln	Asp	Leu	Tyr	Ser	Ala	Arg	Asp	Leu	Gln	Gly	Leu	Thr
145				150					155						160
Val	Glu	His	Ala	Ile	Asp	Ser	Phe	Arg	Glu	Gly	Glu	Thr	Met	Ile	Leu
			165					170					175		
Thr	Leu	Lys	Asp	Lys	Gly	Val	Leu	Gln	Glu	Glu	Glu	Asp	Val	Leu	Val
		180						185					190		
Asn	Val	Asn	Leu	Val	Asp	Lys	Glu	Arg	Ala	Glu	Lys	Asn	Val	Glu	Leu
	195						200					205			

Arg Lys Lys Lys Pro Asp Tyr Leu Pro Tyr Ala Glu Asp Glu Ser Val
 210 215 220
 Asp Asp Leu Ala Gln Gln Lys Pro Arg Ser Ile Leu Ser Lys Tyr Asp
 225 230 235 240
 Glu Lys Leu Glu Gly Glu Arg Pro His Ser Phe Arg Leu Glu Gln Gly
 245 250 255
 Gly Thr Ala Asp Gly Leu Arg Glu Arg Glu Leu Glu Glu Ile Arg Ala
 260 265 270
 Lys Leu Arg Leu Gln Ala Gln Ser Leu Ser Thr Val Gly Pro Arg Leu
 275 280 285
 Ala Ser Glu Tyr Leu Thr Pro Glu Glu Met Val Thr Phe Lys Lys Thr
 290 295 300
 Lys Arg Arg Val Lys Lys Ile Arg Lys Lys Glu Lys Glu Val Val Val
 305 310 315 320
 Arg Ala Asp Asp Leu Leu Pro Leu Gly Asp Gln Thr Gln Asp Gly Asp
 325 330 335
 Phe Gly Ser Arg Leu Arg Gly Arg Gly Arg Arg Arg Val Ser Glu Val
 340 345 350
 Glu Glu Glu Lys Glu Pro Val Pro Gln Pro Leu Pro Ser Asp Asp Thr
 355 360 365
 Arg Val Glu Asn Met Asp Ile Ser Asp Glu Glu Glu Gly Gly Ala Pro
 370 375 380
 Pro Pro Gly Ser Pro Gln Val Leu Glu Glu Asp Glu Ala Glu Leu Glu
 385 390 395 400
 Leu Gln Lys Gln Leu Glu Lys Gly Arg Arg Leu Arg Gln Leu Gln Gln
 405 410 415
 Leu Gln Gln Leu Arg Asp Ser Gly Glu Lys Val Val Glu Ile Val Lys
 420 425 430
 Lys Leu Glu Ser Arg Gln Arg Gly Trp Glu Glu Asp Glu Asp Pro Glu
 435 440 445
 Arg Lys Gly Ala Ile Val Phe Asn Ala Thr Ser Glu Phe Cys Arg Thr
 450 455 460
 Leu Gly Glu Ile Pro Thr Tyr Gly Leu Ala Gly Asn Arg Glu Glu Gln
 465 470 475 480
 Glu Glu Leu Met Asp Phe Glu Arg Asp Glu Glu Arg Ser Ala Asn Gly
 485 490 495
 Gly Ser Glu Ser Asp Gly Glu Glu Asn Ile Gly Trp Ser Thr Val Asn
 500 505 510
 Leu Asp Glu Glu Lys Gln Gln Gln Asp Val Arg Ala Thr Pro Leu Gly
 515 520 525

Gly Gly Arg Leu Gly Val Leu Lys Leu Glu Met Ser Thr Gly Leu Gly
530 535 540

Val Gln Ser Leu Ser Leu Leu Ile Gln Ser Gly Leu Cys Arg Pro Pro
545 550 555 560

Arg Ala Ile

<210> 177

<211> 1790

<212> DNA

<213> Homo sapiens

<400> 177

```

ctggtaccca cccctgtctc tggacttctt tatgaacact gagtatttga ctccccaccc 60
ttcatgggtca gctcctttta acaccctaga tatcattgct gtccttttca ttgacatgct 120
agagacattt ccaattttca tgggagtcag taagatagag atgatgaaa ttcagagtgt 180
tcaaagaaaag aggaggctc atcagttgat agaataaca acagattgtc taatctcagt 240
ttctttcact ttcttaccca ttacagtggg aaaaagaacc cttaagggaag ccagtgtctc 300
tcaacattgg ttacaaaacc atcagtgccg gcacttaatt taaaatcttc tgaaaatcca 360
tcagttgggt ttcatgtctc actctaaact cagctccaac caggctgctt cctagattct 420
tgtctgtgtg tgtttctgtt ttggctcttt gttacttctt tctgtaaact taagtaaact 480
cagtttgttg acctagtga ttgattcttt ccattttcta tactattcac cagcatatca 540
aagatgcttc ttccctaactc tgatttccag agtaactgtg ctatcagctt ttcataacta 600
gaactaggtt ttacactttt cacatttctt gcctcctacc cagttcgtaa gccaaaattag 660
actcgtactc ttgtgaagtg ccttttagat actagtgtaa aatatactgc gtaattctcc 720
attgccctac atatatatac cttttaagaa aatatactac cagtctattt tatcttcgca 780
gaacttgccc aggtatgagc cagcatttat taccactcat tctagaaggt gccagttagg 840
tagctatcca ggtttacatt ttaaggcagt cgagacctta ttcacgtggg atataaaca 900
ccacatttct cttttatatg gaacaagact ttctatttgc ggctggcctt ctcactagct 960
tatgtctttt ttttttaaga cctttcttag cactcgtgta actcctcccc ctcacaatca 1020
atctcagcaa ctcagcagag tcgcttcagt tcacagctct taatccttca ctccagacta 1080
aagctaactt tatgtcctca aacagctaca actcactgtt atcacagttc cgactgcaga 1140
gactccattt aagaggaaat ttaaaaaata aacaatgctc tatttctgtc cacatcaaag 1200
gtacctctaa cagaaaactg tctttgcttc tgtctctctg ctactggact ctcagctcaa 1260
ggtaatgtaa gatgatctca tagatgcact ttagcctctt gatgttagca gttgagctct 1320
ccctgcgcac aaaatacatt tgtggtaatg aatgcccgaa atctcactgg cacaccaag 1380
agtgtttaa ctaaaattgc tactaacaca acattgtaga attcgttgcc ttaatggatg 1440
tgatcttcta ggcttactgt gtcttaacaa gcagtcttat atatatatat cttgcccaga 1500
aagtacactt gtagaattta acatttgctc ctttctccaa acagattaaa gggtatgcag 1560
ctgaatgatt ctgtgtccca ttttaaaggt ccaggctatc ccctagattt agattagttc 1620
ctaaaaaatc ttccagtctg aatcctgaca aatggatagc taaattaatt tccactcctc 1680
ttctctcttg cactgtttgg tacattacag ccaccattag ccttaaaagc tgcagcagcc 1740
aaattcgggt acagtaacat aaagcctgag cttaaaaaaa aaaaaaaaaa 1790

```

<210> 178

<211> 115

<212> PRT

<213> Homo sapiens

<400> 178

```

Met Glu Gln Asp Phe Ser Phe Ala Ala Gly Leu Leu Thr Ser Leu Cys
  1              5              10             15

Phe Phe Phe Leu Arg Pro Phe Leu Ala Leu Ala Glu Leu Leu Pro Leu
  20              25             30

Thr Ile Asn Leu Ser Asn Ser Ala Glu Ser Leu Gln Phe Thr Ala Leu

```

35 40 45
 Asn Pro Ser Leu Gln Thr Lys Ala Asn Leu Met Ser Ser Asn Ser Tyr
 50 55 60
 Asn Ser Leu Leu Ser Gln Phe Arg Leu Gln Arg Leu His Leu Arg Gly
 65 70 75 80
 Asn Leu Lys Asn Lys Gln Cys Ser Ile Ser Val His Ile Lys Gly Thr
 85 90 95
 Ser Asn Arg Asn Leu Ser Leu Leu Leu Ser Leu Cys Tyr Trp Thr Leu
 100 105 110
 Ser Ser Arg
 115

<210> 179

<211> 2026

<212> DNA

<213> Homo sapiens

<400> 179

```

cattgagttg ccctaacaac attgtctcca gtgtcagaac catattaagg ttcgtttctc 60
agattgggag cctgcaacac catacagcca acattgcctt tgccacgcca ctgccaccat 120
ccccaccatt gccctatggt gggcagatga attccagaaa ccctcagggg gccaggataa 180
ttaggcaacc catctgaatt ggccacgtaa gtgacaggca cttatctctc gggttcttgc 240
ttttgcagac tccaggggaag tcctgtctag aggtcgatgg cagagactcc tactctttcc 300
catgaggggt tgataggaat caaattggga ttcttttggc ttggggtttt gtttttttgt 360
tgttgttttt ggttttcagt ttgttttttg gtgtatgggg ggtgatcttg tttctgaata 420
agaaaaagaa gaggcaacca tggcccttat gtgggtttat cctttttgag caatgtttta 480
gccacaagta aggaatcttg aaagtctttt gtccagcaag cagtcttaaa aatgtttttc 540
ctaactcctt ttgcaggtga ctaagtacaa aaaaatagtt ttctcattgt attcaaaata 600
gtgagtaggt tccttgata atacacagtg gtagttagca ttttttctca aaacacaacc 660
agaaaaccca ctctcggtat ttgtaaatca cttttcaagg gaaaaagtga acacgtattc 720
cttgatttcc tagtttgatt accaaacctg atgttacaaa gaaacctccg ttctgtagac 780
agaatttctt ttatttttct tcttttactc ctcaaatca ctttccaggt gccatcacca 840
tctataaggt ctgagagcag aggtattatc atggtataaa gtgggggtgt ggtgcagcca 900
ttccagtaac acccacaaga ggacagctgt tctgaatgtc cccaccacc cctctttcag 960
tacaggtgag acattttcag ttcatgagct ccagaccaa tcccaggcca gcccttgcac 1020
caaaagcctt ttttagaagg cttatcagtc tattaggaat gtctcaggaa agatgagcca 1080
tttctttggg gagaaatata ttacagatg gaagtgtgtg actgcgtgtc tgtgtgtgtg 1140
tgtgtgtgtg gtgcgacgt gagtgcgtgt gttcatctat gtgcatttca cttccataaa 1200
gacccagccc aagctgctgg gaaccatgtt ttctgagta ttctcagagg ttaaacaagt 1260
gacaagttag cttctgaaat tagtgtctca gcaagctggc tttaggaatg agccccattt 1320
tatcaagcag agaaaaaaaa taacagcaga aaagataaag ataaaccaa aatatatacc 1380
ccccaatgga aaataatgtt gattcagcaa ttccatagg atgtattaca tgctctaatt 1440
tattatatta ttatttatct gtctttgac tttgccatt gtactcttaa aaagatgttg 1500
ggatgttgat tgcgattttt aaacaactag ataatgtata aatcagcagt ggaaatcagt 1560
tttaatgtgt ggatgtgtct gattattgtt aaatgcctct ttttttactt tttttttttt 1620
tagatgtata atgtttcata aaccctggca ctggtcacaa agctcagctg tgaaaatgaa 1680
attttagtag tttttaaaca tgaatgtcaa ttccaagtgt atttgaaatg gttcctccag 1740
gagagatatt tgtgcaccat taggaaaatc ttctctgcag aggaagttag cttcttttga 1800
gaaaatggaa aatgggttct gatatgtgat ctgagatag cccatttctt agggcaccat 1860
ggaaaacaca aatgtgatct ttaagtatac ctcttcccca gtttggggag gaaaggactc 1920
agtttgcacc ctttttgtat gtaaaataaa atgtcttacc tttcttggtt aaaaaaaaaa 1980
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 2026

```

<210> 180

<211> 52
 <212> PRT
 <213> Homo sapiens

<400> 180
 Met Leu Gln Arg Asn Leu Arg Ser Val Asp Arg Ile Ser Phe Ile Phe
 1 5 10 15
 Leu Leu Leu Leu Leu Thr Ile Thr Phe Pro Val Pro Ser Pro Ser Ile
 20 25 30
 Arg Ser Gln Ser Arg Gly Leu Phe Met Val Ile Ser Gly Gly Val Val
 35 40 45
 Gln Pro Phe Gln
 50

<210> 181
 <211> 1138
 <212> DNA
 <213> Homo sapiens

<400> 181
 ggggtctgtgg agagccgggt gcgagcggcg gcagcacgag gggaaaagag ctgagcggag 60
 accaaagtca gccgggagac agtgggtctg tgagagaccg aatagagggg ctggggccac 120
 gagcgccatt gacaagcaat ggggaagaaa cagaaaaaca agagcgaaga cagcaccaag 180
 gatgacattg atcttgatgc cttggctgca gaaatagaag gagctggtgc tgccaaagaa 240
 caggagcctc aaaagtcaaa agggaaaaag aaaaaagaga aaaaaaagca ggactttgat 300
 gaagatgata tcctgaaaga actggaagaa ttgtctttgg aagctcaagg catcaaagct 360
 gacagagaaa ctggttcagc gaagccaaca gaaaacaatg aagaggaatt cacctcaaaa 420
 gataaaaaaa agaaaggaca gaagggcaaa aaacagagtt ttgatgataa tgatagcgaa 480
 gaattggaag ataaagattc aaaatcaaaa aagactgcaa aaccgaaagt ggaaatgtac 540
 tctgggagtg atgatgatga tgattttaac aaacttccta aaaaagctaa agggaaagct 600
 caaaaatcaa ataagaagtg ggatgggtca gaggaggatg aggataacag taaaaaatt 660
 aaagagcggt caagaataaa ttcttctggt gaaagtgggt atgaatcaga tgaatttttg 720
 caatctaaaa ggacagaaaa aaaatcagaa aaacaagcca ggtcctaaca tagaaagtgg 780
 gaatgaagat gatgacgct ctttcaaat taagacagtg gccccaaaga aggcagaaaa 840
 gaaggagcgc gagagaaaaa agcgagatga agaaaagcg aaactgcgga agctgaaaga 900
 aaaagaagag ttagaaacag gtaaaaagga tcagagtaaa caaaaggaat ctcaaaggaa 960
 atttgaagaa gaaactgtaa aatccaaagt gactgttgat actggagtaa ttcctgcctc 1020
 tgaagagaaa gcagagactc ccacagctgc agaagatgac aatgaaggag acaaaaagaa 1080
 gaaagataag aagaaaaaga aaggagaaaa ggaagaaaaa gagaaaaaaa aaaaaaaa 1138

<210> 182
 <211> 209
 <212> PRT
 <213> Homo sapiens

<400> 182
 Met Gly Lys Lys Gln Lys Asn Lys Ser Glu Asp Ser Thr Lys Asp Asp
 1 5 10 15
 Ile Asp Leu Asp Ala Leu Ala Ala Glu Ile Glu Gly Ala Gly Ala Ala
 20 25 30
 Lys Glu Gln Glu Prc Gln Lys Ser Lys Gly Lys Lys Lys Lys Glu Lys
 35 40 45
 Lys Lys Gln Asp Phe Asp Glu Asp Asp Ile Leu Lys Glu Leu Glu Glu

50 55 60
 Leu Ser Leu Glu Ala Gln Gly Ile Lys Ala Asp Arg Glu Thr Val Ala
 65 70 75 80
 Val Lys Pro Thr Glu Asn Asn Glu Glu Glu Phe Thr Ser Lys Asp Lys
 85 90 95
 Lys Lys Lys Gly Gln Lys Gly Lys Lys Gln Ser Phe Asp Asp Asn Asp
 100 105 110
 Ser Glu Glu Leu Glu Asp Lys Asp Ser Lys Ser Lys Lys Thr Ala Lys
 115 120 125
 Pro Lys Val Glu Met Tyr Ser Gly Ser Asp Asp Asp Asp Phe Asn
 130 135 140
 Lys Leu Pro Lys Lys Ala Lys Gly Lys Ala Gln Lys Ser Asn Lys Lys
 145 150 155 160
 Trp Asp Gly Ser Glu Glu Asp Glu Asp Asn Ser Lys Lys Ile Lys Glu
 165 170 175
 Arg Ser Arg Ile Asn Ser Ser Gly Glu Ser Gly Asp Glu Ser Asp Glu
 180 185 190
 Phe Leu Gln Ser Lys Arg Thr Glu Lys Lys Ser Glu Lys Gln Ala Arg
 195 200 205
 Ser

<210> 183
 <211> 912
 <212> DNA
 <213> Homo sapiens

<400> 183
 atccagctct ttcagagaga agagatggga ataagataag aacatcattt cctgaagagt 60
 ttactaagtt gatgttttct tttggtgaag ggcaatgttg aatttaaatac atatagttca 120
 tgtgtgcacc tgtcctgcag tttgaggcag atgattttta accagcatct atagacactt 180
 cctgtgaagg agagcttcaa gttggcaaag gagatgaagt cacaattaca ctgccacata 240
 tccctagctg agggcagcag taaaatccag gcccgaaatgg aacagcagcc cactcgtcct 300
 ccacagacgt cacagccacc accacctcca ccacctatgc cattcagagc tccaacgaag 360
 cctccagttg gacccaaaac ttctcccttg aaagataaacc cgtcacctga acctcagttg 420
 gatgacatca aaagagagct gagggctgaa gttgacatta ttgaacaaat gagcagcagc 480
 agtgggagca gctcttcaga ctctgagagc tcttcgggaa gtgatgacga tagctccagc 540
 agtggaggsg aggamaatgg cccagcytct cytccgcagc yttmacacca gcagccytac 600
 aacagtaggc ctgccgttgc caatggaacc agccggccac aaggaagcaa ccagytyatg 660
 aacaccctca gaaatgactt gcagttgagt gagtytgga gtgacagtga tgactagtgc 720
 tggatytttc gaaacctact ttttggtgca caaacatgcc gcaagactga gctacttttg 780
 ccgtggagtc cattgcaaga ggaaaatggt atggatcagt gactgtagta ggagtttgag 840
 gctytggaac tctcacatat tcaagtcttt aacttagtggt tgatgggtga aaaaaaaaaa 900
 aaaaaaaaaa aa 912

<210> 184
 <211> 167
 <212> PRT
 <213> Homo sapiens

```
<210> 185
<211> 4582
<212> DNA
<213> Homo sapiens
```

<400> 185

```

ggggaagcta atcaaactctg gttcctatgt agctctcact gttcagggac gcccacctgg 60
gtcgccccag attccacttg ccgactctga agtagagccg tcagtcattg gacatatgtc 120
tcccatcatg acatctcttc attcaccttg agcatctggg aatatggaga gaatcactag 180
tcctgtgtctc atggggggagg aaaacaatgt ggttcataac cagaaagtag aaattctgag 240
aaaaatgtta cagaaagaac aggaacggct acagttattg caggaagatt acaaccgaac 300
acctgccccaa agattgtctaa aagagatcca agaggccaag aaacacattc ctccagctgca 360
agagcaggtta tccaaagcca caggctctgc tcaggatgga gctgtagtta caccctccag 420
accttttaggg gacaccctaa cagtcagtga ggcagaaaca gatcctggag atgtactggg 480
caggactgac tgtagcagtg gagatgcttc tcggcccagt agtgacaatg cagatagtcc 540
caagagtggc ccaaaagaga gaatttatct agaggaaaac ccagagaaaa gtgaaacaat 600
tcaggacact gacactcaat cacttgctcg aagtccctca acccgatatg cacctcatat 660
tattggagca gaagatgatg attttggtac tgaacatgaa cagatcaatg gacagtgcag 720
ctgtttccag agcattgaaat tactaaaaac tcgcccggct catttggctg ttttcttaca 780
ccatgtagtgt tcacaatttg accctgcgac tttgctctgt tatctctatt cagacctgta 840
taaacatacc aattccaaag aaactcgtcg catcttcctt gaggtttcac agttctttct 900
aaatcgatca gcacacctga aagtttctgt tcctgatgaa atgtctgcag atctagaaaa 960
gagaagacct gagctcattc ctgaggatct gcacgcgcc tatatccaaa ctatgcaaga 1020
aagagtcctat ccagaagttc aaaggcactt aaaagatttt cggcagaaac gtagtatggg 1080
actgaccttg gctgaaaagg agctgactaa acttgatgca gagcgagaca aggaccgatt 1140
gacttttgagg aaggagcggg catgtgcaga acagattgtt gccaaaattg aagaagtatt 1200
gatgaytgct caggctgtar aggaagataa gagctccacc atgcagtatg ttattctcat 1260
gtatatgaag catttgaggag taaaagtga agagcctcga aatttgagc acaaacgggg 1320
tcggattgga tttcttccca aaatcaagca aagtatgaag aaagataaag aaggggaaga 1380
aaaagggaag cgaagaggat tccccagcat cctgggaccc ccacggagac caagccgtca 1440
tgacaacagt gcaattggca gagccatgga actacagaag gcgcgccacc ctaagcactt 1500
atccacaccc tcactgtgga gtccatgaacc tcaggactct gccaaagttg gccagagtgg 1560
gttagcaaat gaaggaacag acgctggata cctgcctgcc aattccatgt cttctgtagc 1620
ttcaggggccc tctttttccc aggaaggagg gaaagagaat gatacaggat caaagcaagt 1680
tgagaaaca tcagcacctg gagcacacct agatggcaca cctcgactc tcaatactgt 1740
ctttgatctc ccaccacctc cattagacca agtgcaggag gaggaatgtg aagtagaaag 1800
gggtgactga catgggacac caaagccctt tcgaaagttt gacagtgtag cttttggaga 1860
aagtcaaatg gaggatgaac aatttgaaaa tgacttagag acagatccac ccaactggca 1920
gcagcttggt agtcgagaag tgttactggg actaaaacct tgtgaaatca aaagacagga 1980
agtgattaat gaattgttct acactgaaag agctcatgtt cgaacactga aggttcttga 2040
tcaagtgttc tatcagcgag tatccagaga aggaattctg tcaccctcag agctacggaa 2100
aattttttca aacttggaag atattcttca acttcacatt ggattgaatg aacaaatgaa 2160
ggctgttcca aagagaaatg agacctctgt tatcgatcag attggggaag atttgcctgac 2220
atggttcacg ggaccaggag aggagaaatt gaaacatgct gctgctacct tttgcagtaa 2280
ccaaccttcc gccctggaaa tgatcaaatc tcgtcagaaa aaggattctc gatttcagac 2340
ttttgtgcaa gatgctgaaa gtaatccact gtgtcgtcgt cttcaactga aggatattat 2400
tcccactcaa atgcaaaggc ttactaagta ccacttctg ttggataata ttgccaaata 2460
cacagaatgg ccaacagaaa gggagaagggt gaagaaagct gcagatcact gtcgtcagat 2520
cttaaatat gtaaatcagg ctgtcaagga ggcagaaaac aagcagcgcc tagaagatta 2580
tcagcgtcgc cttgatacct ccagcctgaa gttgtcagag taccctaatg ttgaagagct 2640
caggaatttg gatttaacaa aaaggaagat gattcatgaa gggccattgg tttggaagg 2700
gaatagagat aaaactattg atttatcac gttgctgctg gaagacattc ttgtattgtt 2760
acaaaagcag gatgatagac tggttttaag gtgtcatagt aagattctgg catctacagc 2820
tgatagcaaa cacacgttta gccctgtcat taagttagt acagtgttg ttcgacaagt 2880
ggcaacagat aacaaagctt tattcgtcat ttccatgtca gacaatggcg ctccagattta 2940
tgaactgggt gcacagacag tttctgaaaa gactgtctgg caggacctaa tctgtcggat 3000
ggctgcatca gtgaaggagc aatccacaaa gccgattcca ttaccacagt caacacctgg 3060
cgaaggagat aatgatgaag aagatccttc aaaattaaaa gaggagcagc atggcatttc 3120
agtcactggg ttgcagagtc cagacagaga tttgggatta gaatctacct taatatcgtc 3180
aaaacctcag tctcattcac tgagtacctc tgggaaatca gaggtacgtg atctgtttgt 3240
ggctgagaga cagtttgcaa aggaacaaca tacagatggg acactaaagg aagttggaga 3300
agattatcaa atcgcaatcc cagattcaca cctgcctgtc tcagaagaac ggtgggcatt 3360
ggatgcacta agaaatttg gtttgttgaa gcagtgtgtg gtgcaacagc taggtttgac 3420
tgagaagagc gttcaggaag actggcaaca ttcccaaga tacagaacag cctctcaggg 3480

```

```

gccgcagaca gacagtgtca tccagaactc tgaaaatatt aaggcctatc attctggtga 3540
aggacatatg ccccttagaa ctggaactgg tgacattgca acttggtaca gtccacggac 3600
ttcaactgaa tcttttgctc cacgggatcc agtgggactg gcaccccagg atagccaggc 3660
aagtaacatt ttagtaatgg accacatgat tatgaccca gagatgccta ccatggagcc 3720
agaagggggt cttgatgaca gtggagagca cttttttgat gcccggaag cacatagtga 3780
tgagaatcca tcagaagggt atggagcagt taacaaggaa gagaaggatg ttaatttacg 3840
catctcagga aactatttga tccttgatgg ctatgaccca gtgcaggaga gttccacaga 3900
tgaggaggtt gcttcctcac ttaccctgca gcccatgaca ggcatccctg ctgtggaatc 3960
caccaccag cagcaacatt ctctcagaa tactcactcc gatggggcaa tttcaccatt 4020
caccgccgaa tttctggtcc agcagcgctg gggagctatg gagtattcct gttttgagat 4080
ccagagtccc tcctcttggt cagattcaca gagccagatc atggagtaca ttcataagat 4140
agaggctgac cttgaacact taaaggaagg tggaggaaag ttaacaccat tctttgcaa 4200
aggctggctg gatcagccct cacagacaag cactcagata aaagttagag ccgcatgtcc 4260
tggaggtgac tgcaggttgt tggatttggg gtatcgggcg tgtctacca catcctggct 4320
ccagtgtgga tgcagagaga gtgtgacaga ggatctgcct gtgaaccacc tgggattagt 4380
caagtcccaa ggtgcccaga gtgggactag ttyttcacag tgtggcagct gcactaatct 4440
gtttgtgagg gaatatccat tccctcactc tactctcctc actatcgaa attcattttg 4500
attcagaata aaaaccaa gtatagagct ttgggttcta ggatatgaaa ttgtacttag 4560
atttaagaaa aaaaaaaaaa aa 4582

```

<210> 186

<211> 1461

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (364)

<220>

<221> UNSURE

<222> (369)

<220>

<221> UNSURE

<222> (1433)

<400> 186

```

Met Ser Pro Ile Met Thr Ser Pro His Ser Pro Gly Ala Ser Gly Asn
  1                      5                      10                      15

```

```

Met Glu Arg Ile Thr Ser Pro Val Leu Met Gly Glu Glu Asn Asn Val
      20                      25                      30

```

```

Val His Asn Gln Lys Val Glu Ile Leu Arg Lys Met Leu Gln Lys Glu
      35                      40                      45

```

```

Gln Glu Arg Leu Gln Leu Leu Gln Glu Asp Tyr Asn Arg Thr Pro Ala
      50                      55                      60

```

```

Gln Arg Leu Leu Lys Glu Ile Gln Glu Ala Lys Lys His Ile Pro Gln
      65                      70                      75                      80

```

```

Leu Gln Glu Gln Leu Ser Lys Ala Thr Gly Ser Ala Gln Asp Gly Ala
      85                      90                      95

```

```

Val Val Thr Pro Ser Arg Pro Leu Gly Asp Thr Leu Thr Val Ser Glu
      100                      105                      110

```

```

Ala Glu Thr Asp Pro Gly Asp Val Leu Gly Arg Thr Asp Cys Ser Ser

```


115	120	125
Gly Asp Ala Ser Arg Pro Ser Ser Asp Asn Ala Asp Ser Pro Lys Ser		
130	135	140
Gly Pro Lys Glu Arg Ile Tyr Leu Glu Glu Asn Pro Glu Lys Ser Glu		
145	150	155
Thr Ile Gln Asp Thr Asp Thr Gln Ser Leu Val Gly Ser Pro Ser Thr		
165	170	175
Arg Ile Ala Pro His Ile Ile Gly Ala Glu Asp Asp Asp Phe Gly Thr		
180	185	190
Glu His Glu Gln Ile Asn Gly Gln Cys Ser Cys Phe Gln Ser Ile Glu		
195	200	205
Leu Leu Lys Ser Arg Pro Ala His Leu Ala Val Phe Leu His His Val		
210	215	220
Val Ser Gln Phe Asp Pro Ala Thr Leu Leu Cys Tyr Leu Tyr Ser Asp		
225	230	235
Leu Tyr Lys His Thr Asn Ser Lys Glu Thr Arg Arg Ile Phe Leu Glu		
245	250	255
Phe His Gln Phe Phe Leu Asn Arg Ser Ala His Leu Lys Val Ser Val		
260	265	270
Pro Asp Glu Met Ser Ala Asp Leu Glu Lys Arg Arg Pro Glu Leu Ile		
275	280	285
Pro Glu Asp Leu His Arg His Tyr Ile Gln Thr Met Gln Glu Arg Val		
290	295	300
His Pro Glu Val Gln Arg His Leu Lys Asp Phe Arg Gln Lys Arg Ser		
305	310	315
Met Gly Leu Thr Leu Ala Glu Ser Glu Leu Thr Lys Leu Asp Ala Glu		
325	330	335
Arg Asp Lys Asp Arg Leu Thr Leu Glu Lys Glu Arg Thr Cys Ala Glu		
340	345	350
Gln Ile Val Ala Lys Ile Glu Glu Val Leu Met Xaa Ala Gln Ala Val		
355	360	365
Xaa Glu Asp Lys Ser Ser Thr Met Gln Tyr Val Ile Leu Met Tyr Met		
370	375	380
Lys His Leu Gly Val Lys Val Lys Glu Pro Arg Asn Leu Glu His Lys		
385	390	395
Arg Gly Arg Ile Gly Phe Leu Pro Lys Ile Lys Gln Ser Met Lys Lys		
405	410	415
Asp Lys Glu Gly Glu Glu Lys Gly Lys Arg Arg Gly Phe Pro Ser Ile		
420	425	430
Leu Gly Pro Pro Arg Arg Pro Ser Arg His Asp Asn Ser Ala Ile Gly		

435	440	445
Arg Ala Met Glu Leu Gln Lys	Ala Arg His Pro Lys	His Leu Ser Thr
450	455	460
Pro Ser Ser Val Ser Pro Glu Pro Gln Asp Ser Ala Lys Leu Arg Gln		
465	470	475 480
Ser Gly Leu Ala Asn Glu Gly Thr Asp Ala Gly Tyr Leu Pro Ala Asn		
485	490	495
Ser Met Ser Ser Val Ala Ser Gly Ala Ser Phe Ser Gln Glu Gly Gly		
500	505	510
Lys Glu Asn Asp Thr Gly Ser Lys Gln Val Gly Glu Thr Ser Ala Pro		
515	520	525
Gly Asp Thr Leu Asp Gly Thr Pro Arg Thr Leu Asn Thr Val Phe Asp		
530	535	540
Phe Pro Pro Pro Pro Leu Asp Gln Val Gln Glu Glu Glu Cys Glu Val		
545	550	555 560
Glu Arg Val Thr Glu His Gly Thr Pro Lys Pro Phe Arg Lys Phe Asp		
565	570	575
Ser Val Ala Phe Gly Glu Ser Gln Ser Glu Asp Glu Gln Phe Glu Asn		
580	585	590
Asp Leu Glu Thr Asp Pro Pro Asn Trp Gln Gln Leu Val Ser Arg Glu		
595	600	605
Val Leu Leu Gly Leu Lys Pro Cys Glu Ile Lys Arg Gln Glu Val Ile		
610	615	620
Asn Glu Leu Phe Tyr Thr Glu Arg Ala His Val Arg Thr Leu Lys Val		
625	630	635 640
Leu Asp Gln Val Phe Tyr Gln Arg Val Ser Arg Glu Gly Ile Leu Ser		
645	650	655
Pro Ser Glu Leu Arg Lys Ile Phe Ser Asn Leu Glu Asp Ile Leu Gln		
660	665	670
Leu His Ile Gly Leu Asn Glu Gln Met Lys Ala Val Arg Lys Arg Asn		
675	680	685
Glu Thr Ser Val Ile Asp Gln Ile Gly Glu Asp Leu Leu Thr Trp Phe		
690	695	700
Ser Gly Pro Gly Glu Glu Lys Leu Lys His Ala Ala Ala Thr Phe Cys		
705	710	715 720
Ser Asn Gln Pro Phe Ala Leu Glu Met Ile Lys Ser Arg Gln Lys Lys		
725	730	735
Asp Ser Arg Phe Gln Thr Phe Val Cln Asp Ala Glu Ser Asn Pro Leu		
740	745	750
Cys Arg Arg Leu Gln Leu Lys Asp Ile Ile Pro Thr Gln Met Gln Arg		

755	760	765
Leu Thr Lys Tyr Pro Leu Leu Leu Asp Asn Ile Ala Lys Tyr Thr Glu 770 775 780		
Trp Pro Thr Glu Arg Glu Lys Val Lys Lys Ala Ala Asp His Cys Arg 785 790 795 800		
Gln Ile Leu Asn Tyr Val Asn Gln Ala Val Lys Glu Ala Glu Asn Lys 805 810 815		
Gln Arg Leu Glu Asp Tyr Gln Arg Arg Leu Asp Thr Ser Ser Leu Lys 820 825 830		
Leu Ser Glu Tyr Pro Asn Val Glu Glu Leu Arg Asn Leu Asp Leu Thr 835 840 845		
Lys Arg Lys Met Ile His Glu Gly Pro Leu Val Trp Lys Val Asn Arg 850 855 860		
Asp Lys Thr Ile Asp Leu Tyr Thr Leu Leu Leu Glu Asp Ile Leu Val 865 870 875 880		
Leu Leu Gln Lys Gln Asp Asp Arg Leu Val Leu Arg Cys His Ser Lys 885 890 895		
Ile Leu Ala Ser Thr Ala Asp Ser Lys His Thr Phe Ser Pro Val Ile 900 905 910		
Lys Leu Ser Thr Val Leu Val Arg Gln Val Ala Thr Asp Asn Lys Ala 915 920 925		
Leu Phe Val Ile Ser Met Ser Asp Asn Gly Ala Gln Ile Tyr Glu Leu 930 935 940		
Val Ala Gln Thr Val Ser Glu Lys Thr Val Trp Gln Asp Leu Ile Cys 945 950 955 960		
Arg Met Ala Ala Ser Val Lys Glu Gln Ser Thr Lys Pro Ile Pro Leu 965 970 975		
Pro Gln Ser Thr Pro Gly Glu Gly Asp Asn Asp Glu Glu Asp Pro Ser 980 985 990		
Lys Leu Lys Glu Glu Gln His Gly Ile Ser Val Thr Gly Leu Gln Ser 995 1000 1005		
Pro Asp Arg Asp Leu Gly Leu Glu Ser Thr Leu Ile Ser Ser Lys Pro 1010 1015 1020		
Gln Ser His Ser Leu Ser Thr Ser Gly Lys Ser Glu Val Arg Asp Leu 1025 1030 1035 1040		
Phe Val Ala Glu Arg Gln Phe Ala Lys Glu Gln His Thr Asp Gly Thr 1045 1050 1055		
Leu Lys Glu Val Gly Glu Asp Tyr Gln Ile Ala Ile Pro Asp Ser His 1060 1065 1070		
Leu Pro Val Ser Glu Glu Arg Trp Ala Leu Asp Ala Leu Arg Asn Leu		

1075	1080	1085
Gly Leu Leu Lys Gln Leu Leu Val Gln Gln Leu Gly Leu Thr Glu Lys		
1090	1095	1100
Ser Val Gln Glu Asp Trp Gln His Phe Pro Arg Tyr Arg Thr Ala Ser		
1105	1110	1115 1120
Gln Gly Pro Gln Thr Asp Ser Val Ile Gln Asn Ser Glu Asn Ile Lys		
	1125	1130 1135
Ala Tyr His Ser Gly Glu Gly His Met Pro Phe Arg Thr Gly Thr Gly		
	1140	1145 1150
Asp Ile Ala Thr Cys Tyr Ser Pro Arg Thr Ser Thr Glu Ser Phe Ala		
	1155	1160 1165
Pro Arg Asp Ser Val Gly Leu Ala Pro Gln Asp Ser Gln Ala Ser Asn		
	1170	1175 1180
Ile Leu Val Met Asp His Met Ile Met Thr Pro Glu Met Pro Thr Met		
1185	1190	1195 1200
Glu Pro Glu Gly Gly Leu Asp Asp Ser Gly Glu His Phe Phe Asp Ala		
	1205	1210 1215
Arg Glu Ala His Ser Asp Glu Asn Pro Ser Glu Gly Asp Gly Ala Val		
	1220	1225 1230
Asn Lys Glu Glu Lys Asp Val Asn Leu Arg Ile Ser Gly Asn Tyr Leu		
	1235	1240 1245
Ile Leu Asp Gly Tyr Asp Pro Val Gln Glu Ser Ser Thr Asp Glu Glu		
	1250	1255 1260
Val Ala Ser Ser Leu Thr Leu Gln Pro Met Thr Gly Ile Pro Ala Val		
1265	1270	1275 1280
Glu Ser Thr His Gln Gln Gln His Ser Pro Gln Asn Thr His Ser Asp		
	1285	1290 1295
Gly Ala Ile Ser Pro Phe Thr Pro Glu Phe Leu Val Gln Gln Arg Trp		
	1300	1305 1310
Gly Ala Met Glu Tyr Ser Cys Phe Glu Ile Gln Ser Pro Ser Ser Cys		
	1315	1320 1325
Ala Asp Ser Gln Ser Gln Ile Met Glu Tyr Ile His Lys Ile Glu Ala		
	1330	1335 1340
Asp Leu Glu His Leu Lys Glu Gly Gly Gly Lys Leu Thr Pro Phe Phe		
1345	1350	1355 1360
Ala Lys Gly Trp Leu Asp Gln Pro Ser Gln Thr Ser Thr Gln Ile Lys		
	1365	1370 1375
Val Arg Ala Ala Cys Pro Gly Gly Asp Cys Arg Leu Leu Asp Leu Glu		
	1380	1385 1390
Tyr Arg Pro Cys Leu Thr Thr Ser Trp Leu Gln Cys Gly Cys Arg Glu		

1395 1400 1405
 Ser Val Thr Glu Asp Leu Pro Val Asn His Leu Gly Leu Val Lys Ser
 1410 1415 1420
 Gln Gly Ala Gln Ser Gly Thr Ser Xaa Ser Gln Cys Gly Ser Cys Thr
 1425 1430 1435 1440
 Asn Leu Phe Val Arg Glu Tyr Pro Phe Pro His Ser Thr Leu Leu Thr
 1445 1450 1455
 Ile Gly Asn Ser Phe
 1460

<210> 187

<211> 2837

<212> DNA

<213> Homo sapiens

<400> 187

tcttttgaca gatataattg ctgcttatca aagattctgt tctcgacccc caaaaggatt 60
 tgaaaaatac tttcctaattg gaaaaaatgg aaaaaaagct agtgaacctt aagaagttat 120
 gggagagaaa aaagaatcaa agccagctgc taccacacgc tcttctggag gaggagggtg 180
 ttgcggtgga aaacgaggtg gcaagaaaaga tgattctcac tgggtggcca gggttcagaa 240
 ggggtgacatt ccatgggacg acaaggattt caggatgttc ttctcttga ctgctctgtt 300
 ctgggggtgga gtcatgtttt acttgctgct caagagatcc gggagagaaa tcaacttgaa 360
 ggactttgtc aataactatc tttcaaaagg agtagtagac agattggaag tcgtcaacaa 420
 gcgttttgtt cgagtgcctt ttacaccagg aaaaactcct gttgatgggc aatacgtttg 480
 gtttaatatg ggcagtgttg acacctttga acggaatctg gaaactttac agcaggaatt 540
 gggcatagaa ggagaaaatc ggtgtcctgt tgtctacatt gctgaaagt atggctcttt 600
 tctgctgagc atgtctgcct cgtgctcat catcgccctc ttgctctaca ccatcagaag 660
 agggcctgct ggcattggcc ggacaggccg agggatgggc ggactcttca gtgtcggaga 720
 aaccactgcc aaggtcttaa aggatgaaat tgatgtgaag ttcaaagatg tggctggctg 780
 tgaggaggcc aagctagaga tcatggaatt tgtgaatttc ttgaaaaacc caaagcagta 840
 tcaagacctt ggagcaataa tcccaaaggg tgccattctc actggtcctc caggcactgg 900
 gaagacgtg ctagttaagg ccacagccgg agaagccaat gtcccttca tcaccgttag 960
 tggatctgag tttttggaga tgttcgcttg tgtgggccct gctagagtc gagacttatt 1020
 tgcccttgct cggaagaatg ccccttgcat cctcttcac gatgaaatcg atgcggtggg 1080
 aaggaagaga ggaagaggca actttggagg gcagagttag caggagaaca cactcaacca 1140
 gctgctggtg gatagtgatg gttttaatac aacaacaaat gtcgtcattt tggccggcac 1200
 caatcgacca ggaccaccag acataaaagg aagagcttct attttcaaag ttcatctccg 1260
 accgctaaaa ctggacagta ccctggagaa ggataaattg gcaagaaaac tggcatcttt 1320
 aactccaggg ttttcaggtg ctgatgttgc taatgtctgt aatgaagctg cgttgattgc 1380
 tgcaaggcat ctgtcagatt ccataaatca gaaacacttt gaacaggcaa ttgagcgagt 1440
 gatttggtgc ttaaagaaaa aaacgcaggt tctgcagcct gaggagaaaa agactgtggc 1500
 ataccacgaa gcaggccatg cgttgcccg ctggtatctg gagcacgcag acccgctttt 1560
 aaaggtatcc atcatccac gtggcacaagg actaggttat gctcagtatt taccaaaaga 1620
 acaatacttc tataccaaag agcagctctt ggataggatg tgtatgactt taggtggtcg 1680
 agtctctgaa gaaatcttct ttggaagaat tacaactggt gctcaagatg acttgagaaa 1740
 agtaactcag agtgcatatg cccaaattgt tcagtttggc atgaatgaaa aggttgggca 1800
 aatctctttt gacctccac gtcaggggga catggtattg gagaaacctt acagtgaagc 1860
 cactgcaaga ttgatagatg atgaagtacg aatacttatt aatgatgctt ataaaagaac 1920
 agtagctctt ctacagaaa agaaagctga cgtggagaag gttgctcttc tgttgtaga 1980
 aaaagaagta ttagataaga atgatatggt tgaacttttg gggccagac catttgcgga 2040
 aaaatctacc tatgaagaat ttgtggaagg cactggcagc ttggatgagg acacctcact 2100
 tccagaaggc cttaaggat ggaacaagg gcgggaaaag gagaagagg agccccggg 2160
 tgagaaagtt gccaactaga gggcagagg gagggccatc cagtctgtcc actgtgggtt 2220
 cagctggtgc attatttcag ctgtggcttt cagaagaatg ggaatgctgc gctgatttta 2280
 gccagccact ggccagctg aaatgatggg gaaaggagtc cttagtcctt tcagcctcag 2340

```

aggtcacagt ggggtggcagg tgactttccg gaggccttga gggaaatgca cactgtccca 2400
tagcctcatt gggttcccag acgtgctgga aaggttgagc ccagagtggc cgaggctgga 2460
ccctgtggca ccaagtgggg tcggctgacc gtgtggcagg gattgttgca mtggawtttt 2520
ggcgtgtggg aagggatgct ttttttttgc cgcccayttt tcattcctgt ttttcctcag 2580
ttccccckkg cagatgggct gtgaaattaa attggagttt tgataagaac attttaattt 2640
gacttaatat tttaaagatt gaatccagat cacttggtgc tgttttaatg gaatggtttt 2700
ttacaggagc tgaacatam ttaaaaatat gaatgtatta tgtaaatatg gcttctttac 2760
ataaaaaata aaatgtcaac actgtaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2820
aaaaaaaaaa aaaaaaa 2837

```

<210> 188

<211> 686

<212> PRT

<213> Homo sapiens

<400> 188

```

Met Gly Glu Lys Lys Glu Ser Lys Pro Ala Ala Thr Thr Arg Ser Ser
  1              5              10              15

Gly Gly Gly Gly Gly Gly Gly Gly Lys Arg Gly Gly Lys Lys Asp Asp
      20              25              30

Ser His Trp Trp Ser Arg Phe Gln Lys Gly Asp Ile Pro Trp Asp Asp
      35              40              45

Lys Asp Phe Arg Met Phe Phe Leu Trp Thr Ala Leu Phe Trp Gly Gly
      50              55              60

Val Met Phe Tyr Leu Leu Leu Lys Arg Ser Gly Arg Glu Ile Thr Trp
      65              70              75              80

Lys Asp Phe Val Asn Asn Tyr Leu Ser Lys Gly Val Val Asp Arg Leu
      85              90              95

Glu Val Val Asn Lys Arg Phe Val Arg Val Thr Phe Thr Pro Gly Lys
      100              105              110

Thr Pro Val Asp Gly Gln Tyr Val Trp Phe Asn Ile Gly Ser Val Asp
      115              120              125

Thr Phe Glu Arg Asn Leu Glu Thr Leu Gln Gln Glu Leu Gly Ile Glu
      130              135              140

Gly Glu Asn Arg Val Pro Val Val Tyr Ile Ala Glu Ser Asp Gly Ser
      145              150              155              160

Phe Leu Leu Ser Met Leu Pro Thr Val Leu Ile Ile Ala Phe Leu Leu
      165              170              175

Tyr Thr Ile Arg Arg Gly Pro Ala Gly Ile Gly Arg Thr Gly Arg Gly
      180              185              190

Met Gly Gly Leu Phe Ser Val Gly Glu Thr Thr Ala Lys Val Leu Lys
      195              200              205

Asp Glu Ile Asp Val Lys Phe Lys Asp Val Ala Gly Cys Glu Glu Ala
      210              215              220

Lys Leu Glu Ile Met Glu Phe Val Asn Phe Leu Lys Asn Pro Lys Gln
      225              230              235              240

```

Tyr Gln Asp Leu Gly Ala Ile Ile Pro Lys Gly Ala Ile Leu Thr Gly
 245 250 255
 Pro Pro Gly Thr Gly Lys Thr Leu Leu Ala Lys Ala Thr Ala Gly Glu
 260 265 270
 Ala Asn Val Pro Phe Ile Thr Val Ser Gly Ser Glu Phe Leu Glu Met
 275 280 285
 Phe Val Gly Val Gly Pro Ala Arg Val Arg Asp Leu Phe Ala Leu Ala
 290 295 300
 Arg Lys Asn Ala Pro Cys Ile Leu Phe Ile Asp Glu Ile Asp Ala Val
 305 310 315 320
 Gly Arg Lys Arg Gly Arg Gly Asn Phe Gly Gly Gln Ser Glu Gln Glu
 325 330 335
 Asn Thr Leu Asn Gln Leu Leu Val Glu Met Asp Gly Phe Asn Thr Thr
 340 345 350
 Thr Asn Val Val Ile Leu Ala Gly Thr Asn Arg Pro Gly Pro Pro Asp
 355 360 365
 Ile Lys Gly Arg Ala Ser Ile Phe Lys Val His Leu Arg Pro Leu Lys
 370 375 380
 Leu Asp Ser Thr Leu Glu Lys Asp Lys Leu Ala Arg Lys Leu Ala Ser
 385 390 395 400
 Leu Thr Pro Gly Phe Ser Gly Ala Asp Val Ala Asn Val Cys Asn Glu
 405 410 415
 Ala Ala Leu Ile Ala Ala Arg His Leu Ser Asp Ser Ile Asn Gln Lys
 420 425 430
 His Phe Glu Gln Ala Ile Glu Arg Val Ile Gly Gly Leu Lys Lys Lys
 435 440 445
 Thr Gln Val Leu Gln Pro Glu Glu Lys Lys Thr Val Ala Tyr His Glu
 450 455 460
 Ala Gly His Ala Val Ala Gly Trp Tyr Leu Glu His Ala Asp Pro Leu
 465 470 475 480
 Leu Lys Val Ser Ile Ile Pro Arg Gly Lys Gly Leu Gly Tyr Ala Gln
 485 490 495
 Tyr Leu Pro Lys Glu Gln Tyr Leu Tyr Thr Lys Glu Gln Leu Leu Asp
 500 505 510
 Arg Met Cys Met Thr Leu Gly Gly Arg Val Ser Glu Glu Ile Phe Phe
 515 520 525
 Gly Arg Ile Thr Thr Gly Ala Gln Asp Asp Leu Arg Lys Val Thr Gln
 530 535 540
 Ser Ala Tyr Ala Gln Ile Val Gln Phe Gly Met Asn Glu Lys Val Gly
 545 550 555 560

Gln Ile Ser Phe Asp Leu Pro Arg Gln Gly Asp Met Val Leu Glu Lys
565 570 575

Pro Tyr Ser Glu Ala Thr Ala Arg Leu Ile Asp Asp Glu Val Arg Ile
580 585 590

Leu Ile Asn Asp Ala Tyr Lys Arg Thr Val Ala Leu Leu Thr Glu Lys
595 600 605

Lys Ala Asp Val Glu Lys Val Ala Leu Leu Leu Glu Lys Glu Val
610 615 620

Leu Asp Lys Asn Asp Met Val Glu Leu Leu Gly Pro Arg Pro Phe Ala
625 630 635 640

Glu Lys Ser Thr Tyr Glu Glu Phe Val Glu Gly Thr Gly Ser Leu Asp
645 650 655

Glu Asp Thr Ser Leu Pro Glu Gly Leu Lys Asp Trp Asn Lys Glu Arg
660 665 670

Glu Lys Glu Lys Glu Glu Pro Pro Gly Glu Lys Val Ala Asn
675 680 685

<210> 189

<211> 627

<212> DNA

<213> Homo sapiens

<400> 189

gattggagtt cttttcttaa atatccaaat aagattatag ctatgttttt accaaatggt 60
tgtgcggttaa atagagagtg gatggaaatt aaccctagaa aggatagttg taacttttaa 120
aaagttgatt aactatttcg tgtgctaatt tgagtttttc tgaatactcc aatatgggtt 180
cctttaacac ctgctctcag tttacaatca cctaacttcc cagcgttggt gtctttttct 240
ctgtctgacc ctgtcttatt tctctacaa agacatatcc tgcgctgtac ttcagatact 300
tttttcgagg aacatttggtg atttgtggca taaagtaact gtctaaagga aatcttctga 360
gaggatcttg tcattttatg aaaggggcaa ttaaggggaa atggaagcag atctttttaa 420
gaaggagcat ttgaaattag cccaggaatc atgtccggcg agtcctgctc ttttgtacct 480
gggcataata gtcagccaca cagagctaga gttagttcaa gaattgtctt tcttgatcgt 540
gctatatttt tggaaacacg ttagatacag aggtaagatg tcaaaattct gaaatacaca 600
caatatagga tcaaaaaaaaa aaaaaaa 627

<210> 190

<211> 63

<212> PRT

<213> Homo sapiens

<400> 190

Met Glu Ala Asp Leu Leu Lys Lys Glu His Leu Lys Leu Ala Gln Glu
1 5 10 15

Ser Cys Pro Ala Ser Pro Ala Leu Leu Tyr Leu Gly Ile Ile Val Ser
20 25 30

His Thr Glu Leu Glu Leu Val Gln Glu Leu Ser Phe Leu Ile Val Leu
35 40 45

Tyr Phe Trp Lys His Val Arg Tyr Arg Gly Lys Met Ser Lys Phe

50

55

60

<210> 191
 <211> 868
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (733)

<400> 191
 attctgtccc tgcccttaga aaacctaaaa taccaagggt gtcattgctgg caactccctg 60
 cccagtcctg cacaagcct tggctgtgtg tggcaccctc tgcctcctac cccagagcag 120
 ctggctccat tggcttctcc ctgcaccagc cctgtcctca ggggtcagga aaaagcacac 180
 agctttcttt cctctcctcc agaggcctgg aaggagggtg gaggtccagt aagggcctgg 240
 ctgccttggg tttcttgggc ctgccttgcc aactgcaccc tgtagctcct gctccctgtg 300
 accccagAAC agagggtgtg ccttccctgt ctcttagaca aagcacaag ggatgccctg 360
 cttggcttga gcctgcccaa ctgaaggatt ttctctgccc caggacctt ccatccctga 420
 atacaaggct ctaggcaact tctctctggg tggtagacac tagaatgcct ggcattagcc 480
 ctgaaagga ggttgggggtg tatgggtagt gagctagggt gggagaaagg tgggtgtgaa 540
 aggacagatg ctagtgttag ttctactcac tcattcattc attagtcaa cagtactgag 600
 caccacctgc actagaggca gaggggtgaa caagataccc ttttgcctgg ggggacgtcc 660
 acttcccatg ggttgggcta tttccaggaa agccctcag tcctcctccc tgttctggct 720
 gtgtgtgaag gangtgtgtg agcaggccca atcctttgca gcaagaatga gaggtcagag 780
 tattccattg cacacgcacc ctggggctga cagacttgtg ccccttagcc ttcattgcatg 840
 cccaagcact ggcagctttg cagccctc 868

<210> 192
 <211> 107
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (62)

<400> 192
 Met Leu Val Val Val Ser Leu Thr His Ser Phe Ile Ser Ala Thr Val
 1 5 10 15
 Leu Ser Thr Thr Cys Thr Arg Gly Arg Gly Val Asn Lys Ile Pro Phe
 20 25 30
 Cys Leu Gly Gly Arg Pro Leu Pro Met Gly Leu Ala Ile Ser Arg Lys
 35 40 45
 Ala Pro Gln Ser Ser Ser Leu Phe Trp Leu Cys Val Lys Xaa Val Cys
 50 55 60
 Glu Gln Ala Gln Ser Phe Ala Ala Arg Met Arg Gly Gln Ser Ile Pro
 65 70 75 80
 Leu His Thr His Pro Gly Ala Asp Arg Leu Val Pro Pro Ser Leu His
 85 90 95
 Ala Cys Pro Ser Thr Gly Ser Phe Ala Ala Pro
 100 105

<210> 193
 <211> 467
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (57)

<220>
 <221> unsure
 <222> (254)

<400> 193
 tggctatccc tagaagcact tcatccatct taaccacccc aaacgggata ccttcanatc 60
 tcatacccag taagatgcaa gaaaggaata tttgagagca agcagccctg ttccaggggc 120
 cccaggtatg tgtagaggcc cagtgggggt ggccacttgg tgtttctacc accccctgcc 180
 atccagtctg gccaggtacc tacctgggag gttggtgtac ttggcttaag tacttcatgc 240
 tttattcagg ctgnttcccc acagcaccgg caggaaatga aggtgcactt atatgcatcc 300
 ctgcaggaat aaagagtggg tggcctgccc agcccagcac cacagccttt ccccagccag 360
 gagagaccac ctaaggatca aggcagctcc tgttttcttg gttctgtgac actcagagtct 420
 gagccagccc ctcaggaatt gcctcaaaag agaaaaaaaa aaaaaaaa 467

<210> 194
 <211> 1035
 <212> DNA
 <213> Homo sapiens

<400> 194
 aatctttttg ttgtcaagct tgagggtgtg caggcttgag atctggccat acacttgagt 60
 gacaatgaca tccactttgc ctttctctcc acaggtgtcc actcccaggt ccaactgcag 120
 acttcgaatt cggccttcat ggcctagatg attgcaagtc aatggaagga gctgcagagg 180
 caaatcaaac gccagcacag ctggattctc agggctcttg ataccatcaa agccgagata 240
 ctggctactg atgtgtctgt ggaggatgag gaagggactg gaagccccc aaagctgaggtt 300
 caactatgct acctggaagc acaaagagat gctgttgagc agatgtccct caagctgtrc 360
 agcgagcagt ataccagcag cagcaagcga aaggaagagt ttgctgatat gtcaaaagtt 420
 tcattcagtg ggaagcaatg ggcttctgga ctttgattca gaatatcmgg agctctggga 480
 ttggctgatt gacatggagt cccttctgat ggacagccac gacctgatga tgtcagagga 540
 gcagcagcag catctttaca agcgatacag tgtggaaatg tccatcagac acctgaaaaa 600
 gacggagctg cttagtaagg ttgaagcttt gaagaaaggt ggcgttttac taccaaatga 660
 tctccttgaa aaagtggatt caattaatga aaaatgggaa ctrcttgggg tatttgcat 720
 tttattactg tttgtagggt atgtgtacat tttttgcgta gtgaagtact ctgtccgatt 780
 tctaatttga ggcacaaata tctctctctt tcaattcact acctacgttt caaacaagct 840
 attcatgcta ttatgggaaa gacactgctt ttctctctct gttgattttt ttttttctctg 900
 agcttgctcc ctctcagatt ttaataatct tggttcttta atacatgaaa aagtaagtaa 960
 aatatgccat gtattatggg tatgcaccaa gtcaactata atacagtata tctgatata 1020
 aaaaaaaaaa aaaaaa 1035

<210> 195
 <211> 179
 <212> PRT
 <213> Homo sapiens

<220>
 <221> UNSURE
 <222> (37)

<220>
 <221> UNSURE

<222> (73)

<400> 195

Met Cys Leu Trp Arg Met Arg Lys Gly Leu Glu Ala Pro Arg Leu Arg
 1 5 10 15

Phe Asn Tyr Ala Thr Trp Lys His Lys Glu Met Leu Leu Ser Arg Cys
 20 25 30

Pro Ser Ser Cys Xaa Ala Ser Ser Ile Pro Ala Ala Ala Ser Glu Arg
 35 40 45

Lys Ser Leu Leu Ile Cys Gln Lys Phe His Ser Val Gly Ser Asn Gly
 50 55 60

Leu Leu Asp Phe Asp Ser Glu Tyr Xaa Glu Leu Trp Asp Trp Leu Ile
 65 70 75 80

Asp Met Glu Ser Leu Val Met Asp Ser His Asp Leu Met Met Ser Glu
 85 90 95

Glu Gln Gln Gln His Leu Tyr Lys Arg Tyr Ser Val Glu Met Ser Ile
 100 105 110

Arg His Leu Lys Lys Thr Glu Leu Leu Ser Lys Val Glu Ala Leu Lys
 115 120 125

Lys Gly Gly Val Leu Leu Pro Asn Asp Leu Leu Glu Lys Val Asp Ser
 130 135 140

Ile Asn Glu Lys Trp Glu Leu Leu Gly Val Phe Ala Phe Leu Leu Leu
 145 150 155 160

Phe Val Gly Tyr Val Tyr Ile Phe Cys Val Val Lys Tyr Ser Val Arg
 165 170 175

Phe Leu Ile

<210> 196

<211> 3831

<212> DNA

<213> Homo sapiens

<400> 196

gcagcaggtc aaacaaggca tctcctagta ttgcataccta cagatgtgct gtaaacaatca 60
 aaagaagacg gtgggatcag gagatgctgt tttggaaaga agtgagggtt agacttctcc 120
 atgttaacca tgagcgtgac actttccccc ctgagggtcac aggacctgga tcccatggct 180
 actgatgctt caccatggc catcaacatg acaccactg tggagcaggg tgagggagaa 240
 gaggcaatga aggacatgga ctctgaccag cagtatgaaa agccaccccc actacacaca 300
 ggggctgact ggaagattgt cctccactta cctgaaattg agacctggct ccggatgacc 360
 tcagagaggg tccgagacct aacctattca gtccagcagg attcggacag caagcatgtg 420
 gatgtacatc tagttcaact aaaggacatt tgtgaagata tttctgatca tgttgagcaa 480
 atccatgccc tccttgaaac agagtctctc ctaaagctgc tgtcttactc tgtcaacgtg 540
 atagtggaca tccacgcagt gcagctcttc tggcaccagc ttcgagtctc agtgctgggt 600
 ctgcgggagc gcattctgca aggtctgcag gacgccaatg gcaactacac taggcagacg 660
 gacattctgc aagctttctc tgaagagaca aaagagggcc ggcttgattc tctaacagaa 720
 gtggatgact caggacaatt aaccatcaaa tgttctcaaa attacttgct tctggattgt 780
 ggcattactg cattcgaact gcttgactac agtccaagtg aggatttgct cagtgggcta 840

```

gggtgacatga cctctagcca agtcaaaacc aaaccctttg actcttggag ctacagttag 900
atggaaaagg agtttctcga gcttatccga agtggttggtt tacttacggg agctgctgac 960
tctatctcta ccaatggcag tgaagcagtt actgaggagg tatctcaagt atctctctca 1020
gtagacgaca aaggtggatg tgaggaagac aatgcttctg cagtccaaga gcaaccaggc 1080
ttaacactgg ggggtgcacg atcttcagga gaagctctga caaatgctgc tcaaccctcc 1140
tctgagactg tgcagcaaga atccagtctc tcctcccatc atgatgcaaa gaatcagcag 1200
cctgttccct gtgaaaatgc aacccccaaa cgaaccatca gagattgctt taattataac 1260
gaggactctc ccacacagcc tacattgcca aaaagaggac tttttcttaa agaggaaact 1320
tttaagaatg atctgaaagg caatggtgga aagaggcaaa tgggtgatct aaagcctgag 1380
atgagcagaa gcaccctctc gctagtagat cctcctgaca gatccaaact ttgcctggta 1440
ttgcagtctt ctaccccaa cagccctctc gctgccagcc agtcttatga gtgtttacac 1500
aagggtggga atgggaacct tgaaaacaca gtcaaatttc acattaaaga aatttcttcc 1560
agcctgggaa ggcttaacga ctgctataaa gagaaatctc gacttaaaaa gccacacaag 1620
acctcagaag aggtgcctcc atgccgaaca cctaaacggg ggactgggtc agggcaacaa 1680
gctaaaaata caaagagctc agcagtgcc aatggagagc tttcttatac ttccaaggcc 1740
atagaggggc cacaacaaaa ttctgcttcc acatcctcac ttgagccttg taatcagaga 1800
agttggaatg ccaaattgca attgcagtca gaaacatcca gttcaccagc ttttactcag 1860
agcagtgaat cctctgttgg ctacagacaac atcatgtctc cgggtgccact tctttcaaaa 1920
cacaaaagca aaaaagggtc agcctcctct ccaagtccag tcactaggaa tgggtagggt 1980
gtggaggcct ggtatggctc tgatgaatac ctgactgctc cctctcacct taagcagaca 2040
gaagtattgg ctttgaagtt ggaaaacctc acaaagcttc tgcttcagaa acccagagga 2100
gaaaccatcc agaattattg tgaactggaa ctgtctgaaa tgaattcaga ttctgaaatc 2160
tatccaacct atcatgtcaa aaagaagcat acaaggctag gcagggtgtc tccaagctca 2220
tctagtgaac tagctctctc actaggggag agcattgaat ctgggcccct gagtgcatt 2280
ctttctgatg aggaagtcag tatgcctctc gctggcatga aaaagtatgc tgatgagaag 2340
tcagaaagag cttcatcctc tgagaaaaat gagagccatt ctgccactaa atcagcttta 2400
attcagaaac tgatgcaaga tattcagcac caagacaact atgaagccat atgggaaaaa 2460
atagaggggt ttgtaaacaa actggatgaa ttcattcaat gggttaaatga agccatggaa 2520
actacagaga attggactcc ccctaaagca gagatggatg accttaaaact gtatctggag 2580
acacacttga gttttaagtt gaatgtagac agtcattgtg ctctcaagga agctgtggag 2640
gagggaaggac accaactctc tgagcttatt gcatctcaca aagcaggact gaaggacatg 2700
ctgcggatga ttgcaagtca atggaaggag ctgcagaggc aaatcaaacg gcagcacagc 2760
tggtattctc gggctctgga taccatcaaa gccgagatac tggctactga tgtgtctgtg 2820
gaggatgagg aagggactgg aagccccaag gctgagggtc aactatgcta cctggaagca 2880
caaagagatg ctgttgagca gatgtccctc aagctgtaca gcgagcagta taccagcagc 2940
agcaagcgaa aggaagagtt tgctgatatg tcaaaagttc attcagtggg aagcaatggg 3000
cttctggact ttgattcaga atatcaggag ctctgggatt ggctgattga catggagtcc 3060
cttgtgatgg acagccacga cctgatgatg tcagaggagc agcagcagca tctttacaag 3120
cgatacagtg tggaatgtc catcagacac ctgaaaaaga cggagctgct tagtaagggt 3180
gaagctttga agaaagggtg cgttttacta ccaaatgatc tccttgaaaa agtggtattca 3240
attaatgaaa aatgggaact gcttggggta tttgcatttt tattactggt tgtaggttat 3300
gtgtacattt tttgcgtagt gaagtactct gtccgatttc taatttgagg cacaatatc 3360
tctctctttc aattcactac ctacgtttca aacaagctat tcatgctatt atgggaaaga 3420
cactgctttt cctcttctgt tgattttttt ttttctgag cttgtccctc ctcatattt 3480
aataattttg gttctttaat acatgaaaaa gtaagtaaaa tatgccatgt attatgggta 3540
tgaccaagt caactataat acagtatatc tgatatatac tgactgtcat gcttgaatga 3600
atgttaatag aatttattct gaaggtacat gtgagagaca tctactgttt aactatttac 3660
tgtaccctta agatgaaaag tggagttgtc actacagctt tcaagtcaca ctaaagccac 3720
caaaacaaa agtgcaattt gacccaaate tgaattgcag aattgaatca gcctgtgttt 3780
tgtgcctcaa tttccagctc acttttaaca aaagccaaaa aaaaaaaaaa a 3831

```

<210> 197

<211> 1075

<212> PRT

<213> Homo sapiens

<400> 197

```

Met Leu Thr Met Ser Val Thr Leu Ser Pro Leu Arg Ser Gln Asp Leu
  1             5             10             15

```

Asp Pro Met Ala Thr Asp Ala Ser Pro Met Ala Ile Asn Met Thr Pro
 20 25 30
 Thr Val Glu Gln Gly Glu Gly Glu Glu Ala Met Lys Asp Met Asp Ser
 35 40 45
 Asp Gln Gln Tyr Glu Lys Pro Pro Leu His Thr Gly Ala Asp Trp
 50 55 60
 Lys Ile Val Leu His Leu Pro Glu Ile Glu Thr Trp Leu Arg Met Thr
 65 70 75 80
 Ser Glu Arg Val Arg Asp Leu Thr Tyr Ser Val Gln Gln Asp Ser Asp
 85 90 95
 Ser Lys His Val Asp Val His Leu Val Gln Leu Lys Asp Ile Cys Glu
 100 105 110
 Asp Ile Ser Asp His Val Glu Gln Ile His Ala Leu Leu Glu Thr Glu
 115 120 125
 Phe Ser Leu Lys Leu Leu Ser Tyr Ser Val Asn Val Ile Val Asp Ile
 130 135 140
 His Ala Val Gln Leu Leu Trp His Gln Leu Arg Val Ser Val Leu Val
 145 150 155 160
 Leu Arg Glu Arg Ile Leu Gln Gly Leu Gln Asp Ala Asn Gly Asn Tyr
 165 170 175
 Thr Arg Gln Thr Asp Ile Leu Gln Ala Phe Ser Glu Glu Thr Lys Glu
 180 185 190
 Gly Arg Leu Asp Ser Leu Thr Glu Val Asp Asp Ser Gly Gln Leu Thr
 195 200 205
 Ile Lys Cys Ser Gln Asn Tyr Leu Ser Leu Asp Cys Gly Ile Thr Ala
 210 215 220
 Phe Glu Leu Ser Asp Tyr Ser Pro Ser Glu Asp Leu Leu Ser Gly Leu
 225 230 235 240
 Gly Asp Met Thr Ser Ser Gln Val Lys Thr Lys Pro Phe Asp Ser Trp
 245 250 255
 Ser Tyr Ser Glu Met Glu Lys Glu Phe Pro Glu Leu Ile Arg Ser Val
 260 265 270
 Gly Leu Leu Thr Val Ala Ala Asp Ser Ile Ser Thr Asn Gly Ser Glu
 275 280 285
 Ala Val Thr Glu Glu Val Ser Gln Val Ser Leu Ser Val Asp Asp Lys
 290 295 300
 Gly Gly Cys Glu Glu Asp Asn Ala Ser Ala Val Glu Glu Gln Pro Gly
 305 310 315 320
 Leu Thr Leu Gly Val Ser Ser Ser Ser Gly Glu Ala Leu Thr Asn Ala
 325 330 335

Ala Gln Pro Ser Ser Glu Thr Val Gln Gln Glu Ser Ser Ser Ser Ser
 340 345 350
 His His Asp Ala Lys Asn Gln Gln Pro Val Pro Cys Glu Asn Ala Thr
 355 360 365
 Pro Lys Arg Thr Ile Arg Asp Cys Phe Asn Tyr Asn Glu Asp Ser Pro
 370 375 380
 Thr Gln Pro Thr Leu Pro Lys Arg Gly Leu Phe Leu Lys Glu Glu Thr
 385 390 395 400
 Phe Lys Asn Asp Leu Lys Gly Asn Gly Gly Lys Arg Gln Met Val Asp
 405 410 415
 Leu Lys Pro Glu Met Ser Arg Ser Thr Pro Ser Leu Val Asp Pro Pro
 420 425 430
 Asp Arg Ser Lys Leu Cys Leu Val Leu Gln Ser Ser Tyr Pro Asn Ser
 435 440 445
 Pro Ser Ala Ala Ser Gln Ser Tyr Glu Cys Leu His Lys Val Gly Asn
 450 455 460
 Gly Asn Leu Glu Asn Thr Val Lys Phe His Ile Lys Glu Ile Ser Ser
 465 470 475 480
 Ser Leu Gly Arg Leu Asn Asp Cys Tyr Lys Glu Lys Ser Arg Leu Lys
 485 490 495
 Lys Pro His Lys Thr Ser Glu Glu Val Pro Pro Cys Arg Thr Pro Lys
 500 505 510
 Arg Gly Thr Gly Ser Gly Lys Gln Ala Lys Asn Thr Lys Ser Ser Ala
 515 520 525
 Val Pro Asn Gly Glu Leu Ser Tyr Thr Ser Lys Ala Ile Glu Gly Pro
 530 535 540
 Gln Thr Asn Ser Ala Ser Thr Ser Ser Leu Glu Pro Cys Asn Gln Arg
 545 550 555 560
 Ser Trp Asn Ala Lys Leu Gln Leu Gln Ser Glu Thr Ser Ser Ser Pro
 565 570 575
 Ala Phe Thr Gln Ser Ser Glu Ser Ser Val Gly Ser Asp Asn Ile Met
 580 585 590
 Ser Pro Val Pro Leu Leu Ser Lys His Lys Ser Lys Lys Gly Gln Ala
 595 600 605
 Ser Ser Pro Ser His Val Thr Arg Asn Gly Glu Val Val Glu Ala Trp
 610 615 620
 Tyr Gly Ser Asp Glu Tyr Leu Ala Leu Pro Ser His Leu Lys Gln Thr
 625 630 635 640
 Glu Val Leu Ala Leu Lys Leu Glu Asn Leu Thr Lys Leu Leu Pro Gln
 645 650 655

Lys Pro Arg Gly Glu Thr Ile Gln Asn Ile Asp Asp Trp Glu Leu Ser
 660 665 670
 Glu Met Asn Ser Asp Ser Glu Ile Tyr Pro Thr Tyr His Val Lys Lys
 675 680 685
 Lys His Thr Arg Leu Gly Arg Val Ser Pro Ser Ser Ser Ser Asp Ile
 690 695 700
 Ala Ser Ser Leu Gly Glu Ser Ile Glu Ser Gly Pro Leu Ser Asp Ile
 705 710 715 720
 Leu Ser Asp Glu Glu Ser Ser Met Pro Leu Ala Gly Met Lys Lys Tyr
 725 730 735
 Ala Asp Glu Lys Ser Glu Arg Ala Ser Ser Ser Glu Lys Asn Glu Ser
 740 745 750
 His Ser Ala Thr Lys Ser Ala Leu Ile Gln Lys Leu Met Gln Asp Ile
 755 760 765
 Gln His Gln Asp Asn Tyr Glu Ala Ile Trp Glu Lys Ile Glu Gly Phe
 770 775 780
 Val Asn Lys Leu Asp Glu Phe Ile Gln Trp Leu Asn Glu Ala Met Glu
 785 790 795 800
 Thr Thr Glu Asn Trp Thr Pro Pro Lys Ala Glu Met Asp Asp Leu Lys
 805 810 815
 Leu Tyr Leu Glu Thr His Leu Ser Phe Lys Leu Asn Val Asp Ser His
 820 825 830
 Cys Ala Leu Lys Glu Ala Val Glu Glu Glu Gly His Gln Leu Leu Glu
 835 840 845
 Leu Ile Ala Ser His Lys Ala Gly Leu Lys Asp Met Leu Arg Met Ile
 850 855 860
 Ala Ser Gln Trp Lys Glu Leu Gln Arg Gln Ile Lys Arg Gln His Ser
 865 870 875 880
 Trp Ile Leu Arg Ala Leu Asp Thr Ile Lys Ala Glu Ile Leu Ala Thr
 885 890 895
 Asp Val Ser Val Glu Asp Glu Glu Gly Thr Gly Ser Pro Lys Ala Glu
 900 905 910
 Val Gln Leu Cys Tyr Leu Glu Ala Gln Arg Asp Ala Val Glu Gln Met
 915 920 925
 Ser Leu Lys Leu Tyr Ser Glu Gln Tyr Thr Ser Ser Ser Lys Arg Lys
 930 935 940
 Glu Glu Phe Ala Asp Met Ser Lys Val His Ser Val Gly Ser Asn Gly
 945 950 955 960
 Leu Leu Asp Phe Asp Ser Glu Tyr Gln Glu Leu Trp Asp Trp Leu Ile
 965 970 975

Asp Met Glu Ser Leu Val Met Asp Ser His Asp Leu Met Met Ser Glu
 980 985 990
 Glu Gln Gln Gln His Leu Tyr Lys Arg Tyr Ser Val Glu Met Ser Ile
 995 1000 1005
 Arg His Leu Lys Lys Thr Glu Leu Leu Ser Lys Val Glu Ala Leu Lys
 1010 1015 1020
 Lys Gly Gly Val Leu Leu Pro Asn Asp Leu Leu Glu Lys Val Asp Ser
 1025 1030 1035 1040
 Ile Asn Glu Lys Trp Glu Leu Leu Gly Val Phe Ala Phe Leu Leu Leu
 1045 1050 1055
 Phe Val Gly Tyr Val Tyr Ile Phe Cys Val Val Lys Tyr Ser Val Arg
 1060 1065 1070
 Phe Leu Ile
 1075

<210> 198
 <211> 3113
 <212> DNA
 <213> Homo sapiens

<400> 198
 gcaaacacat tgaagtacag gtagcccagg aaactagaaa tgtatctact ggctctgctg 60
 aaaaatgaaga aaagtcagaa gttcaagcaa tcatcgaatc tactcctgag ctggatatgg 120
 acaaagatct cagtggatat aaaggttcaa gcactccac caaaggcata gagaacaaag 180
 cttttgatcg caatacagaa tctctctttg aagaactgtc ttcagctggc tcaggcctaa 240
 taggagatgt ggatgaagga gcagatttac taggaatggg tcgggaagtt gagaatctta 300
 tattagaaaa tacacaactg ttggaaacca aaaatgcttt gaacatagtg aagaatgatt 360
 tgatagcaaa agtggatgaa ctgacctgtg agaaagatgt gctgcaaggg gaattggagg 420
 ctgtgaagca agccaaactg aaactagagg aaaagaacag agaattggag gaagagctta 480
 ggaaagctcg ggcagaagct gaagatgcag ggcaaaaagc aaaagatgac gatgatagtg 540
 atattccac agcccagagg aaacgggtta ctagagtaga aatggcccgt gttctcatgg 600
 agcgaaccca gtataaagag agattgatgg agcttcagga agctgttcga tggacagaga 660
 tgattcgggc atcacagaaa aatccagcca tgcaggaaaa aaaaaggcca agcatttggc 720
 agtttttcag ccgacttttc agctcctcaa gtaacacgac taagaagcct gaaccacctg 780
 ttaatctgaa gtacaatgca cccacgtctc atgttactcc gtccgtcaag aaaagaagca 840
 gcaccttatc tcagctccct ggggataagt ccaaagcctt tgatttcctt agtgaagaaa 900
 ctgaagctag tttagcctca cgcagagaaac aaaagagaga gcagtatcgt caggtaaaaag 960
 cacatgttca gaaggaagac ggtagagtgc aggccttttg ctggagtctg cctcagaagt 1020
 acaaacaggt aaccaatggg caaggtgaaa ataagatgaa aaatttacct gtgcctgtct 1080
 atctcagacc tctggatgaa aaagatacat caatgaagct gtggtgtgct gttggagtca 1140
 atttatctgg tgggaagacc agagatggg gttctgttgt tggagcaagt gtattttaca 1200
 aggatgttgc tggtttggat acagaaggca gtaaacagcg aagtgcctct cagagtagtt 1260
 tagataagtt agatcaggaa cttaaggaaac agcagaagga gttaaaaaat caagaagaat 1320
 tatccagtct agtttggatc tgtaccagca ctcattcggc tacaaaagtt cttattattg 1380
 atgctgttca acctggcaac atcctagaca gtttactgt ttgcaactct catgttctgt 1440
 gcattgcaag tgtgccaggt gcacgagaaa cagactaccc tgcaggagaa gatctttcag 1500
 aatctggtca ggtagacaaa gcatctttat gtggaagtat gacaagcaac agctcagcag 1560
 agacagacag cctgttagga ggcacacag tggttggttg ttctgcagaa ggtgtgacgg 1620
 gagctgccac ttcccctagt acaaatgggtg cttctccagt gatggataaa ccaccagaaa 1680
 tggaaagcaga aaatagttag gttgatgaaa atgttccaac agcagaagaa gcaactgaag 1740
 ctacagaagg gaatgcgggg tcagctgaag acacagtgga catctcccaa actggcgtct 1800
 acacagagca tgtctttaca gatcctttgg gagttcagat ccagagaagac ctctcccag 1860
 tgtatcagtc gagcaatgac tcagatgcat ataaagatca aatatcagta ctgccaaatg 1920


```

aacaagactt ggtgagagaa gaagcccaga aaatgagtag tcttttacc aactatgtggc 1980
ttggagctca aaatggctgt ttgtatgtcc attcatctgt agcccagtgg aggaaatgtc 2040
tccattccat taaacttaaa gattcgattc tcagtattgt acacgtgaag ggaatcgtgt 2100
tagtagccct ggctgacggc acccttgcaa tctttcacag aggagtggat gggcagtggg 2160
atttgtcaaa ctatcacctc ttagaccttg gacggcctca tcattccatc cgttgcatga 2220
ctgtggtaca tgacaaagtc tgggtgtggt ataggaacaa aatctatgtg gtgcagccaa 2280
aggccatgaa aatagagaaa tctttttagt cacatcccag gaaggagagc caagtgcgac 2340
agcttgctg ggtgggggat ggcgtgtggg tctccattcg cttggattct acgctccgtc 2400
tctatcatgc acacacttat caacatctac aggatgtgga cattgagcct tatgtaagca 2460
aaatgttagg tactggaaaa ctgggcttct cttttgtgag aattacagct cttatggtgt 2520
cttgtaatcg tttgtgggtg gggacaggaa atggtgtcat tatctccatc ccattgacag 2580
aaagtaagta tttttttaga taactgccat ggaacaaata ggagaattat aggaaatagt 2640
tatgtctgaa attcaaactc gtgttagctc aagaaaatca gtgatggcaa agatctagct 2700
atagttaact tcaactcaat gaaacttata aattctagga aaataagttt tcatgtatca 2760
agaccaatac tcataattaa aagtttttct ttgtcaaaac attttcacag ttcaggaatc 2820
acagtcctat ttctctttt gaaataaaaag ttaaagtcca tttttcttaa gccttattat 2880
accattatcc aaaaacacta cctagcattc atcttagttt ttcatacttc cttagcagat 2940
agtttagtat attccatttg tcttatttga ggtttaaaaa aagttgtgga agaagtttct 3000
caggagtcca gcaagccttt ccattctatc tctattttta aagaatgtga tttattgttt 3060
ttatgaataa aatatacctc gtgtgtttct caaaaaaaa aaaaaaaaaa aaa 3113

```

<210> 199

<211> 828

<212> PRT

<213> Homo sapiens

<400> 199

```

Met Asp Lys Asp Leu Ser Gly Tyr Lys Gly Ser Ser Thr Pro Thr Lys
  1             5             10             15

Gly Ile Glu Asn Lys Ala Phe Asp Arg Asn Thr Glu Ser Leu Phe Glu
      20             25             30

Glu Leu Ser Ser Ala Gly Ser Gly Leu Ile Gly Asp Val Asp Glu Gly
      35             40             45

Ala Asp Leu Leu Gly Met Gly Arg Glu Val Glu Asn Leu Ile Leu Glu
      50             55             60

Asn Thr Gln Leu Leu Glu Thr Lys Asn Ala Leu Asn Ile Val Lys Asn
      65             70             75             80

Asp Leu Ile Ala Lys Val Asp Glu Leu Thr Cys Glu Lys Asp Val Leu
      85             90             95

Gln Gly Glu Leu Glu Ala Val Lys Gln Ala Lys Leu Lys Leu Glu Glu
      100            105            110

Lys Asn Arg Glu Leu Glu Glu Glu Leu Arg Lys Ala Arg Ala Glu Ala
      115            120            125

Glu Asp Ala Gly Gln Lys Ala Lys Asp Asp Asp Asp Ser Asp Ile Pro
      130            135            140

Thr Ala Gln Arg Lys Arg Phe Thr Arg Val Glu Met Ala Arg Val Leu
      145            150            155            160

Met Glu Arg Asn Gln Tyr Lys Glu Arg Leu Met Glu Leu Gln Glu Ala
      165            170            175

```

Val Arg Trp Thr Glu Met Ile Arg Ala Ser Arg Glu Asn Pro Ala Met
 180 185 190
 Gln Glu Lys Lys Arg Ser Ser Ile Trp Gln Phe Phe Ser Arg Leu Phe
 195 200 205
 Ser Ser Ser Ser Asn Thr Thr Lys Lys Pro Glu Pro Pro Val Asn Leu
 210 215 220
 Lys Tyr Asn Ala Pro Thr Ser His Val Thr Pro Ser Val Lys Lys Arg
 225 230 235 240
 Ser Ser Thr Leu Ser Gln Leu Pro Gly Asp Lys Ser Lys Ala Phe Asp
 245 250 255
 Phe Leu Ser Glu Glu Thr Glu Ala Ser Leu Ala Ser Arg Arg Glu Gln
 260 265 270
 Lys Arg Glu Gln Tyr Arg Gln Val Lys Ala His Val Gln Lys Glu Asp
 275 280 285
 Gly Arg Val Gln Ala Phe Gly Trp Ser Leu Pro Gln Lys Tyr Lys Gln
 290 295 300
 Val Thr Asn Gly Gln Gly Glu Asn Lys Met Lys Asn Leu Pro Val Pro
 305 310 315 320
 Val Tyr Leu Arg Pro Leu Asp Glu Lys Asp Thr Ser Met Lys Leu Trp
 325 330 335
 Cys Ala Val Gly Val Asn Leu Ser Gly Gly Lys Thr Arg Asp Gly Gly
 340 345 350
 Ser Val Val Gly Ala Ser Val Phe Tyr Lys Asp Val Ala Gly Leu Asp
 355 360 365
 Thr Glu Gly Ser Lys Gln Arg Ser Ala Ser Gln Ser Ser Leu Asp Lys
 370 375 380
 Leu Asp Gln Glu Leu Lys Glu Gln Gln Lys Glu Leu Lys Asn Gln Glu
 385 390 395 400
 Glu Leu Ser Ser Leu Val Trp Ile Cys Thr Ser Thr His Ser Ala Thr
 405 410 415
 Lys Val Leu Ile Ile Asp Ala Val Gln Pro Gly Asn Ile Leu Asp Ser
 420 425 430
 Phe Thr Val Cys Asn Ser His Val Leu Cys Ile Ala Ser Val Pro Gly
 435 440 445
 Ala Arg Glu Thr Asp Tyr Pro Ala Gly Glu Asp Leu Ser Glu Ser Gly
 450 455 460
 Gln Val Asp Lys Ala Ser Leu Cys Gly Ser Met Thr Ser Asn Ser Ser
 465 470 475 480
 Ala Glu Thr Asp Ser Leu Leu Gly Gly Ile Thr Val Val Gly Cys Ser
 485 490 495

Ala Glu Gly Val Thr Gly Ala Ala Thr Ser Pro Ser Thr Asn Gly Ala
 500 505 510
 Ser Pro Val Met Asp Lys Pro Pro Glu Met Glu Ala Glu Asn Ser Glu
 515 520 525
 Val Asp Glu Asn Val Pro Thr Ala Glu Glu Ala Thr Glu Ala Thr Glu
 530 535 540
 Gly Asn Ala Gly Ser Ala Glu Asp Thr Val Asp Ile Ser Gln Thr Gly
 545 550 555 560
 Val Tyr Thr Glu His Val Phe Thr Asp Pro Leu Gly Val Gln Ile Pro
 565 570 575
 Glu Asp Leu Ser Pro Val Tyr Gln Ser Ser Asn Asp Ser Asp Ala Tyr
 580 585 590
 Lys Asp Gln Ile Ser Val Leu Pro Asn Glu Gln Asp Leu Val Arg Glu
 595 600 605
 Glu Ala Gln Lys Met Ser Ser Leu Leu Pro Thr Met Trp Leu Gly Ala
 610 615 620
 Gln Asn Gly Cys Leu Tyr Val His Ser Ser Val Ala Gln Trp Arg Lys
 625 630 635 640
 Cys Leu His Ser Ile Lys Leu Lys Asp Ser Ile Leu Ser Ile Val His
 645 650 655
 Val Lys Gly Ile Val Leu Val Ala Leu Ala Asp Gly Thr Leu Ala Ile
 660 665 670
 Phe His Arg Gly Val Asp Gly Gln Trp Asp Leu Ser Asn Tyr His Leu
 675 680 685
 Leu Asp Leu Gly Arg Pro His His Ser Ile Arg Cys Met Thr Val Val
 690 695 700
 His Asp Lys Val Trp Cys Gly Tyr Arg Asn Lys Ile Tyr Val Val Gln
 705 710 715 720
 Pro Lys Ala Met Lys Ile Glu Lys Ser Phe Asp Ala His Pro Arg Lys
 725 730 735
 Glu Ser Gln Val Arg Gln Leu Ala Trp Val Gly Asp Gly Val Trp Val
 740 745 750
 Ser Ile Arg Leu Asp Ser Thr Leu Arg Leu Tyr His Ala His Thr Tyr
 755 760 765
 Gln His Leu Gln Asp Val Asp Ile Glu Pro Tyr Val Ser Lys Met Leu
 770 775 780
 Gly Thr Gly Lys Leu Gly Phe Ser Phe Val Arg Ile Thr Ala Leu Met
 785 790 795 800
 Val Ser Cys Asn Arg Leu Trp Val Gly Thr Gly Asn Gly Val Ile Ile
 805 810 815

Ser Ile Pro Leu Thr Glu Ser Lys Tyr Ile Phe Arg
820 825

<210> 200
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 200
tnaggtttgc acttcttctt gcaaattcc 29

<210> 201
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 201
cnaatatctg agtctgtcct gcgggactc 29

<210> 202
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 202
cnggctctgt gatggctgaa tggaaagaa 29

<210> 203
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 203
angagtccaa ttaaagaacg aaaagcggg 29

<210> 204
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 204
cntcagcttg ggcaagagta aagagggca 29

<210> 205
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 205
tnatttaaga tgctgtgtaa gcctctcgc 29

<210> 206
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 206
tntctgtgtg cacagagctt tggagtggc 29

<210> 207
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 207
gnttggattc tctggcagag tacacctgc 29

<210> 208
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 208
gnaggcatat cttttacagg ttcagctcc 29

<210> 209
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 209
cncagtgaga caggtagtag gagccagat 29

<210> 210
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 210
angtaaattc cagccaggat gatggccgc 29

<210> 211
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 211
tngagtgtgg aattgcgctt cttacctat

29

<210> 212
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 212
gngtgaagtc ttcctgggaa ccataatct

29

<210> 213
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 213
tnttcctga agagctggag aggtgctaa

29

<210> 214
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 214
gntctaccat gtgaagaagg aacgcaaaa 29

<210> 215
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 215
tnggcaaagc tgtgctgcag gatagagtg 29

<210> 216
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 216
angtgaagga ctgtaagccc acacaagct 29

<210> 217
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 217
tnatgacagc ccaggctgtg ttatctgca 29

<210> 218
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 218
cngggccttc gttttcacta ccttcagaa 29

<210> 219
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 219
tnaagacggt gactgagagg cagacaagg 29

<210> 220
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 220
tncctagtcc ctttaggtag ctctgcaga 29

<210> 221
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 221
gngcacgaga aagaaggcga ggatggcta 29

<210> 222
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 222
tnggacactg tggattctga gtcttggtg 29

<210> 223
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 223
ancaggaggc agtgggttct atagagtcc 29

<210> 224
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 224
cncttggtg cctgtttctc aattgcaga 29

<210> 225
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 225
cnctagctgt tcggctttac caaggagtt 29

<210> 226
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 226
ancacgctct gacgaatgag caacctctc

29

<210> 227
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 227
cnaacacaga ctggaagcct agagtcgct

29

<210> 228
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 228
gnaaaccagg aaatgtacac tgacacaat

29

<210> 229
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 229
tnatcaatac tgggtgacagt ctgctccaa 29

<210> 230
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 230
gncataaccc ttgacagaa gctgatctt 29

<210> 231
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 231
anatttaagc atttaacgtg aaggatagg 29

<210> 232
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 232
tnagagactg ggaaaccaac agccactgc 29

<210> 233
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 233
anggagcaag ttcctgagag ttcctgtt

29

<210> 234
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 234
anggagcaag ttcctgagag ttcctgtt

29

<210> 235
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 235
gnatggaggg agagagaaca caaattaac

29

<210> 236
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 236
gngacagtgt tgtgctgaag caggattcc

29

<210> 237
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 237
cntctccaac ttccacaatt tgctcacta

29

<210> 238
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 238
tnggaggaag agtggcagag gagcttata

29

<210> 239
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 239
cngtggtttg ttcttaggaa ggtagaagc

29

<210> 240
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 240
cngccagcca gcagcaaagc aaattaaga

29

<210> 241
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 241
tngaggaaag aagggtggag gtctgcaca

29

<210> 242
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 242
cggtgggatt tatttaacat gatcttggc

29

<210> 243
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 243
ccaatgggta tgatgatggt tcttcct

27

<210> 244
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 244
anggatgcca tccagatgag gccacagct

29

<210> 245
<211> 29
<212> DNA
<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 245
anctttcttct gcagctcatt caactcctg 29

<210> 246
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 246
gntcctccta gaccacaaag tagaaagca 29

<210> 247
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 247
gnctgacttt ctctccagtt tgtgaaatc 29

<210> 248
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 248
gngtggggtg aaaagggaat gattagga 29

<210> 249
<211> 29

<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 249
tnattcatgg gcagtcatt ggcgggctc 29

<210> 250
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 250
cngctgagac tcacacactg aatttcctt 29

<210> 251
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 251
cntcttggtc cggatgatgt ttttatcac 29

<210> 252
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 252

tntctcggtta tatggaggac gaatagact 29

<210> 253
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 253
ctctttatga gcatgatatg gcttcag 27

<210> 254
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 254
cntgctgccc gacttaaact gagaaccaa 29

<210> 255
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 255
gnngacaact cccatagttg aggtttgtc 29

<210> 256
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 256
tnggaagagt gacccagggg cccaaagca 29

<210> 257
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 257
cncgtgcaagc gagtgggtttt cagggctgt

29

<210> 258
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 258
tngagccaag gagagagcag acaggctga

29

<210> 259
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 259
ancccccaga agtacaagga catgccaag

29

<210> 260
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)

<223> biotinylated phosphoramidite residue

<400> 260

cnatagcatc agcatcaaca tggtgacaa

29

<210> 261

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 261

ctccattctc cttgactatc ttctccc

27

<210> 262

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 262

tncctggtgc catcatgatt gtatgaggc

29

<210> 263

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 263

anaggctcct ccattcctac agccatctt

29

<210> 264

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 264
cngtccaact gcttgtaggt tatagcaga 29

<210> 265
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 265
tnctctactt cacccttttc ggtgcatcg 29

<210> 266
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 266
tncataaaat gacccacac ctcctactg 29

<210> 267
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 267
gngaggtagc caggcatctc tgtgcattt 29

<210> 268
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 268
ancacagcag gaggtgagga ggaggagat 29

<210> 269
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 269
cnttgatctg agagcttttc accgtcaag 29

<210> 270
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 270
anctgccaaag aagaggagag tggtaaatt 29

<210> 271
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 271
angaaaaaga agtgatcatga tccagtaca 29

<210> 272
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 272
tnatacttgg acaggataga gcgaggttt 29

<210> 273
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 273
cncgactgcc ttaaaatgta aacctggat 29

<210> 274
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 274
tngcactggg aaagtgattg tgaggagta 29

<210> 275
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 275
tngtgctgcc aaagaacagg agcctcaaa 29

<210> 276
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 276
ctgctgctca tttgttcaat aatgtcaac

29

<210> 277
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 277
tngcactgtt gtcacgacgg cttggtctc

29

<210> 278
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 278
anaacagagc agtccagagg aagaacatc

29

<210> 279
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<220>
<221> misc_feature
<222> (2)
<223> biotinylated phosphoramidite residue

<400> 279
anatgaccag atcctctcag aagatttcc

29

<210> 280

<211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide

<220>
 <221> misc_feature
 <222> (2)
 <223> biotinylated phosphoramidite residue

<400> 280
 tnccagtgc tgggcatgca tgaaggcta

29

<210> 281
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide

<220>
 <221> misc_feature
 <222> (2)
 <223> biotinylated phosphoramidite residue

<400> 281
 ancagcaaac tcttcctttc gcttgctgc

29

<210> 282
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide

<220>
 <221> misc_feature
 <222> (2)
 <223> biotinylated phosphoramidite residue

<400> 282
 tntttctcac aggtcagttc atccacttt

29

<210> 283
 <211> 57
 <212> PRT
 <213> Homo sapiens

<400> 283
 Met Cys Arg Pro Phe Leu Ser Leu Leu Leu Cys Ser Thr Lys Cys Phe
 1 5 10 15

Leu Leu Cys Gly Leu Gly Gly Thr His Val Thr Phe Val Ser Cys Ser
 20 25 30

Lys Ala Gly Ala Thr Pro Ser Ser Leu Phe Ser Thr Gln His Gln Ala


35	40	45
Leu Ser Arg His Pro Ile Asn His Cys		
50	55	

Applicant's or agent's file reference 1290.1001010	International application No. PCT/US 00/25135
--	---

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on pages <u>333</u> to <u>335</u> , line <u>35</u> to <u>36</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) American Type Culture Collection (ATCC) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit see Attachment A	Accession Number see Attachment A
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input checked="" type="checkbox"/>	
In respect of those designations for which a European patent is sought, the Applicant(s) hereby informs the International Bureau that the Applicant wishes that, until the publication of the mention of the grant of a European patent or for 20 years from the date of filing if the application is refused or withdrawn or deemed to be withdrawn, the biological material deposited with the American Type Culture Collection under Accession No. <u>see Attachment A</u>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input checked="" type="checkbox"/> For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer 	<input type="checkbox"/> For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer
--	--

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM
(Additional Sheet)

C. ADDITIONAL INDICATIONS (Continued)

shall be made available as provided in ~~Rule 28(3) EPC only~~ by the issue of a sample to an expert nominated by the requester (Rule 28(4) EPC).

In respect of the designation of Australia in the subject PCT application, and in accordance with Regulation 3.25(3) of the Australian Patents Regulations, the Applicant hereby gives notice that the furnishing of a sample of the biological material deposited with the American Type Culture Collection under Accession No. ~~Attachment A~~ shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention and who is nominated in a request for the furnishing of a sample.

In respect of the designation of Canada in the subject PCT application, the Applicant hereby informs the International Bureau that the Applicant wishes that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the biological material deposited with the American Type Culture Collection under Accession No. ~~Attachment A~~ and referred to in the application to an independent expert nominated by the Commissioner.

Attachment A

-1-

Deposit of Clones

Clones AX65_22, BD335_14, BG241_1, BL187_4, BL249_18, BO71_1, BO365_2, BV51_1, BV140_3, BV141_2, CC194_4, and DA136_11 were deposited on October 3, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98196, from which each clone comprising a particular polynucleotide is obtainable.

Clones AR415_4, AS63_29, BG160_1, BO432_4, BO538_2, BR595_4, CI490_2, CI522_1, CN238_1, CO390_1, and AY304_1 (an additional isolate of clone AY304_14) were deposited on October 25, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98232, from which each clone comprising a particular polynucleotide is obtainable. Clone AY304_14 was deposited on October 23, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 98561.

Clones AJ20_2, AR440_1, AS164_1, AX8_1, BD176_3, BD339_1, BD427_1, BL229_22, BV123_16, and CH377_1 were deposited on November 15, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98261, from which each clone comprising a particular polynucleotide is obtainable.

Clones BD441_1, BD441_2, BG102_3, BK158_1, BP163_1, BZ16_3, CC182_1, CG109_1 and CJ397_1 were deposited on November 20, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98264, from which each clone comprising a particular polynucleotide is obtainable.

Clones AM795_4, AT340_1, BG132_1, BG219_2, BG366_2, BV172_2, CC247_10, CI480_9, CO722_1, and CT748_2 were deposited on December 5, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98271, from which each clone comprising a particular polynucleotide is obtainable.

Clones AJ1_1, AQ73_3, BG142_1, BV66_1, BV291_3, CK201_1, CQ331_2, CT550_1, CT585_1 and CT797_3 were deposited on December 13, 1996 with the ATCC

Attachment A

-2-

(American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98278, from which each clone comprising a particular polynucleotide is obtainable.

Clones CB107_1, CG300_3, CJ145_1, CJ160_11, CO20_1, CO223_1, CO310_2, CP258_3, CW1155_3 and CZ247_2 were deposited on December 17, 1996 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98279, from which each clone comprising a particular polynucleotide is obtainable. Clone CO223_3 was deposited on January 9, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number 98291.

Clones AM666_1, BN387_3, BQ135_2, CR678_1, CW420_2, CW795_2, CW823_3, DF989_3, DL162_2, DL162_1, and EC172_1 were deposited on January 10, 1997 with the ATCC (American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number 98292, from which each clone comprising a particular polynucleotide is obtainable.

INTERNATIONAL SEARCH REPORT

International Application No

I US 00/25135

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N1/21 C12N5/10 C12Q1/68
A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, STRAND

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, L	WO 98 17687 A (GENETICS INST) 30 April 1998 (1998-04-30) the document throws doubt on the priority of the application abstract; claims 20-22 see SEQ ID NO: 8 and 9 (pp.73-77) page 18, line 30 -page 20, line 2 page 23, line 12 -page 24, line 14 page 31, line 12 -page 64, line 16 -----	1-11

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 December 2000

Date of mailing of the international search report

30.01.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US 00/25135

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-11

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11

An isolated polynucleotide comprising SEQ ID NO: 41 which encodes a protein of SEQ ID NO: 42 (BG160_1). A host cell, a process for producing said protein, a protein produced by said process, a composition comprising said protein.

2. Claims: 12, 13

An isolated polynucleotide comprising SEQ ID NO: 129. A protein encoded by said polynucleotide having amino acid sequence SEQ ID NO: 130 (C0722_1).

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

F US 00/25135

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9817687 A	30-04-1998	AU 5004097 A EP 0960199 A	15-05-1998 01-12-1999
